



A Report on Non-Ionizing Radiation

Colorectal Cancer Soaring in Young Adults; Are Smartphones in the Mix?

Epidemiologist De-Kun Li Wants To Know

[Facebook](#) [Twitter](#) [Print](#) [Email](#)

June 3, 2019
Last updated July 31, 2019

De-Kun Li wants to change the conversation on cell phones and cancer. Li, a senior epidemiologist and veteran EMF researcher, believes that brain tumors have been getting too much attention at the expense of other types of cancer, notably colorectal cancer.

Efforts to reduce colon and rectal cancers have been a striking success story for those over 50 years old. Incidence among older Americans declined 32% between 2000 and 2013, due largely to better screening. But the story for young adults is very different. Those born around 1990 now face *four times* the risk of developing rectal cancer and *twice* the risk of colon cancer in their 20s, compared to those born around 1950, according to the [American Cancer Society](#). (Rates among young adults are still relatively low; see graphs below.)


Microwave News The Web

[Return to Main Article Archive](#)

VIEW ARTICLES BY YEAR

- [2020 Articles](#)
- [2019 Articles](#)
- [2018 Articles](#)
- [2017 Articles](#)
- [2016 Articles](#)
- [2015 Articles](#)
- [2014 Articles](#)
- [2013 Articles](#)
- [2012 Articles](#)
- [2011 Articles](#)
- [2010 Articles](#)
- [2009 Articles](#)
- [2008 Articles](#)
- [2007 Articles](#)
- [2006 Articles](#)
- [2005 Articles](#)
- [2004 Articles](#)

Help Support
Microwave News



U.S. Trends in Age-Specific Colon and Rectal Cancer Incidence Rates
Source: R.L. Siegel, et al., *JNCI*, Vol.109, 2017 — Supplementary Figure 4

According to the National Cancer Institute's most recent [annual report](#), released last week, colorectal cancer is the most common cancer among men between the age of 20 and 49.

"No one can explain this apparent contradiction," Li told *Microwave News*. Known risk factors for colorectal cancer include obesity, an unhealthy diet and lack of physical activity, but Li doesn't think they can resolve the paradox. "If anything," he said, "younger generations are more health conscious with better diets."

Li offers an alternative hypothesis: Young people's habit of carrying their cell phones in the front or back pockets of their jeans. "When placed in trouser pockets, the phones are in the vicinity of the rectum and the distal colon and these are the sites of the largest increases in cancer," he says.

RF Exposures from Phones on Standby

Li's hypothesis assumes that phones continue to emit RF radiation when placed in a pocket. They do, especially smartphones, but exposures are difficult to estimate.

When not in use, phones quickly go to standby mode, and RF transmissions are greatly reduced. Yet, the phone stays active because the system must always know its location in order for the network to direct calls or texts to the nearest cell tower for delivery.

A Swedish-Danish [study](#), published in 2012, found that phones on standby were only in touch with the local base station once every two-to-five hours —and then for just a few seconds. "The exposure in standby mode can be considered negligible," concluded [Kjell Hansson Mild](#) of Umeå University, the lead author of the study. [Jørgen Bach Andersen](#) of Aalborg University was a coauthor.

Today, Mild sees things differently because smartphones —which were still relatively new when he was working on his study— contact the network much more frequently, if only to update their location. Mild wants research. "We need more measurements and dosimetry on how a phone radiates in a pocket," Mild said in an interview, adding that, "I keep my phone in my briefcase and when I turn it on, I use a hands-free set."

When a phone is in a pocket, the separation distance from the body is less than that recommended by [Om Gandhi](#), Mild said. Gandhi, a professor emeritus at the University of Utah, has published a stream of papers on cell phone exposures, estimated by measurements and calculations (here's [one](#) from 2002).

In an e-mail exchange, Gandhi noted that, in compliance tests, cell phones are placed about 10 mm away from the body. Without that spacing —for instance, when a phone is in a pocket next to the body — exposure (SAR) would be about twice that cited in safety specifications. The title of Gandhi's most recent [paper](#), posted in April by the IEEE, the electrical engineering society, spells out the problem: "Microwave Emissions from Cell Phones Exceed Safety Limits in Europe and the U.S. When Touching the Body."

In his [paper](#), Mild highlighted that exposures go up if a phone on standby is in motion:

"[W]hen a person carries the phone in a pocket and moves in a city, the phone updates when entering a new

base station area, and this can be as often as once every few minutes depending if one is in [a] city center or suburban area. For [those] periodic updates the GSM phones operate at full power.”

A later [investigation](#) by Damiano Urbinello and [Martin Rösli](#) of the University of Basel showed that exposures are even more significant when a phone is in a moving vehicle, whether a car, train or bus.¹ They predicted that smartphones would likely increase exposures because of their need for “regular location updates.”

There was much more to learn. Urbinello and Rösli wrote:

“There is an urgent need to evaluate more thoroughly how personal RF-EMF exposure is affected by one’s own phone in standby mode.”

That was six years ago. There’s been little, if any, follow-up since then. “I am not aware of much further research,” Rösli told *Microwave News*.

Exposures are “quite variable and the amount of uplink may depend on the type of mobile phone, the network, the provider and the apps installed,” Rösli said. “In particular, push functions are expected to increase the amount of uplink in standby mode.”

Phones in Pockets Below the Waist

Do young adults keep their phones in their pockets? Li points to a 2017 [survey](#) of Australian women, aged 15 to 40, by [Mary Redmayne](#), who found that a majority routinely put their phones in a pocket below the waist. Importantly, the youngest women do this the most. She also reported that 96% of the women keep their phones on standby during the day; at night it’s 83%.

In an e-mail, Redmayne said that she has no information about where young men keep their phones, nor is she aware of anyone who had looked into it.

Colorectal Rates Among the Young Rising in Many Countries

Increases of colorectal cancer among the young are being reported around the world, not just in the U.S. — including many European countries, as well as Australia and New Zealand. Just a few weeks ago, [IARC](#) reported that, between 2008 and 2015, the incidence of colon cancer among 20-29-year-olds increased by about 18% per year in Denmark, 8% in Australia and 4% in Ireland. While the risk declined in Canada and Norway, the rate of rectal cancer in those countries rose by 3.5% and 10.6% per year, respectively.

Writing in *The Lancet – Gastroenterology & Hepatology*, the IARC team advises:

“Although the incidence of colorectal cancer in adults younger than 50 years remains low compared with that in older age groups, our findings are of concern and highlight the need for action to counteract the rising burden of disease in younger cohorts.”

Also in May, a second research group reported similar trends across 20 European countries in *Gut*, a *BMJ* journal. On average, colorectal cancer rose by 7.9% per year among 20-29-year-olds from 2004 to 2016. The effect gets smaller with age: The increase among 30-39-

year-olds was 4.9%, and 1.6% for 40-49-year-olds.

In 1990, the rate was 0.8 cases per 100,000 Europeans in their 20s. By 2016, it was 2.3/100,000. The big jump came recently: For 1990-2004, incidence went up by 1.7% per year, but it grew to 7.9% per year from 2004 to 2016.

Li: A Plausible Explanation

“My hypothesis could be proven wrong,” Li concedes, “but this potential explanation is more plausible than any other current hypotheses out there.” People should be made aware of the possible risks, he said, so that young adults can decide whether to take precautions and colorectal cancer researchers can weigh the merits of the case.


Over the last 20 years, Li, who is with Kaiser Permanente in Oakland, CA, has completed a number of groundbreaking studies on the effects of power-frequency EMFs. In 2001, he reported that women faced up to a sixfold increased risk of miscarriages when exposed to magnetic fields above 16 mG (see *MWN*, *M/J01* and *J/F02*). That early work has been supported by others and there are now at least **seven studies** that point to an EMF–miscarriage link.

In other projects, Li has shown that magnetic fields can affect the quality of **human sperm** and that prenatal EMF exposure can lead to childhood **asthma** and **obesity**.

July 31, 2019

Similar Trends in Canada

New data, released today, show the same trend among men and women in Canada. Colorectal cancer rates are going down among those over 50 years old and up among those under 50. See the **graphs** below:



These findings appear in *JAMA Open Network*. The **paper** is open access.

1. Another finding: Standby exposures from UMTS (3G) phones were

"considerably lower" than those from GSM (2G) phones.

*De-Kun Li, Kjell Hansson Mild, Jørgen Bach Andersen,
Om Gandhi, Martin Röösli, IARC, colorectal cancer,
cell phones cancer, brain cell phones, standby mode, smartphone,*

© Copyright Microwave News 2003-2020. All Rights Reserved.