

COMMENTS OF BELLSOUTH

CC DOCKET NO. 01-318

JANUARY 22, 2002

ATTACHMENT 2, PART 3

# APPENDIX F

Statistician's Report  
(Louisiana Collaborative)



BellSouth Telecommunications, Inc. 504 528-2050  
Suite 3060 Fax 504 528-2948  
365 Canal Street  
New Orleans, Louisiana 70130-1102

Victoria K. McHenry  
General Counsel - LA

February 29, 2000

**VIA FEDERAL EXPRESS**

Ms. Susan Cowart  
Louisiana Public Service Commission  
Suite 1630  
One American Place  
Baton Rouge, LA 70825

RE: LPSC Docket No. U-22252-C  
Louisiana Public Service Commission, ex parte  
In re: BellSouth Telecommunications, Inc.  
Service Quality Performance Measurements


Dear Ms. Cowart:

Enclosed are the original and one (1) copy each of the following documents to be filed into the record of the referenced matter:

1. Updated BellSouth SQM Report
2. Updated Statistician's Report

These items were not specifically included in the Commission's most recently issued Notice so I am unsure when they are due. In any event, we are providing them as soon as possible.

Sincerely,

  
Victoria K. McHenry

VKM/as  
Encs.

cc: Official Service List (w/enc.)(via email)

DRAFT

**Statistical Techniques  
For The Analysis And Comparison Of  
Performance Measurement Data**

Submitted to Louisiana Public Service Commission (LPSC)  
Docket U-22252 Subdocket C

Revised February 28, 2000

**Introduction and Scope**

The Louisiana Public Service Commission (LPSC) staff has requested Drs. S. Hinkins, E. Mulrow, and F. Scheuren<sup>1</sup> of Ernst & Young LLP (consultants for BellSouth Telecommunications), and Dr. C. Mallows of AT&T Labs-Research to set out their views on the application of a statistical analysis to performance measurement data. The present report is intended to provide a detailed statistical report on appropriate methodology.

The setting for the analysis is crucial to the interpretation of any statistical significance that might be found. There is no doubt that, to quote the Commission staff, "statistical analysis can help reveal the likelihood that reported differences in an ILECs performance toward its retail customers and CLECs are due to underlying differences in behavior rather than random chance" (Staff Final Recommendation, LPSC Docket No. U-22252 - Subdocket C, dated August 12, 1998, pages 15 - 16).

To frame our presentation the next paragraph from the LPSC Docket U-22252 is quoted in its entirety.

**"Statistical tests are effective in identifying those measurements where differences in performance exist. The tests themselves cannot identify the cause of the apparent differences. The differences may be due to a variety of reasons, including: 1) when the ILEC and CLEC processes being measured are actually different and should not be expected to produce the same result, 2) when the ILEC is employing discriminatory practices, or 3) when assumptions necessary for the statistical test to be valid are not being met." (Ibid., page 16)**

Apparent statistically significant differences in BellSouth and CLEC performance can arise when

- the ILEC and CLEC processes being measured are actually different and should not be expected to produce the same result
- the ILEC is employing discriminatory practices, or
- assumptions necessary for the statistical test to be valid are not being met.

---

<sup>1</sup> Dr. Scheuren is now a Senior Fellow at the Urban Institute.

RECEIVED

MAR 01 2000

LOUISIANA PUBLIC SERVICE COMMISSION  
ADMINISTRATIVE HEARINGS DIVISION

00 APR 1 AM 10 39

**DRAFT**

To meet the Louisiana Commission's purpose, we will recommend techniques that are robust in the presence of possible assumption failure, carefully examine BellSouth Telecommunications (BST) and CLEC performance so "like" is compared only to "like," and are still able, in a highly efficient manner, to detect differences. Upon investigation any differences detected might lead to concerns about possible discriminatory practices.

The LPSC staff also states "that a uniform methodology which identifies those items which need to be measured, how they are to be measured, and how the results are to be reported is also desirable and would be beneficial to all parties" (Ibid., page 16). We agree with this goal as well, stipulating only that the use of a single method may not be desirable while a single methodology (or a set of methods) could be.

The statistical process for testing if CLEC and ILEC customers are being treated equally involves more than just a mathematical formula. Three key elements need to be considered before an appropriate decision process can be developed. These are

- the type of data,
- the type of comparison, and
- the type of performance measure.

When examining the various combinations of these elements, we find that there is a set of testing principles that can be applied uniformly. However, the statistical formulae that need to be used change as the situation changes.

To be responsive to the Commission, we have divided our discussion into four sections and five appendices. The contents of each of these are briefly mentioned below -- first for the main report and then for the extensive supporting appendix materials.

For the main report, this section (Section I) introduces our work and sets out the required scope. The next two sections (Sections II and III) discuss the type of comparisons that need to be identified, and the appropriate testing principles. The final section (Section IV) provides an overview of appropriate testing methodologies, based on what we have learned from our examination of BellSouth's performance measure data in Louisiana.

The five appendices provide technical details on the statistical calculations involved in the Truncated Z statistic (Appendix A), the implementation of the methodology for the trunk blocking performance measure (Appendix B), the calculations involved in computing the balancing critical value of a test (Appendix C), examples of ways to present the results using detailed statistical displays so that results can be audited (Appendix D), and the technical details involved in data trimming (Appendix E).

**DRAFT****2. Data Considerations, Comparisons, and Measurement Types**

This section makes general distinctions which apply to the performance measures. These distinctions will be important in the determination of appropriate methodologies.

**Data Set Types.** The type of statistical methodology used depends on the form of the data available. In general, there are two ways to classify the data used for performance measure comparisons. These are:

- transaction level data, and
- aggregated summaries.

Records in a transaction level data set represent a single transaction, e.g. an individual customer order, or the record of a specific trouble reported by a customer. This type of data set allows for deep like-to-like comparisons, and may also allow one to identify the root cause of a problem. A testing methodology needs to be carefully chosen so that it incorporates the comparison levels and does not cover up problem areas.

Records in an aggregated summary data set are typically summaries of related transactions. For example, the total number of blocked calls in a trunk group during the noon hour of a day is a summary statistic. This type of data set may not contain as much information as a transaction level data set, and it therefore needs to be treated differently. While a general methodology may be determined for a transaction level data set, it may not be possible to do so for aggregated summaries. Testing methodology needs to be developed on a case-by-case basis.

**Comparison Types.** An ILEC's performance in providing services to CLEC customers is tested in one of two ways:

- by comparing CLEC performance to ILEC performance when a retail analog exists, or
- by comparing CLEC performance to a benchmark.

The testing methodologies for these two situations will have similarities, but there are differences that need to be understood.

Table 1 categorizes those performance measures that E&Y has examined by data type and comparison type. The table shows that five performance measures with retail analogs have transaction level data, while three others with retail analogs only have summary level data. No performance measures using benchmarks have been studied.

**DRAFT**

**Table 1. Classification of Performance Measures by Data and Comparison Type  
(only measures previously examined by E&Y are included)**

Level of Data	Comparison Type	
	Retail Analog	Benchmark
Transaction Level	Order Completion Interval	No Measures Examined
	Maintenance Average Duration	
	% Missed Installations	
	% Missed Repair	
	Trouble Report Rate	
Summary Level	Billing Timeliness	No Measures Examined
	OSS Response Interval	
	Trunk Blocking	

**Measurement Types.** The performance measures that will undergo testing are of four types: means, proportions (an average of a measure that takes on only the values of 0 or 1), rates, and ratios.

While all four have similar characteristics, proportions and rates are derived from count data while means and ratios are derived from interval measurements. Table 2 classifies the performance measures by the type of measurement.

**Table 2: Classification of Performance Measures by Measurement Type**

Mean	Proportion	Rate	Ratio
Order Completion Interval	Percent Missed Installations	Trouble Report Rate	Billing Accuracy
Maint. Ave. Duration	Percent Missed Repairs		
OSS Response Interval	Billing Timeliness		
	Trunk Blocking		

### 3. Testing Principles

This section describes five general principles which the final methodology should satisfy:

**DRAFT**

1. *When possible, data should be compared at appropriate levels, e.g. wire center, time of month, dispatched, residential, new orders.*
2. *Each performance measure of interest should be summarized by one overall test statistic giving the decision maker a rule that determines whether a statistically significant difference exists.*
3. *The decision system must be developed so that it does not require intermediate manual intervention.*
4. *The testing methodology should balance Type I and Type II Error probabilities.*
5. *Trimming of extreme observations from BellSouth and CLEC distributions is needed in order to ensure that a fair comparison is made between performance measures.*

Like-to-Like Comparisons. *When possible, data should be compared at appropriate levels, e.g. wire center, time of month, dispatched, residential, new orders.*

In particular, to meet this goal the testing process should:

- Identify variables that may affect the performance measure.
- Record important confounding covariates.
- Adjust for the observed covariates in order to remove potential biases and to make the CLEC and the ILEC units as comparable as possible.

It is a well known principle that comparisons should be made on equal footing: apples-to-apples, oranges-to-oranges. Statistical techniques that are addressed in most text books usually assume that this is the case beforehand. Some higher level books address the issue of “designed experiments” and discuss appropriate ways to structure the data collection method so that the text books’ formulae can be used in analyzing the data.

Performance measure testing does not involve data from a designed experiment. Rather, the data is obtained from an observational study. That being the case, one must impose a structure on the data after it is gathered in order to assure that fair comparisons are being made. For example, it is important to disaggregate the data to a fine level so that appropriate like-to-like comparisons of CLEC and ILEC data can be made. Any statistical methodology that ignores important confounding variables can produce biased results.

Aggregate Level Test Statistic. *Each performance measure of interest should be summarized by one overall test statistic giving the decision maker a rule that determines whether a statistically significant difference exists.*

To achieve this goal, the aggregate test statistic should have the following properties:

**DRAFT**

- The method should provide a single overall index, on a standard scale.
- If entries in comparison cells are exactly proportional over a covariate, the aggregated index should be very nearly the same as if comparisons on the covariate had not been done.
- The contribution of each comparison cell should depend on the number of observations in the cell.
- Cancellation between comparison cells should be limited, i.e., positive outcomes should not be allowed to cancel negative ones.
- The index should be a continuous function of the observations.

Since the data are being disaggregated to a very deep level, thousands of like-to-like comparison cells are created. An aggregate summary statistic is needed in order to make an overall judgment.

The aggregate level statistic should be insensitive to small changes in cells values, and its value should not be affected if some of the disaggregation for like-to-like cells is truly unnecessary. Furthermore, individual cell results should be weighted so that those cells with more transactions have larger effects on the overall result.

**Production Mode Process.** *The decision system must be developed so that it does not require intermediate manual intervention.*

Two statistical paradigms are possible for examining performance measure data. In the exploratory paradigm, data are examined and methodology is developed that is consistent with what is found. In a production paradigm a methodology is decided upon before data exploration. For the production paradigm to succeed

- Calculations should be well defined for possible eventualities.
- The decision process should be based on an algorithm that needs no manual intervention.
- Results should be arrived at in a timely manner.
- The system must recognize that resources are needed for other performance measure-related processes that also must be run in a timely manner.
- The system should be both auditable and adjustable over time.

While the exploratory paradigm provides protection against using erroneous data, it requires a great deal of lead time and is unsuitable for timely monthly performance measure testing. A production paradigm will not only promptly produce overall test results but will also provide documentation that can be used to explore the data after the test results are released.

**DRAFT**

**Error Probability Balancing.** *The testing methodology should balance Type I and Type II Error probabilities.*

Specifically, what is required to achieve this goal is

- The probability of a Type I error should equal the probability of a Type II error for well-defined null and alternative hypotheses.
- The formula for a test's balancing critical value should be simple enough to calculate using standard mathematical functions, i.e. one should avoid methods that require computationally intensive techniques.
- Little to no information beyond the null hypothesis, the alternative hypothesis, and the number of observations should be required for calculating the balancing critical value.

The objective of a statistical test is to test a hypothesis concerning the values of one or more population parameters. Usually an inquiry into whether or not there is evidence to support a hypothesis, called the *alternative hypothesis*, is conducted by seeking statistical evidence that the converse of the alternative, the *null hypothesis*, is most likely false. If there is not sufficient evidence to reject the null hypothesis, then a case for accepting the alternative has not been made.

Two types of errors are possible in any decision-making process. These have been summarized in Table 3.

**Table 3: Statistical Testing Errors**

<b>Decision Error</b>	<b>General Description</b>	<b>In terms of Performance Measure Testing</b>
Type I	Rejecting the null hypothesis (accepting the alternative) when the null is true.	Deciding that BST favors its own customers when it does not.
Type II	Accepting the null hypothesis when the alternative is true.	Deciding that BST does not favor its own customers when it does.

In a controlled experimental study where the sample sizes are relatively small, it is generally desirable to control the Type I error closely to avoid making a conclusion that there is a difference when, in fact, there is none. The probability of a Type II error is not directly controlled but is determined by the sample size and the distance between the null and the alternative hypotheses.

**DRAFT**

If a standard of materiality is set by stating a specific alternative for the test, and the distribution of the test statistic under both the null and alternative hypotheses is understood, then a critical value can be determined so that the two error probabilities are equal.

Trimming. *Trimming of extreme observations from BellSouth and CLEC distributions is needed in order to ensure that a fair comparison is made between performance measures.*

Three conditions are needed to accomplish this goal. These are:

- Trimming should be based on a general rule that can be used in a production setting.
- Trimmed observations should not simply be discarded; they need to be examined and possibly used in the final decision making process.
- Trimming should only be used on performance measures that are sensitive to "outliers."

For the purpose of performance measure testing, trimming refers to removing transactions that significantly distort the performance measure statistic for the set of transactions under consideration. For example, the arithmetic average (or mean) is extremely sensitive to "outliers" since a single large value can significantly distort the average.

The term "outliers" refers to:

- 1) extreme data values that may be valid, but since they are rare measurements, they may be considered to be statistically unique; or
- 2) large values that should not be in the analysis data set because of errors in the measurement or in selecting the data.

Trimming is beneficial since it puts both ILEC and CLEC transactions on equal footing with respect to the largest value in each set. Note, though, that it is only needed for performance measures that are distorted by outliers. Of the three types of measures defined in Section 2, only mean (average) measures require trimming. Appendix E sets forth a trimming plan for mean performance measures.

#### **4. Testing Methodology**

This section details the testing methodology that is most appropriate for the various types of performance measures. First, transaction level testing will be discussed when there is a retail analog. Next, transaction level testing against a benchmark. Then, testing when only aggregated summaries are available.

Transaction Level - Retail Analog: The Truncated Z Statistic. When a retail analog is available CLEC performance can be directly compared with ILEC performance. Over

**DRAFT**

the last year. for transaction level data, many test statistics have been examined. We now believe that the "Truncated Z" test statistic provides the best compromise with respect to possessing the desired qualities outlined in Section 3, above.

The Truncated Z is fully described in Appendix A, and formulae for calculation of a balancing critical value are found in Appendix C. The main features of this statistic are:

- A basic test statistic is calculated within each comparison cell.
- The value of a cell's result is left "as is" if the result suggests that "favoritism" may be taking place. Otherwise, the result is set to zero. This is called the truncation step.
- Weights that depend on the volume of both ILEC and CLEC transactions within the cell are determined, and a weighted sum of the "truncated" cell results is calculated.
- The weighted sum is theoretically corrected to account for the truncation, and a final overall statistic is determined.
- This overall test value is compared to a balancing critical value to determine if favoritism is likely.

The test statistic itself is based on like-to-like comparisons, and it possesses all five of the properties of an aggregate test statistic (Section 3). While the test requires a large amount of calculations, our studies of the process on some of BellSouth's performance measure data indicate that the calculations can be completed in a reasonable amount of time. Therefore, the process can be put into production mode. Finally, since a balancing critical value can be calculated, it is possible to balance the error probabilities.

Transaction Level - Benchmark. When a benchmark is used, CLEC performance is not compared with ILEC performance. Like-to-like comparison cells are not needed, thus greatly simplifying the testing process. Statistical testing can be done using a probability model, or non-statistical testing can be done using a deterministic model. No data for this data/comparison class has been studied at this point in time.

Aggregated Summary - Retail Analog or Benchmark. We cannot provide any one single set of rules for the analysis of data in this class. Data that is an aggregated summary of transactions may or may not present problems. For example, BellSouth's trunk blocking data is saved as summaries by hour of the day. Collectively, the summaries do provide sufficient information to proceed with the Truncated Z methodology.

On the other hand, our examination of the data for the OSS response interval revealed that information necessary for computing a Truncated Z was not available. In this case, however, we were able to construct a satisfactory time series method to analyze the measure.

**DRAFT**

Each measure falling into this class needs to be handled on a case-by-case basis. If sufficient information is available to use the Truncated Z method, then we feel it should be used. When the Truncated Z cannot be used, a testing methodology that adheres closely to the principles outlined in Section 3 should be determined and followed.

## Appendix A. The Truncated Z Statistic

The Truncated Z test statistic was developed by Dr. Mallows in order to have an aggregate level test when transaction level data are available that

- provides a single overall index on a standard scale;
- will not change the outcome if the disaggregation is unnecessary,
- incorporates the number of observations in a cell into the determination of the weight for the contribution of each comparison cell,
- limits the amount of “neutralization” between comparison cells, and
- is a continuous function of the observations.

The Ernst & Young statistical team and Dr. Mallows have studied the implementation of the statistic using some of BellSouth’s performance measure data. This has resulted in an overall process for comparing CLEC and ILEC performance such that the following principles hold:

- 1) Like-to-Like Comparisons are made. (See Appendix B for an example based on the trunk blocking measure.)
- 2) Error probabilities are balanced. (See Appendix C)
- 3) Extreme values are trimmed from the data sets when they significantly distort the performance measure statistic. (See Appendix E)
- 4) The testing process is an automated production system. (Discussed here. See Appendix D for reporting guidelines.)
- 5) The determination of ILEC favoritism is based on a single aggregate level test statistic. (Discussed here.)

This appendix provides the details behind computing the Truncated Z test statistic so that principles 4 and 5 hold. We start by assuming that any necessary trimming of the data is complete, and that the data are disaggregated so that comparisons are made within appropriate classes or adjustment cells that define “like” observations.

### Notation and Exact Testing Distributions

Below, we have detailed the basic notation for the construction of the truncated z statistic. In what follows the word “cell” should be taken to mean a like-to-like comparison cell that has both one (or more) ILEC observation and one (or more) CLEC observation.

- $L$  = the total number of occupied cells
- $j$  =  $1, \dots, L$ : an index for the cells
- $n_{1j}$  = the number of ILEC transactions in cell  $j$
- $n_{2j}$  = the number of CLEC transactions in cell  $j$
- $n_j$  = the total number transactions in cell  $j$ ;  $n_j = n_{1j} + n_{2j}$

$$\begin{aligned}
X_{1jk} &= \text{individual ILEC transactions in cell } j; k = 1, \dots, n_{1j} \\
X_{2jk} &= \text{individual CLEC transactions in cell } j; k = 1, \dots, n_{2j} \\
Y_{jk} &= \text{individual transaction (both ILEC and CLEC) in cell } j \\
&= \begin{cases} X_{1jk} & k = 1, \dots, n_{1j} \\ X_{2jk} & k = n_{1j} + 1, \dots, n_j \end{cases}
\end{aligned}$$

$\Phi^{-1}(\cdot)$  = the inverse of the cumulative standard normal distribution function

For Mean Performance Measures the following additional notation is needed.

$$\begin{aligned}
\bar{X}_{1j} &= \text{the ILEC sample mean of cell } j \\
\bar{X}_{2j} &= \text{the CLEC sample mean of cell } j \\
s_{1j}^2 &= \text{the ILEC sample variance in cell } j \\
s_{2j}^2 &= \text{the CLEC sample variance in cell } j \\
\{y_{jk}\} &= \text{a random sample of size } n_j \text{ from the set of } Y_{j1}, \dots, Y_{jn_j}; k = 1, \dots, n_j \\
M_j &= \text{the total number of distinct pairs of samples of size } n_{1j} \text{ and } n_{2j}; \\
&= \binom{n_j}{n_{1j}}
\end{aligned}$$

The exact parity test is the permutation test based on the "modified Z" statistic. For large samples, we can avoid permutation calculations since this statistic will be normal (or Student's t) to a good approximation. For small samples, where we cannot avoid permutation calculations, we have found that the difference between "modified Z" and the textbook "pooled Z" is negligible. We therefore propose to use the permutation test based on pooled Z for small samples. This decision speeds up the permutation computations considerably, because for each permutation we need only compute the sum of the CLEC sample values, and not the pooled statistic itself.

A permutation probability mass function distribution for cell j, based on the "pooled Z" can be written as

$$PM(t) = P\left(\sum_k y_{jk} = t\right) = \frac{\text{the number of samples that sum to } t}{M_j},$$

and the corresponding cumulative permutation distribution is

$$CPM(t) = P\left(\sum_k y_{jk} \leq t\right) = \frac{\text{the number of samples with sum} \leq t}{M_j}$$

For Proportion Performance Measures the following notation is defined

- $a_{1j}$  = the number of ILEC cases possessing an attribute of interest in cell j
- $a_{2j}$  = the number of CLEC cases possessing an attribute of interest in cell j
- $a_j$  = the number of cases possessing an attribute of interest in cell j;  $a_{1j} + a_{2j}$

The exact distribution for a parity test is the hypergeometric distribution. The hypergeometric probability mass function distribution for cell j is

$$HG(h) = P(H = h) = \begin{cases} \frac{\binom{n_{1j}}{h} \binom{n_{2j}}{a_j - h}}{\binom{n_j}{a_j}}, & \max(0, a_j - n_{2j}) \leq h \leq \min(a_j, n_{1j}) \\ 0 & \text{otherwise} \end{cases}$$

and the cumulative hypergeometric distribution is

$$CHG(x) = P(H \leq x) = \begin{cases} 0 & x < \max(0, a_j - n_{2j}) \\ \sum_{h=\max(0, a_j - n_{2j})}^x HG(h), & \max(0, a_j - n_{2j}) \leq x \leq \min(a_j, n_{1j}) \\ 1 & x > \min(a_j, n_{1j}) \end{cases}$$

For Rate Measures, the notation needed is defined as

- $b_{1j}$  = the number of ILEC base elements in cell j
- $b_{2j}$  = the number of CLEC base elements in cell j
- $b_j$  = the total number of base elements in cell j;  $b_{1j} + b_{2j}$
- $\bar{p}_{1j}$  = the ILEC sample rate of cell j;  $n_{1j}/b_{1j}$
- $\bar{p}_{2j}$  = the CLEC sample rate of cell j;  $n_{2j}/b_{2j}$
- $q_j$  = the relative proportion of ILEC elements for cell j;  $b_{1j}/b_j$

The exact distribution for a parity test is the binomial distribution. The binomial probability mass function distribution for cell j is

$$BN(\cdot) = P(B = k) = \begin{cases} \binom{n_j}{k} q_j^k (1 - q_j)^{n_j - k}, & 0 \leq k \leq n_j, \\ 0 & \text{otherwise} \end{cases}$$

and the cumulative binomial distribution is

$$CBN(x) = P(B \leq x) = \begin{cases} 0 & x < 0 \\ \sum_{k=0}^x BN(k), & 0 \leq x \leq n_j. \\ 1 & x > n_j \end{cases}$$

For Ratio Performance Measures the following additional notation is needed.

- $U_{1jk}$  = additional quantity of interest of an individual ILEC transaction in cell j;  $k = 1, \dots, n_{1j}$
- $U_{2jk}$  = additional quantity of interest of an individual CLEC transaction in cell j;  $k = 1, \dots, n_{2j}$
- $\hat{R}_{ij}$  = the ILEC ( $i = 1$ ) or CLEC ( $i = 2$ ) ratio of the total additional quantity of interest to the base transaction total in cell j, i.e.,  $\sum_k U_{ijk} / \sum_k X_{ijk}$

### Calculating the Truncated Z

The general methodology for calculating an aggregate level test statistic is outlined below.

1. **Calculate cell weights,  $W_j$ .** A weight based on the number of transactions is used so that a cell which has a larger number of transactions has a larger weight. The actual weight formulae will depend on the type of measure.

*Mean or Ratio Measure*

$$W_j = \sqrt{\frac{n_{1j} n_{2j}}{n_j}}$$

*Proportion Measure*

$$W_j = \sqrt{\frac{n_{2j} n_{1j}}{n_j} \cdot \frac{a_j}{n_j} \cdot \left(1 - \frac{a_j}{n_j}\right)}$$

*Rate Measure:*

$$W_j = \sqrt{\frac{b_{1j} b_{2j} \cdot n_j}{b_j \cdot b_j}}$$

2. **In each cell, calculate a Z value,  $Z_j$ .** A Z statistic with mean 0 and variance 1 is needed for each cell.

- If  $W_j = 0$ , set  $Z_j = 0$ .
- Otherwise, the actual Z statistic calculation depends on the type of performance measure.

*Mean Measure*

$$Z_j = \Phi^{-1}(\alpha)$$

where  $\alpha$  is determine by the following algorithm.

If  $\min(n_{1j}, n_{2j}) > 6$ , then determine  $\alpha$  as

$$\alpha = P(t_{n_{1j}-1} \leq T_j),$$

that is,  $\alpha$  is the probability that a t random variable with  $n_{1j} - 1$  degrees of freedom, is less than

$$T_j = t_j + \frac{g}{6} \left( \frac{n_{1j} + 2n_{2j}}{\sqrt{n_{1j} n_{2j} (n_{1j} + n_{2j})}} \right) \left( t_j^2 + \frac{n_{2j} - n_{1j}}{2n_{1j} + n_{2j}} \right),$$

where:

$$t_j = \frac{\bar{X}_{1j} - \bar{X}_{2j}}{s_{1j} \sqrt{\frac{1}{n_{1j}} + \frac{1}{n_{2j}}}}$$

and the coefficient  $g$  is an estimate of the skewness of the parent population, which we assume is the same in all cells. It can be estimated from the ILEC values in the largest cells. This needs to be done only once for each measure. We have found that attempting to estimate this skewness parameter for each cell separately leads to excessive variability in the "adjusted"  $t$ . We therefore use a single compromise value in all cells.

Note, that  $t_j$  is the "modified Z" statistic. The statistic  $T_j$  is a "modified Z" corrected for the skewness of the ILEC data.

If  $\min(n_{1j}, n_{2j}) \leq 6$ , and

a)  $M_j \leq 1,000$  (the total number of distinct pairs of samples of size  $n_{1j}$  and  $n_{2j}$  is 1,000 or less).

- Calculate the sample sum for all possible samples of size  $n_{2j}$ .
- Rank the sample sums from smallest to largest. Ties are dealt by using average ranks.
- Let  $R_0$  be the rank of the observed sample sum with respect all the sample sums.

$$\alpha = 1 - \frac{R_0 - 0.5}{M_j}$$

b)  $M_j > 1,000$

- Draw a random sample of 1,000 sample sums from the permutation distribution.
- Add the observed sample sum to the list. There is a total of 1001 sample sums. Rank the sample sums from smallest to largest. Ties are dealt by using average ranks.
- Let  $R_0$  be the rank of the observed sample sum with respect all the sample sums.

$$\alpha = 1 - \frac{R_0 - 0.5}{1001}$$

*Proportion Measure*

$$Z_j = \frac{n_j a_{1j} - n_{1j} a_j}{\sqrt{\frac{n_{1j} n_{2j} a_j (n_j - a_j)}{n_j - 1}}}$$

*Rate Measure*

$$Z_i = \frac{n_{1j} - n_j q_j}{\sqrt{n_j q_j (1 - q_j)}}$$

*Ratio Measure*

$$Z_j = \frac{\hat{R}_{1j} - \hat{R}_{2j}}{\sqrt{V(\hat{R}_{1j}) \left( \frac{1}{n_{1j}} + \frac{1}{n_{2j}} \right)}}$$

$$V(\hat{R}_{1j}) = \frac{\sum_k (U_{1jk} - \hat{R}_{1j} X_{1jk})^2}{\bar{X}_{1j}^2 (n_{1j} - 1)} = \frac{\sum_k U_{1jk}^2 - 2\hat{R}_{1j} \sum_k (U_{1jk} X_{1jk}) + \hat{R}_{1j}^2 \sum_k X_{1jk}^2}{\bar{X}_{1j}^2 (n_{1j} - 1)}$$

3. **Obtain a truncated Z value for each cell,  $Z_j^*$ .** To limit the amount of cancellation that takes place between cell results during aggregation, cells whose results suggest possible favoritism are left alone. Otherwise the cell statistic is set to zero. This means that positive equivalent Z values are set to 0, and negative values are left alone. Mathematically, this is written as

$$Z_j^* = \min(0, Z_j).$$

4. **Calculate the theoretical mean and variance of the truncated statistic under the null hypothesis of parity,  $E(Z_j^* | H_0)$  and  $\text{Var}(Z_j^* | H_0)$ .** In order to compensate for the truncation in step 3, an aggregated, weighted sum of the  $Z_j^*$  will need to be centered and scaled properly so that the final aggregate statistic follows a standard normal distribution.

- If  $W_j = 0$ , then no evidence of favoritism is contained in the cell. The formulae for calculating  $E(Z_j^* | H_0)$  and  $\text{Var}(Z_j^* | H_0)$  cannot be used. Set both equal to 0.
- If  $\min(n_{1j}, n_{2j}) > 6$  for a mean measure,  $\min\left\{a_{1j} \left(1 - \frac{a_{1j}}{n_{1j}}\right), a_{2j} \left(1 - \frac{a_{2j}}{n_{2j}}\right)\right\} > 9$  for a proportion measure,  $\min(n_{1j}, n_{2j}) > 15$  and  $n_{1j}q_j(1 - q_j) > 9$  for a rate measure, or  $n_{1j}$  and  $n_{2j}$  are large for a ratio measure then

$$E(Z_j^* | H_0) = -\frac{1}{\sqrt{2\pi}}, \text{ and}$$

$$\text{Var}(Z_j^* | H_0) = \frac{1}{2} - \frac{1}{2\pi}.$$

- Otherwise, determine the total number of values for  $Z_j^*$ . Let  $z_{ji}$  and  $\theta_{ji}$ , denote the values of  $Z_j^*$  and the probabilities of observing each value, respectively.

$$E(Z_j^* | H_0) = \sum_i \theta_{ji} z_{ji}, \text{ and}$$

$$\text{Var}(Z_j^* | H_0) = \sum_i \theta_{ji} z_{ji}^2 - [E(Z_j^* | H_0)]^2.$$

The actual values of the  $z$ 's and  $\theta$ 's depends on the type of measure.

#### Mean Measure

$$N_j = \min(M_j, 1,000), \quad i = 1, K, N_j$$

$$z_{ji} = \min \left\{ 0, \Phi^{-1} \left( 1 - \frac{R_i - 0.5}{N_j} \right) \right\} \quad \text{where } R_i \text{ is the rank of sample sum } i$$

$$\theta_{ji} = \frac{1}{N_j}$$

#### Proportion Measure

$$z_{ji} = \min \left\{ 0, \frac{n_{1j} i - n_{1j} a_j}{\sqrt{\frac{n_{1j} n_{2j} a_j (n_j - a_j)}{n_j - 1}}} \right\}, \quad i = \max(0, a_j - n_{2j}), K, \min(a_j, n_{1j})$$

$$\theta_{ji} = \text{HG}(i)$$

#### Rate Measure

$$z_{ji} = \min \left\{ 0, \frac{i - n_j q_j}{\sqrt{n_j q_j (1 - q_j)}} \right\}, \quad i = 0, K, n_j$$

$$\theta_{ji} = \text{BN}(i)$$

#### Ratio Measure

The performance measure that is in this class is billing accuracy. The sample sizes for this measure are quite large, so there is no need for a small sample technique. If one does need a small sample technique, then a resampling method can be used.

5. Calculate the aggregate test statistic,  $Z^T$ .

$$Z^T = \frac{\sum_j W_j Z_j^* - \sum_j W_j E(Z_j^* | H_0)}{\sqrt{\sum_j W_j^2 \text{Var}(Z_j^* | H_0)}}$$

### Decision Process

Once  $Z^T$  has been calculated, it is compared to a critical value to determine if the ILEC is favoring its own customers over a CLEC's customers. The derivation of the critical value is found in Appendix C.

This critical value changes as the ILEC and CLEC transaction volume change. One way to make this transparent to the decision maker, is to report the difference between the test statistic and the critical value,  $diff = Z^T - c_B$ . If favoritism is concluded when  $Z^T < c_B$ , then the  $diff < 0$  indicates favoritism.

This make it very easy to determine favoritism: a positive  $diff$  suggests no favoritism, and a negative  $diff$  suggests favoritism. Appendix D provides an example of how this information can be reported for each month.

## **Appendix B. Trunk Blocking**

This Appendix provides an example of how the trunk blocking data can be processed to apply the Truncated Z Statistic. Trunk blocking is defined as the proportion of blocked calls a trunk group experiences in a time interval. It is a ratio of two numbers—blocked and attempted calls, both of which can vary over time and across trunk groups. Since the measure is a proportion where the numerator is a subset of the denominator, the truncated Z statistic, modified for proportions, can be applied here (see Appendix A).

As with other performance measures, data are first assigned to like-to-like cells, and the Z statistic is then computed within each cell. For trunk blocking, cells are defined by three variables: hour, day, and trunk group size or capacity. The next sections will describe the data and the data processing steps in greater detail.

The approach used in this example needs to be reviewed by subject matter expert to determine if it proper to use for trunk blocking.

### **Data Sources**

Two data files are processed for the trunk blocking measure. One is the Trunk Group Data File that contains the Trunk Group Serial Number (TGSN), Common Language Location Identifier (CLLI) , and other characteristics needed to categorize trunk groups and to identify them as BellSouth or CLEC.

The other file is the Blocking Data File (BDF), which contains the actual 24 hour blocking ratios for each weekday. There are 4 or 5 weeks in a monthly report cycle. The current system, however, allows the storage of daily blocking data by hour for a week only. Therefore, the data elements necessary to compute the Truncated Z must be extracted each week.

Two important data fields of interest on the Blocking Data File are the Blocking Ratio and Offered Load. The basic definition of Blocking Ratio is the proportion of all attempted calls that were blocked. For the simplest case of one way trunk groups, this is computed by dividing the number of blocked calls by the total call attempts, given that the data are valid. If they are not valid (e.g., actual usage exceeds capacity), blocking is estimated via the Neal Wilkinson algorithm.

Although the raw data--blocked calls (overflow) and peg counts (total call attempts)--are available, the calculation of the Blocking Ratio may be complicated for two-way trunk groups and trunk groups with invalid data. For this reason, we use the blocking ratios from the BDF instead of computing the ratios from the raw data. In order to reflect different call volumes processed through each trunk group, however, the blocking ratios need to be either weighted by call volume or converted to blocked and attempted calls before they are aggregated.

The measure of call traffic volume recommended for weighting is Offered Load. Offered Load is different from call counts in that it incorporates call duration as well. Since it is not just the number of calls but the total usage—number of calls multiplied by average call duration—that determines the occurrence of any blocking, this pseudo measure, Offered Load, appears to be the best indicator of call volume.

Cells or comparison classes are determined by three factors—hour, day, and trunk group capacity (number of trunks in service). The first two factors represent natural classes because trunk blocking changes over time. The third factor is based on our finding that high blocking tends to occur in small trunk groups. A pattern was found not only in the magnitude of blocking but also in its variability. Both the magnitude and variability of blocking decrease as trunk group capacity increases. Additional work is needed to establish the appropriate number of capacity levels and the proper location of boundaries.

### Data Processing

The data are processed using the five steps below:

1. Merge the two files by TGSN and select only trunk groups listed in both files.
2. Reset the blocking of all high use trunk groups to zero<sup>1</sup>.
3. Assign trunk group categories to CLEC and BellSouth: Categories 1, 3, 4, 5, 10, and 16 for CLEC and 9 for BellSouth<sup>2</sup>. The categories used here for comparison are:

Category	Administrator	Point A	Point B
1	BellSouth	BellSouth End Office	BellSouth Access Tandem
3	BellSouth	BellSouth End Office	CLEC Switch
4	BellSouth	BellSouth Local Tandem	CLEC Switch
5	BellSouth	BellSouth Access Tandem	CLEC Switch
9	BellSouth	BellSouth End Office	BellSouth End Office
10	BellSouth	BellSouth End Office	BellSouth Local Tandem
16	BellSouth	BellSouth Tandem	BellSouth Tandem

4. Recode the missing data. The Blocking Data File assigns all missing data (no valid measurement data) zero blocking. To differentiate true zero blocking from zeroes due to missing data, invalid records were identified and the ratios reset to missing. The blocking value was invalid if both the number of Loaded Days and the Offered Load were 0 for a given hourly period.
5. Form comparison classes based either on the data (i.e., quartiles) or on a predetermined set of values.

<sup>1</sup> The high use trunk groups cannot have any blocking. These are set up such that all overflow calls are automatically routed to other trunk groups instead of being physically blocked.

<sup>2</sup> More detailed information on all categories is described in a report 'Trunk Performance Report Generation' by Ernst & Young (March 1999).

### Calculation of the Proportion of Blocked Calls

Each cell is determined by day of the month, hour of the day, and trunk group capacity. To use the Truncated Z method, we generate summary information, to include the total number of blocked calls and the total number of attempted calls, for each cell.

For the details of each calculation step, the following notation is used. For a given hour of a day, let  $\bar{X}_{1j}$  be the proportion of BellSouth blocked calls for trunk group i in cell j and  $\bar{X}_{2j}$  be the corresponding proportion for CLEC. Then  $\bar{X}_{1j} = X_{1ij} / n_{1ij}$  where  $X_{1ij}$  denotes the number of BellSouth blocked calls and  $n_{1ij}$  denotes the number of BellSouth total call attempts (indicated by Offered Load) for trunk group i in cell j. Likewise,  $\bar{X}_{2j} = X_{2ij} / n_{2ij}$ . For the steps outlined below, only the CLEC notation is provided.

1. Compute the number of blocked calls for trunk group i:  $X_{2ij} = \bar{X}_{2j} * n_{2ij}$
2. Compute total call attempts for all trunk groups in the cell:  $n_{2j} = \sum_i n_{2ij}$
3. Compute mean blocking proportion for cell j:  $\bar{X}_{2j} = \sum_i X_{2ij} / \sum_i n_{2ij}$
4. Compute the total number of BellSouth and CLEC blocked calls in cell j:  $t_j = \sum_i X_{1ij} + \sum_i X_{2ij}$
5. Apply the Truncated Z Statistic for Proportion measures presented in Appendix A.