

observed in female mice. It is concluded that both types of exposure conditions induced moderate elevation of antibody production only in male mice.

Irradiation with electromagnetic waves (8.15–18 GHz, 1 Hz within, $1 \mu\text{W}/\text{cm}^2$) in vivo increases the cytotoxic activity of natural killer cells of rat spleen [59]. In mice exposed for 24–72 h, the activity of natural killer cells increased by 130–150%, the increased level of activity persisting within 24 h after the cessation of treatment. Microwave irradiation of animals in vivo for 3.5 and 5 h, and a short exposure of splenic cells in vitro did not affect the activity of natural killer cells.

Whole body microwave sinusoidal irradiation of male NMRI mice with 8.15–18 GHz (1 Hz within) at a power density of $1 \mu\text{W}/\text{cm}^2$ caused a significant enhancement of TNF production in peritoneal macrophages and splenic T-lymphocytes [60]. Microwave radiation affected T-cells, facilitating their capacity to proliferate in response to mitogenic stimulation. The exposure duration necessary for the stimulation of cellular immunity ranged from 5 h to 3 days. Chronic irradiation of mice for 7 days produced the decreasing of TNF production in peritoneal macrophages. The exposure of mice for 24 h increased the TNF production and immune proliferative response, and these stimulatory effects persisted over 3 days after the termination of exposure. Microwave treatment increased the endogenously produced TNF more effectively than did lipopolysaccharide, one of the most potential stimuli of synthesis of this cytokine. Microwaves, thus, indeed can be a factor interfering with the process of cellular immunity!

A very intriguing investigation was carried out by Gapeev et al. [61], who compared horn, dielectric and channel antennae and their matching with various types of loads, including a biological object. The channel antenna in contrast to dielectric and horn provides the uniform spatial distribution of specific absorbed rating in the frequency range used and wide-band matching with the object both in near field and far field zones of the radiator. It is shown, that low-intensity electromagnetic radiation of extremely high frequency in near field zone of the channel radiator modifies the activity of mouse peritoneal neutrophils on a quasi-resonance manner. The interaction of electromagnetic radiation with the biological object has been revealed in the narrow-band frequencies of 41.8–42.05 GHz and consists in inhibition of luminol-dependent chemiluminescence of neutrophils activated by opsonized zymosan. No frequency dependence has been found of the electromagnetic radiation effects in the far field zone of the radiator. The results obtained suggest, that the quasi-resonance dependence of the biological effect on the frequency of the electromagnetic radiation in the near field zone is conditioned by structure and nature of the electromagnetic radiation in this zone.

In 2003, Gatta et al. [62] studied the effects of in vivo exposure to GSM-modulated 900 MHz radiation on mouse peripheral lymphocytes. The aim of this study was to evaluate whether daily whole-body exposure to 900 MHz

GSM-modulated radiation could affect spleen lymphocytes. C57BL/6 mice were exposed 2 h/day for 1, 2 or 4 weeks in a TEM cell to an SAR of 1 or 2 W/kg. Untreated and sham-exposed groups were also examined. At the end of the exposure, mice were killed and spleen cells were collected. The number of spleen cells, the percentages of B- and T-cells, and the distribution of T-cell subpopulations (CD4 and CD8) were not altered by the exposure. T- and B-cells were also stimulated *ex vivo* using specific monoclonal antibodies or LPS to induce cell proliferation, cytokine production and expression of activation markers. The results did not show relevant differences in either T- or B-lymphocytes from mice exposed to an SAR of 1 or 2 W/kg and sham-exposed mice with few exceptions. After 1 week of exposure to 1 or 2 W/kg, an increase in IFN-gamma (Ifng) production was observed that was not evident when the exposure was prolonged to 2 or 4 weeks. This suggests that the immune system might have adapted to RF radiation as it does with other stressing agents. All together, from their in vivo data, they concluded that the T- and B-cell compartments were not substantially affected by exposure to RF radiation and that a clinically relevant effect of RF radiation on the immune system is unlikely to occur. [Another explanation could be that the cells were unable to deal with the exposure and the obvious follow-up question then will be: What happened with the immune cells after months and years of exposure?]

On the other hand, Kolomytseva et al. [63], in their whole-body exposure experiment designed to study the dynamics of leukocyte number and functional activity of peripheral blood neutrophils under whole-body exposure of healthy mice to low-intensity extremely high-frequency electromagnetic radiation (EHF EMR, 42.0 GHz, $0.15 \text{ mW}/\text{cm}^2$, 20 min daily), showed that such a whole-body exposure of healthy mice to low-intensity EHF EMR has a profound effect on the indices of non-specific immunity. It was shown that the phagocytic activity of peripheral blood neutrophils was suppressed by about 50% ($p < 0.01$ as compared with the sham-exposed control) in 2–3 h after the single exposure to EHF EMR. The effect persisted for 1 day after the exposure, and then the phagocytic activity of neutrophils returned to the normal within 3 days. A significant modification of the leukocyte blood profile in mice exposed to EHF EMR for 5 days was observed after the cessation of exposures: the number of leukocytes increased by 44% ($p < 0.05$ as compared with sham-exposed animals), mostly due to an increase in the lymphocyte content. The supposition was made that EHF EMR effects can be mediated via the metabolic systems of arachidonic acid and the stimulation of adenylate cyclase activity, with subsequent increase in the intracellular cAMP level.

The modification of indices of the humoral immune response to thymus-dependent antigen (sheep erythrocytes) after a whole-body exposure of healthy mice to low-intensity extremely high-frequency electromagnetic radiation was reported by Lushnikov et al. in 2001 [64]. Male NMRI mice were exposed in the far-field zone of horn antenna at a frequency of 42.0 GHz and energy flux density of $0.15 \text{ mW}/\text{cm}^2$

under different regimes: once for 20 min, for 20 min daily during 5 and 20 successive days before immunization, and for 20 min daily during 5 successive days after immunization throughout the development of the humoral immune response. The intensity of the humoral immune response was estimated on day 5 after immunization by the number of antibody-forming cells of the spleen and antibody titers. Changes in cellularity of the spleen, thymus and red bone marrow were also assessed. The indices of humoral immunity and cellularity of lymphoid organs changed insignificantly after acute exposure and a series of five exposures before and after immunization of the animals. However, after repeated exposures for 20 days before immunization, a statistically significant reduction of thymic cellularity by 17.5% ($p < 0.05$) and a decrease in cellularity of the spleen by 14.5% ($p < 0.05$) were revealed. The results show that single low-intensity extremely high-frequency electromagnetic radiation, at the frequency and energy flux density used, does not influence the humoral immune response intensity in healthy mice but influences immunogenesis under multiple repeated exposures.

Experiments have also been conducted to elucidate the effects of chronic low power-level microwave radiation on the immunological systems of rabbits [65]. Fourteen male Belgian white rabbits were exposed to microwave radiation at 5 mW/cm², 2.1 GHz, 3 h daily, 6 days/week for 3 months in two batches of seven each in specially designed miniature anechoic chambers. Seven rabbits were subjected to sham exposure for identical duration. The microwave energy was provided through S band standard gain horns connected to a 4K3SJ2 Klystron power amplifier. The first batch of animals was assessed for T-lymphocyte-mediated cellular immune response mechanisms and the second batch of animals for B-lymphocyte-mediated humoral immune response mechanisms. The peripheral blood samples collected monthly during microwave/sham exposure and during follow-up (5/14 days after termination of exposures, in the second batch animals only) were analysed for T-lymphocyte numbers and their mitogen responsiveness to ConA and PHA. Significant suppression of T-lymphocyte numbers was noted in the microwave group at 2 months ($p < 0.01$) and during follow-up ($p < 0.01$). The first batch of animals was initially sensitised with BCG and challenged with tuberculin (0.03 ml) at the termination of microwave irradiation/sham exposure and the increase in foot pad thickness (delta mm), which is a measure of T-cell-mediated immunity (delayed type hypersensitivity response, DTH) was noted in both the groups. The microwave group revealed a "better" response than the control group (delta % +12.4 versus +7.54).

Nasta et al. [66], very recently examined the effects of *in vivo* exposure to a GSM-modulated 900 MHz RF field on B-cell peripheral differentiation and antibody production in mice. Their results show that exposure to a whole-body average SAR of 2 W/kg, 2 h/day for 4 consecutive weeks does not affect the frequencies of differentiating transitional 1 (T1) and T2 B-cells or those of mature follicular B and marginal zone B-cells in the spleen. IgM and IgG serum levels are also

not significantly different among exposed, sham-exposed and control mice. B-cells from these mice, challenged *in vitro* with LPS, produce comparable amounts of IgM and IgG. Moreover, exposure of immunized mice to RF fields does not change the antigen-specific antibody serum level. Interestingly, not only the production of antigen-specific IgM but also that of IgG (which requires T–B-cell interaction) is not affected by RF-field exposure. This indicates that the exposure does not alter an ongoing *in vivo* antigen-specific immune response. In conclusion, the results of Nasta et al. [66] do not indicate any effects of GSM-modulated RF radiation on the B-cell peripheral compartment and antibody production.

Whole-body microwave sinusoidal irradiation of male NMRI mice, exposure of macrophages *in vitro*, and preliminary irradiation of culture medium with 8.15–18 GHz (1 Hz within) at a power density of 1 μ W/cm² caused a significant enhancement of tumour necrosis factor production in peritoneal macrophages [67]. The role of microwaves as a factor interfering with the process of cell immunity must, thus, be seriously considered. Furthermore, the effect of 8.15–18 GHz (1 Hz within) microwave radiation at a power density of 1 μ W/cm² on the tumour necrosis factor (TNF) production and immune response was tested by Novoselova et al. [68]. A single 5 h whole-body exposure induced a significant increase in TNF production in peritoneal macrophages and splenic T-cells. The mitogenic response in T-lymphocytes increased after microwave exposure. The activation of cellular immunity was observed within 3 days after exposure. A diet containing lipid-soluble nutrients (beta-carotene, alpha-tocopherol and ubiquinone Q9) increased the activity of macrophages and T-cells from irradiated mice. These results demonstrate that irradiation with low-power density microwaves stimulates the immune potential of macrophages and T-cells, and the antioxidant treatment enhances the effect of microwaves, in particular when the effect of irradiation is reduced.

In the experimental study by Çetin et al. [69] in 2006, the hematological effects of pulsed EMFs chronic exposure were investigated on mice. Sixty, 6-week-old male Swiss mice, weighing 40–45 g were used, and were divided into two groups: in one group, animals ($n=30$) were exposed to pulsed EMFs (60 Hz, intensity 3 μ T, 12 h by day) for a 120-day period, whereas the second group ($n=30$) was used as control. On days 15, 30, 90 and 120, samples were taken by cardiac puncture for the hematological analysis (red blood cell and white blood cell counts, leukocyte distribution). Whereas no significant difference was noted between control and exposed animals at the 15th and the 30th days, a macrocytic anemia characterized by decreases in of hemoglobin concentration, hematocrit values and erythrocyte counts and by increases in mean corpuscular volume, occurred in the exposed animals on day 90. Furthermore, they have shown significant reductions of leukocyte, lymphocyte and neutrophil counts, while monocyte counts were increased. On day 120, these leukocyte alterations were still

observed, whereas erythrocyte parameters approached control values. These results suggest that pulsed electromagnetic fields (60 Hz and 3 μ T) affect the hematological parameters of mice, probably by reducing proliferation and differentiation of marrow stem cells.

Obukhan [70] has performed cytologic investigations designed to study bone marrow, peripheral blood, spleen, and thymus of albino rats irradiated by an EMF of 2375, 2450, and 3000 MHz. Structural and functional changes in populations of megakaryocytes, immunocompetent cells as well as of undifferentiated cells, and of other types of cells that are dependent on the intensity of irradiation were revealed. The results permitted establishing the probability-threshold levels of exposure taking account of reactions of perception and physiologic adaptation together with compensatory and regenerative processes and the injury sustained. It was shown that changes in bone marrow cells differentiation and reproduction, rather than integral shifts in the peripheral blood, acquired the utmost significance. The blast cell population increased in low-intensity exposure, along with disturbances in mitosis.

The possibility of genotoxicity of radiofrequency radiation (RFR) applied alone or in combination with X-rays was recently investigated *in vitro* using several assays on human lymphocytes by Stronati et al. [71]. The chosen SAR values are near the upper limit of energy absorbed by localized tissue during the use of some cellular telephones. The purpose of the combined exposures was to examine whether RFR might act epigenetically by reducing the fidelity of repair of DNA damage caused by a well-characterized and established mutagen. Blood specimens from 14 donors were exposed continuously for 24 h to a GSM basic 935 MHz signal. The signal was applied at two SAR; 1 and 2 W/Kg, alone or combined with a 1-min exposure to 1.0 Gy of 250 kVp X-rays given immediately before or after the RFR. The assays employed were the alkaline comet technique to detect DNA strand breakage, metaphase analyses to detect unstable chromosomal aberrations and sister chromatid exchanges, micronuclei in cytokinesis-blocked binucleate lymphocytes and the nuclear division index to detect alterations in the speed of *in vitro* cell cycling. By comparison with appropriate sham-exposed and control samples, no effect of RFR alone could be found for any of the assay endpoints. In addition, RFR did not modify any measured effects of the X-radiation. In conclusion, this study has used several standard *in vitro* tests for chromosomal and DNA damage in Go human lymphocytes exposed *in vitro* to a combination of X-rays and RFR. It has comprehensively examined whether a 24-h continuous exposure to a 935 MHz GSM basic signal delivering SAR of 1 or 2 W/Kg is genotoxic *per se* or whether, it can influence the genotoxicity of the well-established clastogenic X-radiation. Within the experimental parameters of the study in all instances no effect from the RFR signal was observed. [Of course, DNA damage is a well characterized effect of electromagnetic irradiation of other cell types, including lymphoblastoid cells [72], fibroblasts [73], and brain cells [74].]

Despite the important role of the immune system in defending the body against infections and cancer, only rather few investigations on possible effects of RF radiation on function of human immune cells have been undertaken. One of these is the investigation by Tuschl et al. [75] in 2006 where they assessed whether GSM modulated RF fields have adverse effects on the functional competence of human immune cells. Within the frame of a multidisciplinary project "Biological effects of high frequency electromagnetic fields (EMFs)" sponsored by the National Occupation Hazard Insurance Association (AUVA), *in vitro* investigations were carried out on human blood cells. Exposure was performed at GSM basic 1950 MHz, an SAR of 1 mW/g in an intermittent mode (5 min "ON", 10 min "OFF") and a maximum ΔT of 0.06 °C for the duration of 8 h. The following immune parameters were evaluated: (1) the intracellular production of interleukin-2 (IL-2) and interferon (INF) gamma in lymphocytes, and IL-1 and TNF-alpha in monocytes were evaluated with monoclonal antibodies. (2) The activity of immune-relevant genes (IL-1-alpha and beta, IL-2, IL-2-receptor, IL-4, macrophage colony stimulating factor (MCSF)-receptor, TNF-alpha, TNF-alpha-receptor) and housekeeping genes was analyzed with real time PCR. (3) The cytotoxicity of lymphokine activated killer cells (LAK cells) against a tumour cell line was determined in a flow cytometric test. For each parameter, blood samples of at least 15 donors were evaluated. No statistically significant effects of exposure were found and there was no indication that emissions from mobile phones are associated with adverse effects on the human immune system.

Chagnaud and Veyret [76] in 1999 could also not demonstrate an effect of low-level pulsed microwaves on the integrity of the immune system. They investigated the effects of GSM-modulated microwaves on lymphocyte sub-populations of Sprague-Dawley rats and their normal mitogenic responses using flow cytometry analysis and a colorimetric method. No alterations were found in the surface phenotype of splenic lymphocytes or in their mitogenic activity.

[N.B. One must always be very cautious when it comes to negative findings. For example: of the 100 or so papers on genotoxic effects of RF fields, a majority has been done with peripheral blood lymphocytes. Except for special cases, these cells are highly protected from their upregulated repair enzymes. These cells are often used to investigate chemical genotoxicity, because in these cases the toxicity often occurs due to the action of the repair enzymes. Repair-deficient cells remain intact! The mechanisms of action of EMFs may not be clearly understood, but it is unlikely they mimic such chemical enzyme-induced genotoxicity. — Yet another example is the use of mice and rats for the study increases in brain tumour incidence due to mobile telephony exposure. Since the incidence data from human studies point to a needed exposure time of at least 5 years, and mice and rats do not live longer

than 2 years, the rats will die long before they have had a chance to develop the tumours!]

Irradiation by pulsed microwaves (9.4 GHz, 1 μ s pulses at 1000/s), both with and without concurrent amplitude modulation (AM) by a sinusoid at discrete frequencies between 14 and 41 MHz, was assessed for effects on the immune system of Balb/c mice [77]. The mice were immunized either by sheep red blood cells (SRBC) or by glutaric-anhydride conjugated bovine serum albumin (GA-BSA), then exposed to the microwaves at a low rms power density (30 μ W/cm²; whole-body averaged SAR approximately 0.015 W/kg). Sham exposure or microwave irradiation took place during each of 5 contiguous days, 10 h/day. The antibody response was evaluated by the plaque-forming cell assay (SRBC experiment) or by the titration of IgM and IgG antibodies (GA-BSA experiment). In the absence of AM, the pulsed field did not greatly alter immune responsiveness. In contrast, exposure to the field under the combined-modulation condition resulted in significant, AM-frequency-dependent augmentation or weakening of immune responses.

To study the possible RF effects on human lymphocyte activation, Capri et al. [78] analyzed CD25, CD95, CD28 molecules in unstimulated and stimulated CD4+ or CD8+ T-cells *in vitro*. Peripheral blood mononuclear cells (PBMCs) from young and elderly donors were exposed or sham-exposed to RF (1800 MHz, SAR 2 W/kg) with or without mitogenic stimulation. No significant changes in the percentage of these cell subsets were found between exposed and sham-exposed lymphocytes in both young and elderly donors. Nevertheless, after RF exposure they observed a slight, but significant, downregulation of CD95 expression in stimulated CD4+ T-lymphocytes from elderly, but not from young donors. This age-related result is noteworthy given the importance of such a molecule in regulation of the immune response.

In the paper by Yurekli et al. [79], a GHz transverse electromagnetic (GTEM) cell was used as an exposure environment for plane wave conditions of far-field free space EMF propagation at the GSM base transceiver station (BTS) frequency of 945 MHz, and effects on oxidative stress in rats were investigated. When EMFs at a power density of 3.67 W/m² (SAR = 11.3 mW/kg), which is well below current exposure limits, were applied, MDA (malondialdehyde) level was found to increase and GSH (reduced glutathione) concentration was found to decrease significantly ($p < 0.0001$). Additionally, there was a less significant ($p = 0.019$) increase in SOD (superoxide dismutase) activity under EMF exposure.

Since genotoxic effects of the second generation standard GSM have been reported after exposure of human cells *in vitro*, Schwarz et al. [80] decided to use human cultured fibroblasts of three different donors and three different short-term human lymphocyte cultures and expose them to 1950 MHz UMTS below the SAR safety limit of 2 W/kg. The alkaline comet assay and the micronucleus assay

were used to ascertain dose and time-dependent genotoxic effects. Five hundred cells per slide were visually evaluated in the comet assay and comet tail factor (CTF) was calculated. In the micronucleus assay 1000 binucleated cells were evaluated per assay. The origin of the micronuclei was determined by fluorescence labeled anticentromere antibodies. All evaluations were performed under blinded conditions. UMTS exposure increased the CTF and induced centromere-negative micronuclei (MN) in human cultured fibroblasts in a dose and time-dependent way. Incubation for 24 h at an SAR of 0.05 W/kg generated a statistically significant rise in both CTF and MN ($p = 0.02$). At an SAR of 0.1 W/kg the CTF was significantly increased after 8 h of incubation ($p = 0.02$), the number of MN after 12 h ($p = 0.02$). However, under these conditions, no UMTS effect was obtained with lymphocytes, either unstimulated or stimulated with phytohemagglutinin. The authors conclusion was that UMTS exposure may cause genetic alterations in some but not in all human cells *in vitro*.

Simkó and Mattsson [81] have presented a hypothesis of a possible initial cellular event affected by exposure to ELF-EMF, an event that is compatible with the multitude of effects observed after exposure. Based on an extensive literature review, they suggested that ELF-EMF exposure is able to perform such activation by means of increasing levels of free radicals. Such a general activation is compatible with the diverse nature of observed effects. Free radicals are intermediates in natural processes, like mitochondrial metabolism, and are also a key feature of phagocytosis. Free radical release is inducible by ionizing radiation or phorbol ester treatment, both leading to genomic instability. EMFs might be a stimulus to induce an "activated state" of the cell such as phagocytosis, which then enhances the release of free radicals, in turn leading to genotoxic events. Simkó and Mattsson envisaged that EMF exposure can cause both acute and chronic effects that are mediated by increased free radical levels: (1) Direct activation of, for example macrophages (or other cells) by short-term exposure to EMF leads to phagocytosis (or other cell-specific responses) and consequently, free radical production. This pathway may be utilized to positively influence certain aspects of the immune response, and could be useful for specific therapeutic applications. (2) EMF-induced macrophage (cell) activation includes direct stimulation of free radical production. (3) An increase in the lifetime of free radicals by EMF leads to persistently elevated free radical concentrations. In general, reactions in which radicals are involved become more frequent, increasing the possibility of DNA damage. (4) Long-term EMF exposure leads to a chronically increased level of free radicals, subsequently causing an inhibition of the pineal gland hormone melatonin. Taken together, these EMF-induced reactions could lead to a higher incidence of DNA damage and therefore, to an increased risk of tumour development. While the effects on melatonin and the extension of the lifetime of radicals can explain the link between EMF

exposure and the incidence of for example leukaemia, the two additional mechanisms described by them specifically for mouse macrophages, can explain the possible correlation between immune cell system stimulation and EMF exposure.

5.7.4. Effects of EMFs on the immune system at pregnancy

Nakamura et al. [82] have investigated a very important issue, namely what happens to pregnant rats exposed to microwaves. Earlier data had indicated that these microwaves produce various detrimental changes based on actions of heat or non-specific stress, although the effects of microwaves on pregnant organisms were not uniform. This study was therefore designed to clarify the effect of exposure to microwaves during pregnancy on endocrine and immune functions. Natural killer cell activity and natural killer cell subsets in the spleen were measured, as well as some endocrine indicators in blood: corticosterone and adrenocorticotrophic hormone (ACTH) as indices of the hypothalamic–pituitary–adrenal axis; beta-endorphin, oestradiol, and progesterone in six female virgin rats and six pregnant rats (9–11 days gestation) exposed to microwaves at 10 mW/cm² incident power density at 2450 MHz for 90 min. The same measurements were performed in control rats (six virgin and six pregnant rats). Skin temperature in virgin and pregnant rats increased immediately after exposure to microwaves. Although splenic activity of natural killer cells and any of the subset populations identified by the monoclonal antibodies CD16 and CD57 did not differ in virgin rats with or without exposure to microwaves, pregnant rats exposed to microwaves showed a significant reduction of splenic activity of natural killer cells and CD16+ CD57- ones. Although corticosterone and ACTH increased, and oestradiol decreased in exposed virgin and pregnant rats, microwaves produced significant increases in beta-endorphin and progesterone only in pregnant rats. So, in summary, Nakamura et al. [82] showed that microwaves at the power of 10 mW/cm² produced activation of the hypothalamic–pituitary–adrenal axis and increased oestradiol in both virgin and pregnant rats, indicating that microwaves are a great stress in pregnancy.

In the following year, 1998, the same groups of scientists published a new study [83] in which they examined the involvement of opioid systems in reduced natural killer cell activity (NKCA) in pregnant rats exposed to microwaves at a relatively low level (2 mW/cm² incident power density at 2450 MHz for 90 min). They assayed beta-endorphin (betaEP) in blood, pituitary lobes, and placenta as well as splenic NKCA in virgin and/or pregnant rats. Although microwaves elevated colonic temperatures by 0.8 °C for virgin and 0.9 °C for pregnant rats, and betaEP in blood and anterior pituitary lobes (AP) significantly, it did not change blood corticosterone as an index of hypothalamic–pituitary–adrenal axis. There were significant interactions between pregnancy and microwave exposure on splenic NKCA, betaEP in both blood and AP, and blood progesterone.

Intra-peritoneal administration of opioid receptor antagonist naloxone prior to microwave exposure increased NKCA, blood, and placental betaEP in pregnant rats. Alterations in splenic NKCA, betaEP and progesterone in pregnant rats exposed to microwaves may be due to both thermal and non-thermal actions. These results suggest that NKCA reduced by microwaves during pregnancy is mediated by the pituitary opioid system.

To further clarify the effects of microwaves on pregnancy, uterine or uteroplacental blood flow and endocrine and biochemical mediators, including corticosterone, estradiol, prostaglandin E(2) (PGE(2)), and prostaglandin F(2)alpha (PGF(2)alpha), Nakamura et al. published yet another study in 2000 [84]. In this article they measured these parameters and factors in rats exposed to continuous-wave (CW) microwave at 2 mW/cm² incident power density at 2450 MHz for 90 min. Colonic temperature in virgin and pregnant rats was not significantly altered by microwave treatment. Microwaves decreased uteroplacental blood flow and increased progesterone and PGF(2)alpha in pregnant, but not in virgin rats. Intraperitoneal (i.p.) administration of angiotensin II, a uteroplacental vasodilator, before microwave exposure prevented the reduction in uteroplacental blood flow and the increased progesterone and PGF(2)alpha in pregnant rats. Increased corticosterone and decreased estradiol during microwave exposure were observed independent of pregnancy and pretreatment with angiotensin II. These results suggest that microwaves (CW, 2 mW/cm²; 2450 MHz) produce uteroplacental circulatory disturbances and ovarian and placental dysfunction during pregnancy, probably through non-thermal actions. The uteroplacental disturbances appear to be due to actions of PGF(2)alpha and may pose some risk for pregnancy! [Could the above findings be part of the explanation behind the sensational findings of Magras and Xenos [85] from 1997?]

5.7.5. Synchronization of cerebral rhythms. Important for the brain-immune system axis?

Vecchio et al. [86] have reported that EMF from mobile phones affects the synchronization of cerebral rhythms. Their findings suggest that prolonged exposure to mobile phone emissions affect cortical activity and the speed of neural synchronization by interhemispherical functional coupling of EEG rhythms. This may be evidence that such exposure can affect the way in which the brain is able to process information, by interfering with the synchronization rhythms between the halves of the brain, and by deregulating the normal alpha wave 2 (about 8–10 Hz) and alpha 3 (10–12 Hz) bands. [Could such deregulation affect the brain-immune system axis? If so, what implications would it have in the short- as well as in the long-term?]

5.7.6. Classical contact allergy reactions

Finally, in addition, classical contact allergy reactions, such as chromate allergy, have been studied by Seishima et al. [87]. The background for the study was an earlier case

report about a patient with allergic contact dermatitis caused by hexavalent chromium plating on a cellular phone. The new study described the clinical characteristics and results of patch tests (closed patch tests and photopatch tests were performed using metal standard antigens) in eight patients with contact dermatitis possibly caused by handling a cellular phone. The eight patients were four males and four females aged from 14 to 54 years. They each noticed skin eruptions after 9–25 days of using a cellular phone. All patients had erythema, and seven had papules on the hemilateral auricle or in the preauricular region. Three of eight patients had a history of metal allergy. Chromate, aluminium and acrylonitrile–butadiene–styrene copolymer were used as plating on the cellular phones used by these patients. The patch test was positive for 0.5%, 0.1% and 0.05% potassium dichromate in all eight patients. The photopatch test showed the same results. One patient was positive for 2% cobalt chloride and one for 5% nickel sulfate. Based on these data, it is important to consider the possibility of contact dermatitis due to a cellular phone, possibly caused by chromate, when the patients have erythema and papules on the hemilateral auricle or in the preauricular region.

6. Effects of electromagnetic fields on other biological systems

Some parallel investigations, pointing to severe biological effects that need to be mentioned are, for instance, the results of Roux et al. [88] in 2008. Using an especially designed facility, the Mode Stirred Reverberation Chamber, they exposed tomato plants (*Lycopersicon esculentum* Mill. VFN8) to low level (900 MHz, 5 V/m) EMF for a short period (10 min) and measured changes in abundance of three specific mRNA soon after exposure. Within minutes of stimulation, stress-related mRNA (calmodulin, calcium-dependent protein kinase and proteinase inhibitor) accumulated in a rapid, large and 3-phase manner typical of an environmental stress response. Accumulation of these transcripts into the polysomal RNA also took place (indicating that the encoded proteins were translated) but was delayed (indicating that newly-synthesized mRNA was not immediately recruited into polysomes). Transcript accumulation was maximal at normal Ca(2+) levels and was depressed at higher Ca(2+), especially for those encoding calcium-binding proteins. Removal of Ca(2+) (by addition of chelating agents or Ca(2+) channel blocker) led to total suppression of mRNA accumulation. Finally, 30 min after the electromagnetic treatment, ATP concentration and adenylate energy charge were transiently decreased, while transcript accumulation was totally prevented by application of the uncoupling reagent, CCCP. These responses occur very soon after exposure, strongly suggesting that they are the direct consequence of application of radiofrequency fields, and their similarities to wound responses strongly suggests that this radiation is perceived by plants as an injurious stimulus! [Furthermore, it is

impossible to interpret these reactions as “psychological or psychiatric personality disturbances, cognitive malfunction, or likewise”.]

Also, the data from Divan et al. [89] deserve to be mentioned. They examined the association between prenatal and postnatal exposure to cell phones and behavioral problems in young children. Mothers were recruited to the Danish National Birth Cohort early in pregnancy. When the children of those pregnancies reached 7 years of age in 2005 and 2006, mothers were asked to complete a questionnaire regarding the current health and behavioral status of children, as well as past exposure to cell phone use. Mothers evaluated the child's behavior problems using the Strength and Difficulties Questionnaire. Mothers of 13,159 children completed the follow-up questionnaire reporting their use of cell phones during pregnancy as well as current cell phone use by the child. Greater odds ratios for behavioral problems were observed for children who had possible prenatal or postnatal exposure to cell phone use. After adjustment for potential confounders, the odds ratio for a higher overall behavioral problems score was 1.80 (95% confidence interval = 1.45–2.23) in children with both prenatal and postnatal exposure to cell phones. Exposure to cell phones prenatally – and, to a lesser degree, postnatally – was associated with behavioral difficulties such as emotional and hyperactivity problems around the age of school entry. [An obvious follow-up question would be “What about immune function alterations?”.] Naturally, and hopefully, these associations may be non-causal and may be due to unmeasured confounding. But if real, they would be of public health concern given the widespread use of this technology.

The exposure to non-thermal microwave EMF generated by mobile phones affects the expression of many proteins. This effect on transcription and protein stability can be mediated by the MAPK (mitogen-activated protein kinase) cascades, which serve as central signaling pathways and govern essentially all stimulated cellular processes. Indeed, long-term exposure of cells to mobile phone irradiation results in the activation of p38 as well as the ERK (extracellular signal-regulated kinase) MAPKs. Friedman et al. [90] recently have studied the immediate effect of irradiation on the MAPK cascades, and found that ERKs, but not stress-related MAPKs, are rapidly activated in response to various frequencies and intensities. Using signaling inhibitors, they delineated the mechanism that is involved in this activation. They found that the first step is mediated in the plasma membrane by NADH oxidase, which rapidly generates ROS (reactive oxygen species). These ROS then directly stimulate MMPs (matrix metalloproteinases) and allow them to cleave and release Hb-EGF (heparin-binding EGF (epidermal growth factor)). This secreted factor activates the EGF receptor, which in turn further activates the ERK cascade. Thus, their study demonstrates for the first time a detailed molecular mechanism by which electromagnetic irradiation from mobile phones induces the activation of the ERK cascade and thereby induces transcription and other cellular processes.

The terminal deoxynucleotide transferase dUTP nick end labeling (TUNEL) assay, a well known technique widely used for detecting fragmented DNA in various types of cells, was used by Panagopoulos et al. [91] to detect cell death (DNA fragmentation) in a biological model, the early and mid stages of oogenesis of the insect *Drosophila melanogaster*. The flies were exposed in vivo to either GSM 900 MHz or DCS 1800 MHz radiation from a common digital mobile phone, for few minutes per day during the first 6 days of their adult life. The exposure conditions were similar to those to which a mobile phone user is exposed. Previous results from the same group [92–94] had shown a large decrease in the oviposition of the same insect caused by GSM radiation. The recent results suggest that this decrease in oviposition, is due to degeneration of large numbers of egg chambers after DNA fragmentation of their constituent cells, induced by both types of mobile telephony radiation. Induced cell death is recorded for the first time, in all types of cells constituting an egg chamber (follicle cells, nurse cells and the oocyte) and in all stages of the early and mid-oogenesis, from germarium to stage 10, during which programmed cell death does not physiologically occur. Germarium and stages 7–8 were found to be the most sensitive developmental stages also in response to electromagnetic stress induced by the GSM and DCS fields and, moreover, germarium was found to be even more sensitive than stages 7–8.

7. Conclusions

- Both human and animal studies report large immunological changes upon exposure to environmental levels of modern, human-made EMFs. Some of these exposure levels are equivalent to those of wireless technologies in daily life, and often at low or very low (i.e., non-thermal) levels.
- Measurable physiological changes (mast cells increases, for example) that are bedrock indicators of allergic response and inflammatory conditions are stimulated by EMF exposures.
- Chronic exposure to such factors that increase allergic and inflammatory responses on a continuing basis may be harmful to health. The data presented here, as well as the very rapid international increase in incidence of allergies, asthma and other oversensitivities, together form a clear warning signal.
- It is, thus, possible that chronic provocation by exposure to EMF can lead to immune dysfunction, chronic allergic responses, inflammatory responses and ill health if they occur on a continuing basis over time. This is an area that should be investigated immediately.
- Specific findings from studies on exposures to various types of modern equipment and/or EMFs report over-reaction of the immune system; morphological alterations of immune cells; profound increases in mast cells in the upper skin layers, increased degranulation of mast cells and larger size of mast cells in electrohypersensitive indi-

viduals; presence of biological markers for inflammation which are sensitive to EMF exposure at non-thermal levels; changes in lymphocyte viability; decreased count of NK cells; decreased count of T-lymphocytes; negative effects on pregnancy (uteroplacental circulatory disturbances and placental disfunction); suppressed or impaired immune function; and inflammatory responses that can ultimately result in cellular, tissue and organ damage.

- The functional impairment electrohypersensitivity is reported by individuals in the United States, Sweden, Switzerland, Germany, Belgium, Italy, The Netherlands, Norway, Denmark and many other countries of the world. Estimates range from 3% to perhaps 10% of populations, and appear to be a growing condition of ill-health leading to lost work and productivity.
- The WHO and IEEE literature surveys do not include all of the relevant papers cited here, leading to the conclusion that evidence has been ignored in the current WHO ELF Health Criteria Monograph; and the proposed new IEEE C95.1 RF public exposure limits.
- The current international public safety limits for EMFs do not appear to be sufficiently protective of public health at all, based on the studies of immune function. New, biologically based public standards are warranted that take into account low-intensity effects on immune function and health that are reported in the scientific literature. Also the accessibility needs of persons with the functional impairment electrohypersensitivity must be fully addressed and resolved as dictated by the UN 22 “Standard rules on the equalization of opportunities for people with disabilities” (about the UN 22 Standard Rules, see website: <http://www.un.org>; since 2007 they have been upgraded into the UN “Convention on Human Rights for Persons with Functional Impairments”).

The conclusion of the above must be that there are a number of very strong indications of EMFs being capable of disturbing the immune system and thus increasing disease, including cancer, risk. It is somewhat odd that professional epidemiologists for the last 50 years have not addressed the issue of reduced repair but only looked at increased cell damages from different agents and environments when trying to understand trend changes.

Based on this review as well as on the recent Bioinitiative Report [<http://www.bioinitiative.org/>] [1], it must be concluded that the existing public safety limits are inadequate to protect public health. From a public health policy standpoint, new public safety limits, and limits on further deployment of untested technologies, are warranted.

New biologically based public and occupational exposure are recommended to address bioeffects and potential adverse health effects of chronic exposure. These effects are now widely reported to occur at exposure levels significantly below most current national and international limits. Therefore, biologically based exposure standards are needed to prevent disruption of normal body processes. Effects are

reported for DNA damage (genotoxicity that is directly linked to integrity of the human genome), cellular communication, cellular metabolism and repair, cancer surveillance within the body; and for protection against cancer and neurological diseases. Also reported are neurological effects including changes in brainwave activity during cell phone calls, impairment of memory, attention and cognitive function; sleep disorders, cardiac effects; and – as reported here – serious impact on the immune function (allergic and inflammatory responses).

The current recommendation must be a biologically based exposure limit that is completely protective against e.g. extremely low frequency and radiofrequency fields which, with chronic exposure, can reasonably be presumed to result in no adverse impacts on health and well-being. Today, such a completely protective safety limit would, for many exposure situations, be zero.

Finally, attention to the above need would also mean a great gain in future public health costs for the entire electrified world. To do the opposite could turn out to be very expensive.

Acknowledgements

Supported by the Karolinska Institute, the Help Foundation (Hjälpfonden), Stiftelsen Tornspiran, and the Cancer and Allergy Foundation (Cancer- och Allergifonden). Ms Marianne Ekman is warmly acknowledged for her expert help with literature search engines, Mr. Philippe Hug for constructive criticism, and Mr. Jonas Aule for his superb technical support without which this paper would never have been completed.

References

- [1] C.F. Blackman, M. Blank, M. Kundi, C. Sage, D.O. Carpenter, Z. Davanipour, D. Gee, L. Hardell, O. Johansson, H. Lai, K.H. Mild, A. Sage, E.L. Sobel, Z. Xu, G. Chen, The Bioinitiative Report—A Rationale for a Biologically-based Public Exposure Standard for Electromagnetic Fields (ELF and RF), <http://www.bioinitiative.org>, 2007.
- [2] B.R.R. Persson, L.G. Salford, A. Brun, Blood-brain barrier permeability in rats exposed to electromagnetic fields used in wireless communication, *Wireless Networks* 3 (1997) 455–461.
- [3] L.G. Salford, A.E. Brun, J.L. Eberhardt, L. Malmgren, B.R. Persson, Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones, *Environ. Health Perspect.* 111 (2003) 881–883.
- [4] O. Johansson, M. Hilliges, V. Björnhagen, K. Hall, Skin changes in patients claiming to suffer from “screen dermatitis”: a two-case open-field provocation study, *Exp. Dermatol.* 3 (1994) 234–238.
- [5] O. Johansson, S. Gangi, Y. Liang, K. Yoshimura, C. Jing, P.-Y. Liu, Cutaneous mast cells are altered in normal healthy volunteers sitting in front of ordinary TV/PCs—results from open-field provocation experiments, *J. Cutan. Pathol.* 28 (2001) 513–519.
- [6] V. Rajkovic, M. Matavulj, O. Johansson, Histological characteristics of cutaneous and thyroid mast cell populations in male rats exposed to power-frequency electromagnetic fields, *Int. J. Radiat. Biol.* 81 (2005) 491–499.
- [7] V. Rajkovic, M. Matavulj, O. Johansson, The effect of extremely low-frequency electromagnetic fields on skin and thyroid amine- and peptide-containing cells in rats: an immunohistochemical and morphometrical study, *Environ. Res.* 99 (2005) 369–377.
- [8] V. Rajkovic, M. Matavulj, O. Johansson, Light and electron microscopic study of the thyroid gland in rats exposed to power-frequency electromagnetic fields, *J. Exp. Biol.* 209 (2006) 3322–3328.
- [9] M. Peakman, *Basic and Clinical Immunology*, Churchill Livingstone, New York, 2008.
- [10] R.R.A. Coombs, P.G.H. Gell, P.J. Lachmann, *Clinical Aspects of Immunology*, Blackwell Scientific Publisher, Oxford, 1962.
- [11] K. Jokela, L. Puranen, A.P. Sihvonen, Assessment of the magnetic field exposure due to the battery current of digital mobile phones, *Health Phys.* 86 (2004) 56–66.
- [12] C. Sage, O. Johansson, S.A. Sage, Personal digital assistant (PDA) cell phone units produce elevated extremely-low frequency electromagnetic field emissions, *Bioelectromagnetics* 28 (2007) 386–392.
- [13] O. Johansson, Electrohypersensitivity: state-of-the-art of a functional impairment, *Electromagn. Biol. Med.* 25 (2006) 245–258.
- [14] R. Cox, Electrical hypersensitivity—human studies in the UK, in: Conference Presentation at the WHO International Workshop on EMF Hypersensitivity, Prague, Czech Republic, October 25–27, 2004.
- [15] E. Fox, Electrosensitivity symptoms associated with electromagnetic field exposure, in: Conference Presentation at the WHO International Workshop on EMF Hypersensitivity, Prague, Czech Republic, October 25–27, 2004.
- [16] P. Levallois, K. Neutra, G. Lee, L. Hristova, Study of self-reported hypersensitivity to electromagnetic fields in California, *Environ. Health Perspect.* 110 (4) (2002) 619–623.
- [17] Institut für angewandte Sozialwissenschaft GmbH (infas)/Bundesamt für Strahlenschutz, Ermittlung der Befürchtungen und Ängste der breiten Öffentlichkeit hinsichtlich möglicher Gefahren der Hochfrequenten Elektromagnetischen Felder der Mobilfunks—Jährliche Umfrage, <http://www.infas.de>, 2004.
- [18] J. Schütz, C. Peters, U.T. Egle, B. Jansen, R. Kimbel, S. Letzel, W. Nix, L.G. Schmidt, L. Vollrath, The “Mainzer EMF-Wachhund”: results from a watchdog project on self-reported health complaints attributed to exposure to electromagnetic fields, *Bioelectromagnetics* 27 (2006) 280–287.
- [19] M. Roosli, M. Moser, Y. Baldinini, M. Meier, C. Braun-Fahrlander, Symptoms of ill health ascribed to electromagnetic field exposure—a questionnaire survey, *Int. J. Hyg. Environ. Health* 207 (2004) 141–150.
- [20] M. Roosli, Conference Poster at the WHO International Workshop on EMF Hypersensitivity, Prague, Czech Republic, October 25–27, 2004 [as reported in Rapporteur’s Report by KH Mild], 2004.
- [21] E. Lyskov, K.H. Mild, M. Sandstrom, Sensor Reactivity and Autonomous Regulation in Persons with Perceived Electrical Hypersensitivity, Conference Presentation at the WHO International Workshop on EMF Hypersensitivity, Prague, Czech Republic, October 25–27, 2004.
- [22] C.H. Mueller, C. Schierz, Project NEMESIS: double blind study on effects of 50 Hz EMF on sleep quality, physiological parameters and field perception in people suffering from electrical hypersensitivity, in: K.H. Mild, M. Repacholi, E. van Deventer, P. Ravazzani (Eds.), Proceedings of the International Workshop on EMF Hypersensitivity, Prague, Czech Republic, 2004, pp. 107–121.
- [23] E. Marková, L. Hillert, L. Malmgren, B.R. Persson, I.Y. Belyaev, Microwaves from GSM mobile telephones affect 53BP1 and gamma-H2AX foci in human lymphocytes from hypersensitive and healthy persons, *Environ. Health Perspect.* 113 (2005) 1172–1177.
- [24] I.Y. Belyaev, E. Marková, L. Hillert, L.O. Malmgren, B.R. Persson, Microwaves from UMTS/GSM mobile phones induce long-lasting inhibition of 53BP1/gamma-H2AX DNA repair foci in human lymphocytes, *Bioelectromagnetics* 30 (2009) 129–141.
- [25] A.P. Fews, D.L. Henshaw, P.A. Keitch, J.J. Close, R.J. Wilding, Increased exposure to pollutant aerosols under high voltage power lines, *Int. J. Radiat. Biol.* 75 (1999) 1505–1521.

- [26] A.P. Fews, D.L. Henshaw, R.J. Wilding, P.A. Keitch, Corona ions from powerlines and increased exposure to pollutant aerosols, *Int. J. Radiat. Biol.* 75 (1999) 1523–1531.
- [27] G. Nordström, *The Invisible Disease—The Dangers of Environmental Illnesses caused by Electromagnetic Fields and Chemical Emissions*, O Books, Hants and New York, 2004, ISBN 1-903816-71-8.
- [28] O. Johansson, P.-Y. Liu, "Electrosensitivity", "electrosuper-sensitivity" and "screen dermatitis": preliminary observations from on-going studies in the human skin, in: D. Simunic (Ed.), *Proceedings of the COST 244: Biomedical Effects of Electromagnetic Fields – Workshop on Electromagnetic Hypersensitivity*. Brussels/Graz: EU/EC (DG XIII), 1995, pp. 52–57.
- [29] X. Zhuang, A.J. Silverman, R. Silver, Distribution and local differentiation of mast cells in the parenchyma of the forebrain, *J. Comp. Neurol.* 408 (1999) 477–488.
- [30] O. Johansson, M. Hilliges, S.W. Han, A screening of skin changes, with special emphasis on neurochemical marker antibody evaluation, in patients claiming to suffer from screen dermatitis as compared to normal healthy controls, *Exp. Dermatol.* 5 (1996) 279–285.
- [31] S. Gangi, O. Johansson, Skin changes in "screen dermatitis" versus classical UV- and ionizing irradiation-related damage—similarities and differences. Two neuroscientists' speculative review, *Exp. Dermatol.* 6 (1997) 283–291.
- [32] S. Gangi, O. Johansson, A theoretical model based upon mast cells and histamine to explain the recently proclaimed sensitivity to electric and/or magnetic fields in humans, *Med. Hypotheses* 54 (2000) 663–671.
- [33] G. Holmboe, O. Johansson, Description of symptoms as well as occurrence of IgE and positive Phadiatop Combi in persons with the physical impairment electrohypersensitivity [Article in Swedish], *Medicinsk Access* 1 (2005) 58–63.
- [34] D. Leszczynski, S. Joensuu, J. Reivinen, R. Kuokka, Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: molecular mechanism for cancer- and blood-brain barrier-related effects, *Differentiation* 70 (2002) 120–129.
- [35] R. Nyland, D. Leszczynski, Mobile phone radiation causes changes in gene and protein expression in human endothelial cell lines and the response seems to be genome- and proteome-dependent, *Proteomics* 6 (2006) 4769–4780.
- [36] N. Wertheimer, E. Leeper, Electrical wiring configurations and childhood cancer, *Am. J. Epidemiol.* 109 (1979) 273–284.
- [37] L. Hardell, Å. Nilsson, A. Pahlson, A. Hallquist, K.H. Mild, Use of cellular telephones and the risk for brain tumours: a case-control study, *Int. J. Oncol.* 15 (1999) 113–116.
- [38] L. Hardell, K.H. Mild, M. Carlberg, A. Hallquist, Cellular and cordless telephone use and the association with brain tumors in different age groups, *Arch. Environ. Health* 59 (2004) 132–137.
- [39] L. Hardell, M. Carlberg, K.H. Mild, Case-control study on cellular and cordless telephones and the risk for acoustic neuroma or meningioma in patients diagnosed 2000–2003, *Neuroepidemiology* 25 (2005) 120–128.
- [40] S. Lonn, A. Ahlbom, P. Hall, M. Feychting, Mobile phone use and the risk of acoustic neuroma, *Epidemiology* 15 (2004) 653–659.
- [41] Ö. Hallberg, O. Johansson, Melanoma incidence and frequency modulation (FM) broadcasting, *Arch. Environ. Health* 57 (2002) 32–40.
- [42] Ö. Hallberg, O. Johansson, Malignant melanoma of the skin—not a sunshine story!, *Med. Sci. Monit.* 10 (2004) CR336–CR340.
- [43] Ö. Hallberg, O. Johansson, FM broadcasting exposure time and malignant melanoma incidence, *Electromagn. Biol. Med.* 24 (2005) 1–8.
- [44] R. Anane, M. Geffard, M. Taxile, D. Bodet, B. Billaudel, P.E. Dutou, B. Veyret, Effects of GSM-900 microwaves on the experimental allergic encephalomyelitis (EAE) rat model of multiple sclerosis, *Bioelectromagnetics* 24 (2003) 211–213.
- [45] P. Boscol, M.B. Di Scascio, S. D'Ostillo, A. Del Signore, M. Reale, P. Conti, P. Bavazzano, R. Paganelli, M. Di Gioacchino, Effects of electromagnetic fields produced by radiotelevision broadcasting stations on the immune system of women, *Sci. Total Environ.* 273 (2001) 1–10.
- [46] A. Dmoch, P. Moszczynski, Levels of immunoglobulin and subpopulations of T lymphocytes and NK cells in men occupationally exposed to microwave radiation in frequencies of 6–12 GHz [Article in Polish], *Med. Pr.* 49 (1998) 43–49.
- [47] P. Moszczynski, J. Lisiewicz, A. Dmoch, Z. Zabinski, L. Bergier, M. Rucinska, U. Sasiadek, The effect of various occupational exposures to microwave radiation on the concentrations of immunoglobulins and T lymphocyte subsets [Article in Polish], *Wiad. Lek.* 52 (1999) 30–34.
- [48] H. Tuschl, G. Neubauer, H. Garn, K. Duftschmid, N. Winker, H. Brusi, Occupational exposure to high frequency electromagnetic fields and its effect on human immune parameters, *Int. J. Occup. Med. Environ. Health* 12 (1999) 239–251.
- [49] A. Del Signore, P. Boscolo, S. Kouri, G. Di Martino, G. Giuliano, Combined effects of traffic and electromagnetic fields on the immune system of fertile atopic women, *Ind. Health* 38 (2000) 294–300.
- [50] S.F. Cleary, L.M. Liu, R.E. Merchant, In vitro lymphocyte proliferation induced by radio-frequency electromagnetic radiation under isothermal conditions, *Bioelectromagnetics* 11 (1990) 47–56.
- [51] S.F. Cleary, Z. Du, G. Cao, L.M. Liu, C. McCrady, Effect of isothermal radiofrequency radiation on cytolytic T lymphocytes, *FASEB J.* 10 (1996) 913–919.
- [52] E.M. Czerska, E.C. Elson, C.C. Davis, M.L. Swicord, P. Czeraki, Effects of continuous and pulsed 2450-MHz radiation on spontaneous lymphoblastoid transformation of human lymphocytes in vitro, *Bioelectromagnetics* 13 (1992) 247–259.
- [53] M.P. Dabrowski, W. Stankiewicz, R. Kubacki, E. Sobiczewska, S. Szmiigielski, Immunotropic effects in cultured human blood mononuclear cells pre-exposed to low-level 1300 MHz pulse-modulated microwave field, *Electromagn. Biol. Med.* 22 (2003) 1–13.
- [54] M.P. Dabrowski, W. Stankiewicz, E. Sobiczewska, S. Szmiigielski, Immunotropic effects of electromagnetic fields in the range of radio- and microwave frequencies [Article in Polish], *Pol. Merkuz. Lekarski* 11 (2001) 447–451.
- [55] W. Stankiewicz, M.P. Dabrowski, R. Kubacki, E. Sobiczewska, S. Szmiigielski, Immunotropic influence of 900 MHz microwave GSM signal on human blood immune cells activated in vitro, *Electromagn. Biol. Med.* 25 (2006) 45–51.
- [56] M. Donnellan, D.R. McKenzie, P.W. French, Effects of exposure to electromagnetic radiation at 835 MHz on growth, morphology and secretory characteristics of a mast cell analogue, RBL-2H3, *Cell. Biol. Int.* 21 (1997) 427–439.
- [57] C. Harvey, P.W. French, Effects on protein kinase C and gene expression in a human mast cell line, HMC-1, following microwave exposure, *Cell. Biol. Int.* 23 (2000) 739–748.
- [58] E. Elekcs, G. Thuroczy, L.D. Szabo, Effect on the immune system of mice exposed chronically to 50 Hz amplitude-modulated 2.45 GHz microwaves, *Bioelectromagnetics* 17 (1996) 246–248.
- [59] E.E. Fesenko, E.G. Novoselova, N.V. Semiletova, T.A. Agafonova, V.B. Sadovnikov, Stimulation of murine natural killer cells by weak electromagnetic waves in the centimeter range [Article in Russian], *Biofizika* 44 (1999) 737–741.
- [60] E.E. Fesenko, V.R. Makar, E.G. Novoselova, V.B. Sadovnikov, Microwaves and cellular immunity. I. Effect of whole body microwave irradiation on tumor necrosis factor production in mouse cells, *Bioelectrochem. Bioenerg.* 49 (1999) 29–35.
- [61] A.B. Gapeev, V.G. Safronova, N.K. Chemeris, E.E. Fesenko, Modification of the activity of murine peritoneal neutrophils upon exposure to millimeter waves at close and far distances from the emitter [Article in Russian], *Biofizika* 41 (1996) 205–219.
- [62] L. Gatta, R. Pinto, V. Ubaldi, L. Pace, P. Gallodi, G.A. Lovisolo, C. Marino, C. Pioli, Effects of in vivo exposure to GSM-modulated 900 MHz radiation on mouse peripheral lymphocytes, *Radiat. Res.* 160 (2003) 600–605.
- [63] M.P. Kolomytseva, A.B. Gapeev, V.B. Sadovnikov, N.K. Chemeris, Suppression of nonspecific resistance of the body under the effect of extremely high frequency electromagnetic radiation of low intensity [Article in Russian], *Biofizika* 47 (2002) 71–77.

- [64] K.V. Lushnikov, A.B. Gapeev, V.B. Sadovnikov, N.K. Cheremis, Effect of extremely high frequency electromagnetic radiation of low intensity on parameters of humoral immunity in healthy mice [Article in Russian], *Biofizika* 46 (2001) 753–760.
- [65] K.S. Nageswari, K.R. Sarma, V.S. Rajvanshi, R. Sharan, M. Sharma, V. Barathwal, V. Singh, Effect of chronic microwave radiation on T cell-mediated immunity in the rabbit, *Int. J. Biometeorol.* 35 (1991) 92–97.
- [66] F. Nasta, M.G. Prisco, R. Pinto, G.A. Lovisolo, C. Marino, C. Pioli, Effects of GSM-modulated radiofrequency electromagnetic fields on B-cell peripheral differentiation and antibody production, *Radiat. Res.* 165 (2006) 664–670.
- [67] E.T. Novoselova, E.E. Fesenko, Stimulation of production of tumor necrosis factor by murine macrophages when exposed in vivo and in vitro to weak electromagnetic waves in the centimeter range [Article in Russian], *Biofizika* 43 (1998) 1132–1133.
- [68] E.O. Novoselova, E.E. Fesenko, V.R. Makar, V.B. Sadovnikov, Microwaves and cellular immunity. II. Immunostimulating effects of microwaves and naturally occurring antioxidant nutrients, *Bioelectrochem. Bioenerg.* 49 (1999) 37–41.
- [69] N. Çetin, A. Bilgili, G. Eraslan, Effects of pulsed magnetic field chronic exposure on some hematological parameters in mice, *Revue Méd. Vét.* 157 (2006) 68–71.
- [70] K.I. Obukhan, The effect of ultrahigh-frequency radiation on adaptation thresholds and the damages to blood system cells [Article in Ukrainian], *Lik. Sprava* 7 (1998) 71–73.
- [71] L. Stronati, A. Testa, J. Moquet, A. Edwards, E. Cordelli, P. Villani, C. Marino, A.M. Freseghna, M. Appolloni, D. Lloyd, 935 MHz cellular phone radiation. An in vitro study of genotoxicity in human lymphocytes, *Int. J. Radiat. Biol.* 82 (2006) 339–346.
- [72] J.L. Phillips, O. Ivaschuk, T. Ishida-Jones, R.A. Jones, M. Campbell-Beachler, W. Haggren, DNA damage in Molt-4 T-lymphoblastoid cells exposed to cellular telephone radiofrequency fields in vitro, *Bioelectrochem. Bioenerg.* 45 (1998) 103–110.
- [73] S. Ivancsits, E. Diem, A. Pilger, H.W. Rüdiger, O. Jahn, Induction of DNA strand breaks by intermittent exposure to extremely-low-frequency electromagnetic fields in human diploid fibroblasts, *Mutat. Res.* 519 (2002) 1–13.
- [74] H. Lai, N.P. Singh, Acute low-intensity microwave exposure increases DNA single-strand breaks in rat brain cells, *Bioelectromagnetics* 16 (1995) 207–210.
- [75] H. Tuschl, W. Novak, H. Molla-Djafari, In vitro effects of GSM modulated radiofrequency fields on human immune cells, *Bioelectromagnetics* 27 (2006) 188–196.
- [76] J.L. Chagnaud, B. Veyret, In vivo exposure of rats to GSM-modulated microwaves: flow cytometry analysis of lymphocyte subpopulations and of mitogen stimulation, *Int. J. Radiat. Biol.* 75 (1999) 111–113.
- [77] B. Veyret, C. Bouthet, P. Deschaux, R. de Seze, M. Geffard, J. Jousset-Dubien, M. le Diraison, J.M. Moreau, A. Caristan, Antibody responses of mice exposed to low-power microwaves under combined, pulse-and-amplitude modulation, *Bioelectromagnetics* 12 (1991) 47–56.
- [78] M. Capri, S. Salviolo, S. Altília, F. Sevinci, D. Remondini, P. Mesirca, F. Bersani, D. Monti, C. Franceschi, Age-dependent effects of in vitro radiofrequency exposure (mobile phone) on CD95+ T Helper human lymphocytes, *Ann. N. Y. Acad. Sci.* 1067 (2006) 493–499.
- [79] A.I. Yurekli, M. Ozkan, T. Kalkan, H. Saybasili, H. Tuncel, P. Atukeren, K. Gumustas, S. Seker, GSM base station electromagnetic radiation and oxidative stress in rats, *Electromagn. Biol. Med.* 25 (2006) 177–188.
- [80] C. Schwarz, E. Kratochvil, A. Pilger, N. Kuster, F. Adlkofer, H.W. Rüdiger, Radiofrequency electromagnetic fields (UMTS 1 950 MHz) induce genotoxic effects in vitro in human fibroblasts but not in lymphocytes, *Int. Arch. Occup. Environ. Health* 81 (2008) 755–767.
- [81] M. Simkó, M.O. Mattsson, Extremely low frequency electromagnetic fields as effectors of cellular responses in vitro: possible immune cell activation, *J. Cell. Biochem.* 93 (2004) 83–92.
- [82] H. Nakamura, T. Seto, H. Nagase, M. Yoshida, S. Dan, K. Ogino, Effects of exposure to microwaves on cellular immunity and placental steroids in pregnant rats, *Occup. Environ. Med.* 54 (1997) 676–680.
- [83] H. Nakamura, T. Seto, K. Hatta, I. Matsuzaki, H. Nagase, M. Yoshida, K. Ogino, Natural killer cell activity reduced by microwave exposure during pregnancy is mediated by opioid systems, *Environ. Res.* 79 (1998) 106–113.
- [84] H. Nakamura, H. Nagase, K. Ogino, K. Hatta, I. Matsuzaki, Uteroplacental circulatory disturbance mediated by prostaglandin F(2alpha) in rats exposed to microwaves, *Reprod. Toxicol.* 14 (2000) 235–240.
- [85] J.N. Magras, T.D. Xenos, RF radiation-induced changes in the prenatal development of mice, *Bioelectromagnetics* 18 (1997) 455–461.
- [86] F. Vecchio, C. Babiloni, F. Ferreri, G. Curcio, R. Fini, C. Del Percio, P.M. Rossini, Mobile phone emission modulates interhemispheric functional coupling of EEG alpha rhythms, *Eur. J. Neurosci.* 25 (2007) 1908–1913.
- [87] M. Seishima, Z. Oyama, M. Oda, Cellular phone dermatitis with chromate allergy, *Dermatology* 207 (2003) 48–50.
- [88] D. Roux, A. Vian, S. Girard, P. Bonnet, F. Paladian, E. Davies, G. Ledoigt, High frequency (900 MHz) low amplitude (5 V/m) electromagnetic field: a genuine environmental stimulus that affects transcription, translation, calcium and energy charge in tomato, *Planta* 227 (2008) 883–891.
- [89] H.A. Divan, L. Kheifets, C. Obel, J. Olsen, Prenatal and postnatal exposure to cell phone use and behavioral problems in children, *Epidemiology* 19 (2008) 523–529.
- [90] J. Friedman, S. Kraus, Y. Hauptman, Y. Schiff, R. Seger, Mechanism of short-term ERK activation by electromagnetic fields at mobile phone frequencies, *Biochem. J.* 405 (2007) 559–568.
- [91] D.J. Panagopoulos, E.D. Chavdoula, I.P. Nezis, L.H. Margaritis, Cell death induced by GSM 900-MHz and DCS 1800-MHz mobile telephony radiation, *Mutat. Res.* 626 (2007) 69–78.
- [92] D.J. Panagopoulos, L.H. Margaritis, Effects of electromagnetic fields on the reproductive capacity of *D. melanogaster*, in: P. Stavroulakis (Ed.), *Biological Effects of Electromagnetic Fields*, Springer, Berlin/New York, 2003, pp. 543–578.
- [93] D.J. Panagopoulos, N. Messini, A. Karabarbounis, A.L. Philippetis, L.H. Margaritis, Radio frequency electromagnetic radiation within “safety levels” alters the physiological function of insects, in: P. Kostarakis, P. Stavroulakis (Eds.), *Proceedings of the Millennium International Workshop on Biological Effects of Electromagnetic Fields*, Heraklion, Crete, Greece, 2000, pp. 169–175.
- [94] D.J. Panagopoulos, A. Karabarbounis, L.H. Margaritis, Effect of GSM 900-MHz mobile phone radiation on the reproductive capacity of *D. melanogaster*, *Electromagn. Biol. Med.* 23 (2004) 29–43.

Reproductive and developmental effects of EMF in vertebrate animal models

Aris F. Pourlis *

Laboratory of Anatomy Histology & Embryology, Veterinary School, University of Thessaly Karditsa, Greece

Received 10 August 2008; received in revised form 28 August 2008; accepted 30 January 2009

Abstract

This paper reviews the literature data on the effects of electromagnetic fields (EMF), in the reproductive organs as well as in prenatal and postnatal development of vertebrate animals. Review articles which have been published till 2001, regarding the reproductive and developmental effects of the entire range of frequency of electromagnetic fields, were surveyed. Experimental studies which were published from 2001 onwards were summarized. Special focus on the effects of radiofrequencies related to mobile communication in the above mentioned topics has been made. According to the majority of the investigations, no strong effects resulted regarding the exposure to EMF of mobile telephony in the animal reproduction and development. However further research should be done in order to clarify many unknown aspects of the impact of EMF in the living organisms.

© 2009 Elsevier Ireland Ltd. All rights reserved.

Keywords: Electromagnetic fields (EMF); Mobile phones; Teratology; Endometrium; Testis

1. Introduction

During the 20th century, the exposure to electromagnetic fields (EMF) became an important source of concern about the possible effects in the living organisms. The artificial sources of electromagnetic radiation have risen tremendously because of the ongoing needs on electricity, telecommunications, and electronic devices. In this context, World Health Organisation (WHO) established in 1996 the International EMF project in order to assess health and environmental effects of exposure to EMF in the frequency range from 0 to 300 GHz. For the purpose of this paper this range will be divided into static (0 Hz), extremely low frequency (ELF > 0–300 kHz), intermediate frequencies (IF > 300–10 MHz) and radiofrequency (RF 10 MHz–300 GHz) fields [J. Juutilainen, Developmental effects of electromagnetic fields, *Bioelectromagnetics* 7 (2005) S107–S115]. The mobile phone technology is based on radiofrequency radiation with transmission of microwaves carrying frequencies between 880 and 1800 MHz [P.A. Valberg, T.E. van Deventer, M.H. Repacholi, Workgroup report:

base stations and wireless networks-radiofrequency (RF) exposures and health consequences, *Environ. Health Perspect.* 115 (2007) 416–424].

The mobile telephony revolution took place in the last decade. There is an increasing number of cell phone users all over the world. Also, new technologies which use the spectrum of high frequency emissions are incorporated in many aspects of telecommunications. As a consequence, there is a lot of interest about the possible effects of the radiation emitted from the machines which are engaged in the telephony such as hand phones, base stations and transmitters.

The biological effects of EMF have been and are being investigated on different levels of organization. On the level of human populations, epidemiological studies are used whereas, on the level of individuals human, animal and plant *in vivo* experiments are carried out. Furthermore, on the level of organs, tissues and cells *in vitro* investigations are employed. Finally, on the sub-cellular level, biochemical and molecular techniques are utilized.

From another point of view, many studies have been carried out or are in progress about the various effects of radiation emissions regarding the behaviour, cancer, central nervous system, sleep, children, cardiovascular system, immune function, reproduction and development [3].

* Tel.: +30 24410 66013.

E-mail address: apourlis@vet.uth.gr.

The present paper will focus on the existing data about the reproductive and developmental effects of EMF in vertebrates. Reproduction is a critical function of the organisms and involves two body systems the male and female genital system. The development comprises a series of events which begins with fertilization, continues with implantation, embryonic growth and terms with sexual maturity. In the context of systematic zoology, the vertebrates are close to the humans. Therefore, the animal studies could provide useful information on the comprehension of interaction of EMF with the living organism and on the possible commonality with the humans.

The biological effects of EMF of interest can be broadly grouped into thermal and non-thermal [4]. The thermal effects are associated with local heat production just like the mechanism of a microwave oven. The non-thermal mechanism is triggered by an amount of energy absorption, which is not directly associated with temperature change but rather to some other changes produced in the tissues.

The goal of this paper is to present the up to date available data about the EMF and their potential effects on reproduction and development, filling the gap of information from the most recent published reviews. All the bibliographic data which will be presented were collected exclusively from scientific journals published in English and partially in other languages. The survey includes studies which were published from 2001 onward. The studies which relate to the impact of mobile phone electromagnetic fields will be presented thoroughly and independently from the date of their publication.

2. Historical background

The first paper which I found in the medical literature, regarding the effects of EMF on the development of vertebrates, was published in 1893 in an anatomical journal from Windle [5]. The author summarized the observations of three scientists and added his own about the effects of electricity on the chicken embryos. Two years later the same author [6], published an account on the effects of electricity and magnetism on development.

In 1980 two papers were published about the biological effects of microwave radiation. Cook et al. [7] published a comprehensive survey regarding the very early research on the biological effects of electromagnetic fields. The early work on short waves from 1885 to 1940 was presented. Following, the authors summarized the available data from 1940 to 1960. Leach [8] provided an account on the genetic, growth and reproductive effects of microwave radiation including early studies in this field that were published from 1959 to 1979. The majority of revised papers dealt with animals. Later, Algers and Hennichs [9] summarized the biological effects on vertebrates, of electromagnetic fields where the frequency did not exceed 100 Hz. The authors included many studies about the impact of EMF on farm animals. The same

year, a specialized review was published on the effects of non-ionizing radiation on birds [10].

Berman et al. [11], presented the results of a large multinational experimental effort (Henhouse project) regarding the low frequency EMF effects on chick embryos. Juutilainen [12], Chernoff et al. [13], Brent et al. [14] presented detailed reviews of the literature about the effects on reproduction related to low frequency EMF.

Jensh [15] reviewed behavioral teratologic studies using microwave radiation with special interest to continuous wave (CW) 915, 2450, or 6000 MHz radiation.

Verschaeve and Maes [16] reviewed the genetic, carcinogenic and teratogenic effects of RF (300 MHz–300 GHz). Regarding the effects on reproduction and teratogenesis, studies from 1961 to 1991 were surveyed. The majority of these experimental studies dealt with the exposure of animals at 2.45 GHz. The same year, Huuskonen et al. [17] reported on the teratogenic and reproductive effects of low frequency (0–100 kHz) magnetic fields associated with the use or transmission of electric power or emitted from video display terminals. The animal studies that were surveyed, have been published from 1987 to 1997 regarding the effects of alternating magnetic fields on prenatal development of rats and mice. In the same paper, studies on the effects of prenatal exposure of alternating magnetic fields on postnatal development were included. Brent [18] provided a thorough review of *in vivo* and *in vitro* studies on the reproductive and teratologic effects of low frequency EMF. The survey of reproductive effects has involved studies with chick embryos, chickens, cows, mice, and rats from 1969 to 1996. O'Connor [19] recorded the intrauterine effects in animals exposed to radiofrequency and microwave fields with a special feature. The SAR of the surveyed studies was above the limit of 0.4 W/kg.

Experimental studies on the teratologic effects or developmental abnormalities from exposure to RF electromagnetic fields in the range 3 kHz–300 GHz were reviewed from Heynick and Merritt [20]. The review included investigations with insects, birds (chicken, quails, turkeys) and mammalian species (mice, rats) as well as non-human primates which appeared from 1974 to 2000. A brief critical review, on the developmental effects of extremely low frequency (ELF) electric and magnetic fields provided by Juutilainen [21]. Löscher [22] published a survey of the effects of radiofrequency electromagnetic fields on production, health and behaviour of farm animals.

Juutilainen [1] reported on the effects of EMF on animal development. In his review, he surveyed specific topics such as the Henhouse project, the interaction of LF-IMF EMF with known teratogens, and the behavioral teratology of RF. Saunders and McCaig [23] summarized the possible effects on prenatal development of physiologically weak electric fields induced in the body by exposure to extremely low frequency electromagnetic fields and of elevated temperature levels that might result from exposure to radiofrequency (RF) radiation.

Table 1
Overview of investigations on EMF effects on animal genital system.

Animal species	Exposure frequency	Exposure parameters	Exposure duration	Endpoint	Results	Comments	Reference
Mouse Swiss	50 Hz	25 mT	Continuous 90 days	Effects on reproductive ability	No effect on the fertility of male and female mice. The ovarian weight was significantly increased		[27]
Mouse CD1 (BALB/c X DBA/2)	60 Hz	2 mT	Continuous for 72 h or 8 h/day for 10 days	Sperm morphology	No statistically differences were observed	Two groups were treated with mitomycin C. Sperm abnormalities were found in the group exposed versus the group treated with mitomycin C alone	[28]
Mouse BALB/c	60 Hz	0.1 or 0.5 mT	24 h/day for 8 weeks	Germ cell apoptosis in the testes	No significant changes in testicular weights. Decrease of normal seminiferous tubules. Increase of the germ cell death		[29]
Rat Sprague-Dawley	60 Hz	5, 83.3, 500 mT	Continuous 21 h/day from day 6 of gestation to day 21 of lactation	Spermatotoxicity and reproductive dysfunction in the F1 offspring	No detectable alterations in offspring spermatogenesis and fertility		[30]
Rat Sprague-Dawley	50 Hz	25 ± 1 µT	Continuous for 18 weeks	Effects on sperm count, weights of testes, seminal vesicles, preputial glands	No effect on the weight of testes. Significant reduction of the weight of seminal vesicles and preputial glands. Significant reduction in sperm count		[31]
Rat Sprague-Dawley	50 Hz	1.35 ± 0.018 mT	2 h/day, 7 days/week for 2 months	Sperm count, morphological changes of testes	No significant alterations were observed	Funding not mentioned	[32]
Rat Wistar albino ♂	50 Hz	1 mT (mean value)	3 h/day for 50 or 100 days	Morphological evaluation of uterus and ovaries	Ultrastructural alterations in germinal epithelium of ovaries in the experimental group (50 days) as well as in tunica albuginea (100 days)	Ambiguous observations in the uterus	[33]
Rat Sprague-Dawley ♂	20 kHz	6.25 mT	8 h/day, 5 days/week for 90 days	Histopathological examination of various organs	No differences were seen in testis and ovary		[34]

Table 1 (Continued)

Animal species	Exposure frequency	Exposure parameters	Exposure duration	Endpoint	Results	Comments	Reference
Rat Wistar ♂♀	50 Hz		3 weeks in utero and 5 weeks postnatal	Testes	Morphological changes in the seminiferous tubules		[35]
Rat Sprague-Dawley ♂♂	20 kHz sine waves	6.25 mT	8 h/day for 12 or 18 months	Histopathological examination of various organs	No differences were seen in testis and ovary		[36]
Rat Wistar	30–300 GHz	>0.3 mW/cm ²	30 min for 63 days	Spermatogenesis	Morphological changes in spermatozoa	Scanty data presentation	[37]
Rat Wistar	50 Hz		8 h/day for 8 months	Histological evaluation of testes	Mean seminiferous tubule diameter and testicular weight were significantly lower in exposed group. Histologic damage score was threefold in experimental group versus control		[38]

A special topic, the effects of EMF from power lines on avian reproductive biology, was reviewed by Femie and Reynolds [24]. Krewski et al. [25], reviewed studies referring to various disciplines regarding the effects of RF. The included literature was published between 2001 and 2003. A novelty of this paper, was a discussion of the reports of various authorities and committees about the potential health risks associated with exposure to RF fields. A gap in the literature regarding the biological effects of EMF in the intermediate frequency range was covered by the review of Shigemitsu et al. [26].

During the last decade, many reports from authorities (local, national and international) and expert panels have been uploaded on the web [2].

It is suggested that the reader refer to the above-mentioned review articles and electronic addresses, in order to assemble a more complete and detailed view of the biological effects of EMF.

3. Male genital system

The testes are very important organs situated externally to the body and enclosed by the scrotum. The testicular parenchyma is the site of an intense proliferation and differentiation of the germinal cells that will become the sperm cells. The testes are very sensitive to temperature variations and for this reason the scrotum, which contains the testicular parenchyma, has a specialized contractile structure.

Studies that have evaluated EMF effects (mainly LF) on the genital systems of the vertebrates are summarized in Table 1.

Regarding mobile telephony, the first study conducted by Dasdag et al. [39] investigated whether there are adverse effects due to microwave exposure emitted by cellular phones in male Wistar albino rats. The animals ($n = 18$) were divided in three groups (control, standby exposed group, speech exposed group). Specific energy absorption rate (SAR) was 0.141 W/kg. Rats in the experimental groups were exposed for 2 h/day for 1 month in standby position, whereas phones were turned to the speech position three times for 1 min. The decrease of epididymal sperm counts in the speech groups was not found to be significant. Differences in terms of normal and abnormal sperm forms were not observed. Histological changes were especially observed in the testes of rats in the speech group. Seminiferous tubular diameter of rat testes in the standby and speech groups was found to be lower than the sham group. Rectal temperatures of rats in the speech group were found to be higher than the sham and standby groups. The rectal temperatures of rats before and after exposure were also found to be significantly higher in the speech group.

The same group of authors [40], failed to reproduce the results of their previous work. Sixteen Sprague-Dawley rats were separated into two groups (control, experimental). They were exposed to 890–915 MHz pulsed wave (PW) daily for

20 min/day for 1 month. For 250 mW average radiated power, SAR was 0.52 W/kg. No differences were observed in the percentages of epididymal normal and abnormal sperms, the epididymal sperm count, as well as in the seminiferous tubule diameter between control and experimental groups. Also, the testicular biopsy score as evaluated by Johnson's scale did not differ significantly.

Aitken et al. [41] assessed the testis of mice irradiated with 900 MHz in a waveguide, with an exposure condition SAR 90 mW/kg for 7 days at 12 h/day. The authors did not observe abnormalities regarding the sperm number, morphology and vitality. However, they reported significant damage to the mitochondrial genome as well as to the nuclear-globin locus.

Results similar to a previous study [39] regarding the diameter of the seminiferous tubules of rat testes were obtained by Ozguner et al. [42]. During the experiment, 20 male Sprague–Dawley rats (5 months of age) were either exposed to 900 MHz CW (average power density 1 ± 0.4 mW/cm²) or not (control group). Rats exposed 30 min/day, for 5 days/week for 4 weeks. The authors also did not observe significantly different values of weight of testes, testicular biopsy score count and the percentage of interstitial tissue. However, the mean height of the germinal epithelium was found decreased in the group of rats that had been irradiated.

Forgács et al. [43] repeatedly exposed male NMR1 mice to 1800 MHz GSM like microwave radiation at 0.018–0.023 W/kg whole body SAR. 11–12 sham exposed and 11–12 exposed mice were used. The animals were exposed ten times (over 2 weeks) and the duration of exposure was 2 h/day. No microwave exposure-related morphological alterations were found in testis, epididymis and prostate.

Adult male rats were examined after exposure at sub-chronic exposure to RF emitted from a conventional cell phone on their testicular function. Sixteen Wistar rats were used at age 30 days. The animals were exposed for 1 h daily during 11 weeks. The experimental group ($n = 8$) was exposed to 1835–1850 MHz at 0.04–1.4 mW/cm². Total body weight and absolute and relative testicular and epididymal weights did not change significantly. Epididymal sperm count was not significantly different between the groups. Regarding the histomorphological endpoints of the study, no difference was found between the experimental and control arm [44].

The effect of cellular phone emissions on sperm characteristics in 16 Sprague–Dawley rats were studied [45]. The laboratory animals were divided in two groups (experimental, control) and exposed to four cell phones which had a personal communications service code division multiple access frequency band of 1.9 GHz (800 MHz digital and 800 MHz analog). The rats received daily (3 h–30 min rest–3 h) cell phone exposure for 18 weeks. The SAR ranged from 0.9 to 1.80 W/kg whereas the power from 0.00001 to 0.607 W, according to the specific mode of function. The authors analyzed the morphology of the sperm cells from

epididymis of rats. The percentage of deformities for the experimental group was 34.3% and the percentage of deformities for the control group was 32.1%. This difference in the occurrence of deformities between the two groups was not statistically significant ($p > .05$) through a paired *t* test. The total sperm counts from the testes were not significantly different between the two groups. None of the temperature differences between the two groups were statistically significant.

Sixteen Sprague–Dawley rats were used to evaluate the bcl-2 protein (an anti-apoptotic protein) in rat testes. The experimental group ($n = 8$) was exposed to commercial (GSM) cellular phones irradiation for 20 min/day for 1 month. Average power density was measured at 0.047 mW/cm² and SAR levels changed between 0.29 and 0.87 W/kg. The testes were investigated by means of immunohistochemistry. No difference was observed between testes sections of the sham and experimental groups in terms of bcl-2 staining. These results indicate that the radiation emitted from 900 MHz cellular phones did not alter the anti-apoptotic protein in the testes of rats [46].

In order to investigate the apoptosis-inducing effect of mobile phone exposure on spermatogonia in seminiferous tubules, 31 Wistar albino male rats were divided in three groups such as cage control ($n = 10$), sham exposed ($n = 7$), and experimental ($n = 14$). The 2 h/day (7 days/week) exposure of 900 MHz radiation (power density 0.012–0.149 mW/cm² and SAR 0.07–0.57 W/kg) over a period of 10 months was evaluated by means of immunohistochemistry. The long-term radiation did not affect the active caspase-3 levels in testes of rats. Caspase-3 is a typical feature of apoptosis [47].

4. Female genital system

Studies on the impact of RF in the female genital system are scarce. Two studies were conducted in order to evaluate the effects on endometrial apoptosis and the ameliorating effects of a combination of vitamin E and C against EMF damage.

Oral et al. [48], exposed sexually mature female rats (16 weeks old) to 900 MHz radiation, 30 min/day for 30 days. Twenty-four Wistar albino rats were divided in three groups (sham exposed, EMF exposed, EMF exposed treated with vitamin C and E). The animals were exposed at 1.04 mW/cm² (SAR 0.016–4 W/kg). The effect of microwaves was examined in rat endometrium by means of immunohistochemistry. Endometrial apoptosis was observed. Guney et al. [49], repeated the experiment with the addition of another group (control). Histological changes in endometrium, diffuse and severe apoptosis in the endometrial surface, epithelial and glandular cells were reported regarding the group exposed to EMF. Also, eosinophilic leucocyte and lymphocyte infiltration were seen in the endometrial stroma.

Table 2
Overview of investigations on EMF effects on animal development.

Animal species	Exposure frequency	Exposure parameters	Exposure duration	Endpoint	Results	Comments	Reference
Rat Sprague–Dawley	50 Hz	7, 70, 350 mT	22 h/day during 0–7 or 8–15 day of gestation	Effects on teratogenicity and embryonic development	No differences regarding embryonic deaths, fetal weight and teratogenicity		[50]
Mouse ICR	50 Hz	Sham (0.1–1 μ T), 0.5, 5 mT	9 weeks [♂] 2 weeks [♀] prior to mating	Effects on teratogenicity and embryonic development	No differences regarding embryonic deaths, fetal weight and teratogenicity		[51]
Mouse Swiss Webster	0 Hz–25 MHz		1 week beginning from the 18th day of pregnancy	Morphological changes in brain, thymus, adrenal gland during embryonic development	Pathological changes were observed in the neonates		[52]
Rat Sprague–Dawley	60 Hz	0 (sham group), 5, 83.3, 500 mT	22 h/day during 6–20 day of gestation	Developmental toxicity	No differences regarding embryonic deaths, fetal weight and teratogenicity		[53]
Chicken	50 Hz	1.33–7.32 mT	24 h	Effects on teratogenicity and embryonic development	Significant difference in the percentage of abnormal embryos versus control was found in 4.19, 5.32, 5.86, and 6.65 densities. Some embryos with extra ribs, defects in ribs and vertebrae, anuria and abnormal beaks were observed	Funding not mentioned	[54]
Mouse ICR	20 kHz	6.5 mT	8 h/day from 2.5 to 15.5 days post-coitum	Effects on teratogenicity and embryonic development	No statistically significant differences in the number of implantation, embryonic death, resorption, growth retarded fetuses, external and skeletal abnormalities		[55]
Chicken Leghorn HR7	50 Hz	1 μ T, 500 μ T, 1 mT	Continuous for 15 or 21 days	Effects on embryo/fetus	At 15 days of incubation body weight was significantly lower versus controls. At 21 days of incubation the body weight and cranial diameters were smaller versus control. No differences in brain weight were observed in all groups	Funding not mentioned	[56]
Mouse ♀	Static magnetic field	400 mT	6 min/day from 7.5 to 14.5 day of pregnancy	Teratogenic effects	Polydactylism, abdominal fissure, fused ribs, vestigial 13th rib, brain hernia, curled tail		[57]
Mouse ♀	50 Hz	1.2 mT	8 h/day during pregnancy	Body weight of dams, development of offspring	Fetal loss, malformed fetuses, retardation of growth of the offspring in the first 2 weeks after birth	Article in chinese	[58]
Chicken White Leghorn eggs	50 Hz	1.33–7.32 mT	4 days	Morphological evaluation of embryos/fetuses	Abnormal brain cavities, spina bifida, monophthalmia, ocular defects and growth retardation		[59]

5. Developmental effects

The critical phases in the dynamic process of development take place mainly *in utero* (mammals) or *in ovo* (birds) i.e. during the embryonic period. The main bulk of investigations were performed regarding the possible effects on animals after irradiation, during *in utero* or *in ovo* development. The effects on development are determined by endpoints such as weight gain, congenital malformations, resorptions, and number of litters. These endpoints will be considered for various exposure conditions. The effects of EMF (mainly LF) on animal development are summarized in Table 2. Egg production was reduced (8%) when young laying hens have been continuously exposed to CW 915 MHz with an incident power of 800 mW during the first 2.5 weeks, 0 mW during the following week and 200 mW for the rest of experiment. Hatching of fertile and total eggs was not significantly influenced. No macroscopic malformations were observed in the chicks or dead embryos [60].

Jensh et al. [61] irradiated pregnant Wistar albino rats at a power density level of 10 mW/cm², at a frequency of 915 MHz and average SAR 3.57 W/kg. The animals were exposed for 6 h/day from day 1 to day 21 of gestation. No significant teratogenic signs were observed regarding the resorption rate, malformation rate, mean litter size, fetal weight and number of live and dead fetuses. The experiment was repeated and extended in order to analyze the embryonic and postnatal development of offspring [62]. Eleven pregnant rats were irradiated and 19 rats were used as control animals. All animals delivered and raised their offspring (F_{1a}) until weaning at 30 days of age. Ten days later females were rebred and teratologic evaluation was conducted on the resultant F_{1b} fetuses. At 90 days of age, reproductive capability was evaluated and a standard teratologic analysis performed on the resultant F₂ offspring. No significant morphologic changes were revealed.

Pregnant rats were exposed at 970 MHz for 22 h/day from the 1st to 19th day of pregnancy [63]. The SAR values varied from 0.07, 2.4 and 4.8 W/kg. The embryo mortality, fetal weight, skeletal ossification, as well as maternal fertility were evaluated. The exposure with SAR 4.8 W/kg caused reduced (–12%) fetal body weight versus the control. All the other examined parameters were not significantly different.

Klug et al. [64] exposed rat embryos (9.5 days old) for up to 36 h to 900 MHz. The modulation frequency was fixed at 215 Hz and the SAR values were calculated at 0.2, 1 and 5 W/kg. The endpoints of the experiment were crown-rump length, number of somites as well as embryonic malformations. No significant changes were observed on the growth and differentiation parameters of the embryos. Magras and Xenos [65] investigated the possible effects of radiofrequency radiation on prenatal development in mice. The study consisted of *in vivo* experiments at several places around an "antenna park" where the frequency emissions ranged from 88.5 to 950 MHz. At these locations RF power densities between 168 and 1053 nW/cm² were measured. Twelve pairs

of mice, divided in two groups, were placed in locations of different power densities and were repeatedly mated five times. One hundred eighteen newborns were collected. They were measured, weighed, and examined macro- and microscopically. A progressive decrease in the number of newborns per dam was observed, which ended in irreversible infertility. The prenatal development of the newborns, however, evaluated by the crown-rump length, the body weight, and the number of the lumbar, sacral, and coccygeal vertebrae, was improved. Wistar albino rats [15] were exposed through pregnancy for 6 h each day to CW 915 MHz radiation at a power density level of 10 mW/cm². Teratologic evaluation included the following parameters: mean litter size, maternal organ weight and organ weight/body weight ratios, body weight ratios of various organs (brain, liver, kidneys, and ovaries), number of resorptions and resorption rate, number of abnormalities and abnormality rate, mean term fetal weight. Mothers were rebred, and the second, unexposed litters were evaluated for teratogenic effects. Animals exposed to 915 MHz did not exhibit any consistent significant alterations in any of the above parameters.

Wistar rats were continuously exposed [66] during pregnancy to a low-level (0.1 mW/cm²) 900 MHz, 217 Hz pulse modulated EMF. Whole body average SAR values for the freely roaming, pregnant animals were measured in models; they ranged between 17.5 and 75 mW/kg. No differences between exposed and sham exposed dams or offspring were recorded in terms of litter size, evolution of body mass and developmental landmarks of litter mates. The effects of microwaves emitted by cellular phones on birth weights of rats were investigated by Dasdag et al. [67]. Thirty-six Wistar albino rats were divided into four groups. Each experimental or sham exposed group comprised six males or 12 females. The rats were exposed at 890–915 MHz (SAR 0.155 W/kg). Males were exposed daily for 3 × 1 min during 2 h/day for 1 month. Females were exposed in the same way until they gave birth. When the offspring became adult the experiment was repeated on them. No significant differences were observed between rectal temperatures in the sham and experimental groups. The birth weight of offspring in the experimental group was significantly lower than in the sham exposed group. However in the next generation of rats all parameters investigated were normal. Pregnant Sprague–Dawley rats were exposed [68] to ultra wide band (UWB) 0.1–1 GHz radiation in order to determine if teratological changes occur in rat pups as a result of (1) daily UWB exposures during gestation days 3 ± 18, or (2) as a result of both prenatal and postnatal (10 days) exposures. Dams were exposed either to (I) UWB irradiation with average whole body specific absorption rate 45 mW/kg (II) sham irradiation or (III) a positive control. Offspring were examined regarding litter size, sex-ratios, weights, coat appearance, and tooth eruption. The pups postnatally exposed were examined for hippocampal morphology. Generally, no significant differences were found between the exposed and sham group. The medial-to-lateral length of the hippocampus was significantly longer in the

Table 3
Summary of animal studies on effects of EMF (related to mobile telephony), on reproduction and development.

Animal species	Exposure frequency	Endpoint	Effect	Reference
Chicken	915 MHz	Development	No	[60]
Rat	915 MHz	Development	No	[61]
Rat	915 MHz	Development	No	[62]
Rat	970 MHz	Development	No	[63]
Rat	915 MHz	Development	No	[15]
Rat	900 MHz	Development	No	[64]
Mouse	88.5–950 MHz	Fertility/development	Yes/no	[65]
Rat	890–915 MHz	Testes	Yes	[39]
Rat	900 MHz	Development	No	[66]
Rat	0.1–1 GHz	Development	No	[68]
Rat	890–915 MHz	Development	Yes	[67]
Chicken	900 MHz	Development	Yes	[69]
Rat	890–915 MHz	Testes	No	[40]
Chicken		Development	Yes	[70]
Rat	900 MHz	Testes	No	[42]
Mouse	900 MHz	Testes	No	[41]
White stork	900–1800 MHz phone mast	Reproduction	Yes	[74]
Chicken	900 MHz	Kidney development	Yes	[71]
Mouse	1800 MHz	Testes	No	[43]
Rat	900 MHz	Endometrium	Yes	[48]
Rat	900 MHz	Brain development	No	[72]
Rat	1835–1850 MHz	Testes	No	[44]
Rat	1.9 GHz	Sperm	No	[45]
Tit	1200–3000 MHz	Reproduction	No	[75]
Rat	900 MHz	Endometrium	Yes	[49]
Chicken	900 MHz	Development	Yes	[73]
Rat	900 MHz	Testes	No	[46]
Rat	900 MHz	Testes	No	[47]

UWB-exposed pups than in the sham exposed animals but could not correlated with neurological dysfunction. The male offspring exposed *in utero* to UWB mated significantly less frequently than sham exposed males, but when they did mate there was no difference in fertilization and offspring numbers from the sham group.

Bastide et al. [69] reported chicken embryo mortality from day 7 to day 11 of incubation. This mortality reached 64% compared to 11% in controls. The maximum level of embryonic mortality was observed in the eggs placed near the telephone.

Chicken embryos were exposed to EMF from GSM mobile phone during the embryonic development [70]. The embryo mortality rate in the incubation period increased to 75% versus 16% in control group.

Ingole and Ghosh [71] studied by means of light microscopy the developmental effects on the avian kidney of radiation, from a cell phone handset (900 MHz frequency, power of 2 W and SAR of 0.37 W/kg). The authors reported morphological alterations on the epithelium of the renal tubules as well as of the renal corpuscles in E6, E8 and E10 chicken embryos.

The possible impact of cell phone radiation in the developing central nervous system of male Wistar rats was examined [72]. The animals were exposed to 900 MHz signal for 2 h/day on 5 days/week. After 5 weeks of exposure at whole body average SAR of 0.3 or 3 W/kg or sham exposure no degenerative morphological changes were found.

The results about the effects of exposing fertilized chicken eggs to a mobile phone over the entire period of incubation were published recently [73]. In this study, a series of 4 incubations were employed. During each incubation, 4 groups were used (control I, control II, experimental, sham). In the experimental group, the cell phone in call position was placed near (<25 cm) the eggs, whereas in the sham group the cell phone in off position was placed 1.5 m away from the exposed group. A significantly higher percentage of embryo mortality was observed in the experimental compared to the sham group in 2 of the 4 incubations. The lethal effects of embryo development in the experimental group were mainly observed between the 9th and 12th day of incubation.

Another issue that in recent years has attracted the attention of scientists is the effects of radiation from RF antennas on the biology of wild birds.

Balmori [74] investigated the possible effects of EMF from phone masts on a population of White stork (*Ciconia ciconia*). The total productivity in the nests located within 200 m of antennas was 0.86 ± 0.16 versus 1.6 ± 0.14 for those located further than 300 m. Another interesting observation, was that, 40% of the nests within 200 m of the antennae never had any chicks, while only 3.3% located further than 300 m never had chicks.

The influence of a military radar station [75] emitting pulsed modulated microwave radiation of 1200–3000 MHz was examined in tits (*Parus* sp). Experimental nest-boxes

were either exposed to a mean level of 3.41 ± 1.38 or $1.12 \pm 0.84 \text{ W/m}^2$. For control nest-boxes the exposure ranged from 0.001 to 0.01 W/m^2 . No statistically significant differences in the number of eggs or in the number of nestlings were observed between the two series (exposed, control) of tits.

6. Conclusions

The EMF were, are and will be a part of our life. The progress of science will provide the world with new EMF emitting technologies and subsequently with new problems. The monitoring of literature on this scientific field shows a shift of research which follows exactly the new technologies. The era of mobile telephony is beginning.

The evaluation of the possible effects of EMF on the living organism is a complex process that needs the combined contributions of many scientific disciplines. Due to the need for expertise in many different sciences, together with the technical problems of radiation studies, many times the published results are considered deficient in certain aspects. This is inevitable, and not an indication of poor quality. The inability to observe a biological effect in a particular study does not necessarily mean that such effect or/and adverse health effect is not present.

The vertebrate animal studies summarized in the present paper do not suggest strong effects of LF EMF on the male genital system. However, some studies on the development of animals, showed sensitivity, mainly observed in chickens. There is no convincing evidence from studies of mammals (Table 3), that exposure to EMF at levels associated with mobile telecommunications could be harmful for embryonic or postnatal development or for male fertility. On the other hand, the birds appeared to be more sensitive. The effects of EMF on the female genital system need further attention, since two experimental studies cannot lead to definitive conclusions.

The positive findings of the experimental studies with vertebrate animals are mainly attributed to the thermal effects of EMF. No valid evidence was found for the occurrence of non-thermal effects. However the non-thermal mechanisms must be the next target of the research.

The majority of reviewed studies were conducted in laboratories. This fact cannot represent the realistic situation of cell phone communication. On the other hand, the *in vivo* and simultaneously *in situ* studies are very scarce. Only Magras and Xenos conducted an *in situ* experiment which took place near an antenna park. That is because this kind of experiment is very difficult to carry out, and interaction with other exogenous factors could change the results.

One particular deficiency in most studies is that they describe experiments with acute or short-term exposure of animals on EMF. Experiments are needed to perform long-term exposure in order to demonstrate the chronic impact of EMF.

Another point that must be elucidated is that the majority of experimental animals used were small rodents (mice and rats), as well as chicken embryos. Further research is needed with the use of bigger animals such as dog and sheep.

The radiations emitted from masts that are situated in many rural and sylvatic areas could be possibly pathogenic in the wild animals. The wild animal populations could be candidate "experimental material" for closer observation of the possible effects of EMF on vertebrate models.

An important and intriguing aspect of the research is the possible role of the combination of RF with other pollutants such as chemical substances and other forms of radiation, as well as the interaction with drugs.

The potential health effects of EMF should be continually reassessed as new research results become available. EMF exposure guidelines also need to be updated or reconsidered as new scientific information on radiation and health risks is produced. However, additional studies might increase our understanding of the sensitivity of organisms to EMF.

References

- [1] J. Juutilainen, Developmental effects of electromagnetic fields, *Bioelectromagnetics* 7 (2005) S107–S115.
- [2] P.A. Valberg, T.E. van Deventer, M.H. Repacholi, Workgroup report: base stations and wireless networks-radiofrequency (RF) exposures and health consequences, *Environ. Health Perspect.* 115 (2007) 416–424.
- [3] M.H. Repacholi, Health risks from the use of mobile phones, *Toxicol. Lett.* 120 (2001) 323–331.
- [4] L.J. Challis, Mechanisms for interaction between RF fields and biological tissue, *Bioelectromagnetics* 7 (2005) S98–S106.
- [5] B.C.A. Windle, On certain early malformations of the embryo, *J. Anat. Physiol.* 27 (1893) 436–453.
- [6] B.C.A. Windle, On the effects of electricity and magnetism on development, *J. Anat. Physiol.* 29 (1895) 346–351.
- [7] H.J. Cook, N.H. Stenock, A.J. Vander, G.L. Kane, Early research on the biological effects of microwave radiation: 1940–1960, *Ann. Sci.* 37 (1980) 323–351.
- [8] W.M. Leach, Genetic, growth and reproductive effects of microwave radiation, *Bull. N.Y. Acad. Med.* 56 (1980) 249–257.
- [9] B. Algers, K. Hennrichs, Biological effects of electromagnetic fields. A review, *Vet. Res. Commun.* 6 (1983) 265–279.
- [10] T.E. Bryan, R.P. Gilderleeve, Effects of nonionizing radiation on birds, *Comp. Biochem. Physiol.* 89A (1988) 511–530.
- [11] E. Berman, L. Chacon, D. House, Development of chicken embryos in a pulsed magnetic field, *Bioelectromagnetics* 11 (1990) 169–187.
- [12] J. Juutilainen, Effects of low-frequency magnetic fields on embryonic development and pregnancy, *Scand. J. Work Environ. Health.* 17 (1991) 149–158.
- [13] N. Chernoff, J.M. Rogers, R. Kavet, A review of the literature on potential reproductive and developmental toxicity of electric and magnetic fields, *Toxicology* 74 (1992) 91–126.
- [14] R.L. Brent, W.E. Gordon, W.R. Bennett, D.A. Beckman, Reproductive and teratologic effects of electromagnetic fields, *Reprod. Toxicol.* 7 (1993) 535–580.
- [15] R.P. Jensch, Behavioral teratological studies using microwave radiation: is there an increased risk from exposure to cellular phones and microwave ovens? *Reprod. Toxicol.* 11 (1997) 601–611.

- [16] L. Verschaeve, A. Maes, Genetic, carcinogenic and teratogenic effects of radiofrequency fields, *Mutation Res.* 410 (1998) 141–165.
- [17] H. Huuskonen, M.-L. Lindbohm, J. Juutilainen, Teratogenic and reproductive effects of low-frequency magnetic fields, *Mutation Res.* 410 (1998) 167–183.
- [18] R.L. Brent, Reproductive and teratologic effects of low-frequency electromagnetic fields: a review of *in vivo* and *in vitro* studies using animal models, *Teratology* 59 (1999) 261–286.
- [19] M.E. O'Connor, Mammalian teratogenesis and radiofrequency fields, *Proc. IEEE* 68 (1980) 56–60.
- [20] L.N. Heynick, J.H. Merritt, Radiofrequency fields and teratogenesis, *Bioelectromagnetics* 6 (2003) (2003) S174–S186.
- [21] J. Juutilainen, Developmental effects of extremely low frequency electric and magnetic fields, *Radiat. Prot. Dosim.* 106 (2003) 385–390.
- [22] W. Löscher, Die Auswirkungen elektromagnetischer Felder von Mobilfunkendstationen auf Leistung, Gesundheit und Verhalten landwirtschaftlicher Nutztiere: Eine Bestandsaufnahme, *Der praktische Tierarzt* 84 (2003) 850–863.
- [23] R.D. Saunders, C.D. McCaig, Developmental effects of physiologically weak electric fields and heat: an overview, *Bioelectromagnetics* 7 (2005) S127–S132.
- [24] K.J. Fernic, S.J. Reynolds, The effects of electromagnetic fields from power lines on avian reproductive biology and physiology: a review, *J. Toxic Environ. Health B* 8 (2005) 127–140.
- [25] D. Krewski, B.W. Gluckman, R.W.Y. Habash, B. Habbick, W.G. Lotz, R. Mandeville, F.S. Prato, T. Salem, D.F. Weaver, Recent advances in research on radiofrequency fields and health: 2001–2003, *J. Toxic Environ. Health B* 10 (2007) 287–318.
- [26] T. Shigemitsu, K. Yamazaki, S. Nakasono, M. Kakkikawa, A review of studies of the biological effects of electromagnetic fields in the intermediate frequency range, *IEEE Trans. Electr. Electron. Eng.* 2 (2007) 405–412.
- [27] A. Elbeticha, M.A. Al-Akhras, H. Darmani, Long-term exposure of male and female mice to 50 Hz magnetic field: effects on fertility, *Bioelectromagnetics* 23 (2002) 168–172.
- [28] J.A. Heredia-Rojas, D.E. Caballero-Hernandez, A.O. Rodriguez-De La Fuente, G. Ramos-Alfaro, L.E. Rodriguez-Flores, Lack of alterations on meiotic chromosomes and morphological characteristics of male germ cells in mice exposed to a 60 Hz and 2.0 mT magnetic field, *Bioelectromagnetics* 25 (2004) 63–68.
- [29] J.S. Lee, S.S. Ahn, K.C. Jung, Y.W. Kim, S.K. Lee, Effects of 60 Hz electromagnetic field exposure on testicular germ cell apoptosis in mice, *Asian J. Androl.* 6 (2004) 29–34.
- [30] M.K. Chung, S.J. Lee, Y.B. Kim, S.C. Park, D.H. Shin, S.H. Kim, J.C. Kim, Evaluation of spermatogenesis and fertility in F1 male rats after *in utero* and neonatal exposure to extremely low frequency electromagnetic fields, *Asian J. Androl.* 7 (2005) 189–194.
- [31] M.A. Al-Akhras, H. Darmani, A. Elbeticha, Influence of 50 Hz magnetic field on sex hormones and other fertility parameters of adult male rats, *Bioelectromagnetics* 27 (2006) 127–131.
- [32] M.Z. Akdag, S. Dasdag, F. Aksen, B. Isik, F. Yilmaz, Effect of ELF magnetic fields on lipid peroxidation, sperm count, p53 and trace elements, *Med. Sci. Monit.* 12 (2006) 366–371.
- [33] F. Aksen, M.Z. Akdag, A. Ketani, B. Yokus, A. Kaya, S. Dasdag, Effect of 50 Hz 1 mT magnetic field on the uterus and ovaries of rats, *Med. Sci. Monit.* 12 (2006) 215–220.
- [34] S.H. Kim, H.J. Lee, S.Y. Choi, Y.M. Gimm, J.K. Pack, H.D. Choi, Y.S. Lee, Toxicity bioassay in Sprague–Dawley rats exposed to 20 kHz triangular magnetic field for 90 days, *Bioelectromagnetics* 27 (2006) 105–111.
- [35] A.A. Khaki, R.S. Tubbs, M.M. Shoja, J.S. Rad, A. Khaki, R.M. Farahani, S. Zarrintan, T.C. Nag, The effects of an electromagnetic field on the boundary tissue of the seminiferous tubules of the rat: a light and transmission electron microscope study, *Folia Morphol.* 65 (2006) 188–194.
- [36] H.J. Lee, S.H. Kim, S.Y. Choi, Y.M. Gimm, J.K. Pack, H.D. Choi, Y.S. Lee, Long-term exposure of Sprague–Dawley rats to 20 kHz triangular magnetic fields, *Int. J. Radiat. Biol.* 82 (2006) 285–291.
- [37] T.I. Subbotina, O.V. Tereshkina, A.A. Khadartsev, A.A. Yashin, Effect of low-intensity extremely high frequency radiation on reproductive function in Wistar rats, *Bull. Exp. Biol. Med.* 142 (2006) 189–190.
- [38] S. Erpek, M.D. Bilgin, E. Dikicioglu, A. Karul, The effects of low frequency electric field in rat testis, *Revue Méd. Vét* 158 (2007) 206–212.
- [39] S. Dasdag, M.A. Ketani, Z. Akdag, A.R. Ersay, I. Sari, O.C. Demirtas, M.S. Celik, Whole body microwave exposure emitted by cellular phones and testicular function of rats, *Urol. Res.* 27 (1999) 219–223.
- [40] S. Dasdag, M.Z. Akdag, F. Aksen, F. Yilmaz, M. Bashan, M.M. Dasdag, M.S. Celik, Whole body exposure of rats to microwaves emitted from a cell phone does not affect the testes, *Bioelectromagnetics* 24 (2003) 182–188.
- [41] R.J. Aitken, L.E. Bennetts, D. Sawyer, A.M. Wikleand, B.V. King, Impact of radio frequency electromagnetic radiation on DNA integrity in the male germ line *Inter. J. Androl.* 28 (2005) 171–179.
- [42] M. Ozguner, A. Koyu, G. Cesur, M. Ural, F. Ozguner, A. Gokcimen, N. Delibas, Biological and morphological effects on the reproductive organ of rats after exposure to electromagnetic field, *Saudi Med. J.* 26 (2005) 405–410.
- [43] Z. Forgács, Z. Somosy, G. Kubinyi, J. Bakos, A. Hudák, A. Surján, G. Thuróczy, Effects of whole body 1800 MHz GSM-like microwave exposure on testicular steroidogenesis and histology in mice, *Reprod. Toxicol.* 22 (2006) 111–117.
- [44] E.P. Ribeiro, E.L. Rhoden, M.M. Horn, C. Rhoden, L.P. Lima, L. Toniolo, Effects of sub chronic exposure to radio frequency from a conventional cellular telephone on testicular function in adult rats, *J. Urol.* 177 (2007) 395–399.
- [45] J.G. Yan, M. Agresti, T. Bruce, Y.H. Yan, A. Granlund, H.S. Matloub, Effects of cellular phone emissions on sperm motility in rats, *Fertil. Steril.* 88 (2007) 957–964.
- [46] F. Yilmaz, S. Dasdag, M.Z. Akdag, N. Kilinc, Whole body exposure of radiation emitted from 900 MHz mobile phones does not seem to affect the levels of anti-apoptotic bcl-2 protein, *Electromagn. Biol. Med.* 27 (2008) 65–72.
- [47] S. Dasdag, M.Z. Akdag, E. Ulukaya, A.K. Uzumlar, D. Yegin, Mobile phone exposure does not induce apoptosis on spermatogenesis in rats, *Arch. Med. Res.* 39 (2008) 40–44.
- [48] B. Oral, M. Guney, F. Ozguner, N. Karahan, T. Mungan, S. Comlekci, G. Cesur, Endometrial apoptosis induced by a 900 MHz mobile phone: preventive effects of vitamins E and C, *Adv. Ther.* 23 (2006) 957–973.
- [49] M. Guney, F. Ozguner, B. Oral, N. Karahan, T. Mungan, 900 MHz radiofrequency induced histopathologic changes and oxidative stress in rat endometrium: protection by vitamins E and C, *Toxicol. Ind. Health* 23 (2007) 411–420.
- [50] T. Negishi, S. Imai, M. Itabashi, I. Nishimura, T. Sasano, Studies of 50 Hz circularly polarized magnetic fields of up to 350 mT on reproduction and embryo-fetal development in rats: exposure during organogenesis or during preimplantation, *Bioelectromagnetics* 23 (2002) 369–389.
- [51] Y. Ohnishi, F. Mizuno, T. Sato, M. Yasui, T. Kikuchi, M. Ogawa, Effects of power frequency alternating magnetic fields on reproduction and prenatal development of mice, *J. Toxicol. Sci.* 27 (2002) 131–138.
- [52] Z.E. Gagnon, C. Newkirk, J.A. Conetta, M.A. Sama, S. Sisselman, Teratogenic effect of broad-band electromagnetic field on neonatal mice, *J. Environ. Sci. Health A Tox. Hazard Subst. Environ. Eng.* 38 (2003) 2465–2481.
- [53] M.K. Chung, J.C. Kim, S.H. Myung, D.I. Lee, Developmental toxicity evaluation of ELF magnetic fields in Sprague–Dawley rats, *Bioelectromagnetics* 24 (2003) 231–240.
- [54] M.S. Lahijani, K. Sajadi, Development of preincubated chicken eggs following exposure to 50 Hz electromagnetic fields with 1.33–7.32 mT flux densities, *Indian J. Exp. Biol.* 42 (2004) 858–865.
- [55] S.H. Kim, J.E. Song, S.R. Kim, H. Oh, Y.M. Gimm, D.S. Yoo, J.K. Pack, Y.S. Lee, Teratological studies of prenatal exposure of mice to a

- 20 kHz sawtooth magnetic field, *Bioelectromagnetics* 25 (2004) 114–117.
- [56] O. Roda-Murillo, J.A. Roda-Moreno, M.T. Morente-Chiquero, Effects of low-frequency magnetic fields on different parameters of embryo of *Gallus domesticus*, *Electromagn. Biol. Med.* 24 (2005) 55–62.
- [57] K. Saito, H. Suzuki, K. Suzuki, Teratogenic effects of static magnetic field on mouse fetuses, *Reprod. Toxicol.* 22 (2006) 118–124.
- [58] Y.N. Cao, Y. Zhang, Y. Liu, Effects of exposure to extremely low frequency electromagnetic fields on reproduction of female mice and development of offspring, *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 24 (2006) 468–470.
- [59] M.S. Lahijani, S.E. Nojooshi, S.F. Siadat, Light and electron microscope studies of effects of 50 Hz electromagnetic fields on preincubated chick embryo, *Electromagn. Biol. Med.* 26 (2007) 83–98.
- [60] W.F. Krueger, A.J. Giarola, J.W. Bradley, A. Shrekenhamer, Effects of electromagnetic fields on fecundity in the chicken, *Ann. N.Y. Acad. Sci.* 247 (1975) 391–400.
- [61] R.P. Jensh, I. Weinberg, R.I. Brent, Teratologic studies of prenatal exposure of rats to 915 MHz microwave radiation, *Radiat. Res.* 92 (1982) 160–171.
- [62] R.P. Jensh, W.H. Vogel, R.I. Brent, Postnatal functional analysis of prenatal exposure of rats to 915 MHz microwave radiation, *Int. J. Toxicol.* 1 (1982) 73–90.
- [63] E. Berman, C. Weil, P.A. Phillips, H.B. Carter, D.E. House, Fetal and maternal effects of continual exposure of rats to 970 MHz circularly-polarized microwaves, *Electromagn. Biol. Med.* 11 (1992) 43–54.
- [64] S. Klug, M. Hetscher, S. Giles, S. Kohlsman, K. Kramer, The lack of effects of nonthermal RF electromagnetic fields on the development of rat embryos grown in culture, *Life Sci.* 61 (1997) 1789–1802.
- [65] I.N. Magras, T.D. Xenos, RF radiation-induced changes in the prenatal development of mice, *Bioelectromagnetics* 18 (1997) 455–461.
- [66] M. Bornhausen, H. Scheingraber, Prenatal exposure to 900 MHz cell phone electromagnetic fields had no effect on operant-behaviour performances of adult rats, *Bioelectromagnetics* 21 (2000) 566–574.
- [67] S. Dasdag, M.Z. Akdag, O. Ayyıldız, O.C. Demirtas, M. Yayla, C. Sert, Do cellular phones alter blood parameters and birth weight of rats? *Electromagn. Biol. Med.* 19 (2000) 107–113.
- [68] B.L. Cobb, J.R. Jauchem, P.A. Mason, M.P. Dooley, S.A. Miller, J.M. Zirlax, M.R. Murphy, Neural and behavioural teratological evaluation of rats exposed to Ultra-Wideband electromagnetic fields, *Bioelectromagnetics* 21 (2000) 524–537.
- [69] M. Bastide, B.J. Youbicier-Simoa, J.C. Lebecqz, J. Gianninis, Toxicologic study of electromagnetic radiation emitted by television and video display screens and cellular telephones on chickens and mice, *Indoor Built Environ.* 10 (2001) 291–298.
- [70] Y. Grigoryev, Biological effects of mobile phone electromagnetic field on chick embryo (Risk assessment using the mortality rate), *Radiat. Biol. Radioecol.* 43 (2003) 541–543.
- [71] I.V. Ingole, S.K. Ghosh, Cell phone radiation and developing tissues in chicken embryo—a light microscopic study of kidneys, *J. Anat. Soc. India* 55 (2006) 19–23.
- [72] T. Kurnlin, H. Iivonen, P. Miettinen, A. Junonen, T. van Groen, L. Puranen, R. Pitkääho, R. Juutilainen, H. Tanila, Mobile phone radiation and the developing brain: behavioral and morphological effects in juvenile rats, *Radiat. Res.* 168 (2007) 471–479.
- [73] F. Batellier, I. Coury, D. Picard, J.P. Brillard, Effects of exposing chicken eggs to a cell phone in 'call' position over the entire incubation period, *Theriogenology* 69 (2008) 737–745.
- [74] A. Balmori, Possible effects of electromagnetic fields from phone masts on a population of White stork (*Ciconia ciconia*), *Electromagn. Biol. Med.* 24 (2005) 109–119.
- [75] L. Rejt, T. Mazgajski, R. Kubacki, J. Kieliszek, E. Sobiczewska, S. Szmigielski, Influence of radar radiation on breeding biology of tuis (*Parus* sp.), *Electromagn. Biol. Med.* 26 (2007) 235–238.

Electromagnetic pollution from phone masts. Effects on wildlife

Alfonso Balmori

*Dirección General del Medio Natural, Consejería de Medio Ambiente, Junta de Castilla y León, C/Rigoberto Cortezoso,
14, 47014 Valladolid, Spain*

Received 10 August 2008; received in revised form 28 August 2008; accepted 30 January 2009

Abstract

A review on the impact of radiofrequency radiation from wireless telecommunications on wildlife is presented. Electromagnetic radiation is a form of environmental pollution which may hurt wildlife. Phone masts located in their living areas are irradiating continuously some species that could suffer long-term effects, like reduction of their natural defenses, deterioration of their health, problems in reproduction and reduction of their useful territory through habitat deterioration. Electromagnetic radiation can exert an aversive behavioral response in rats, bats and birds such as sparrows. Therefore microwave and radiofrequency pollution constitutes a potential cause for the decline of animal populations and deterioration of health of plants living near phone masts. To measure these effects urgent specific studies are necessary. © 2009 Elsevier Ireland Ltd. All rights reserved.

Keywords: Effects on wildlife; Effects on birds; Electromagnetic radiation; Mammals; Microwaves; Mobile telecommunications; Non-thermal effects; Phone masts; Radiofrequencies

1. Introduction

Life has evolved under the influence of two omnipresent forces: gravity and electromagnetism. It should be expected that both play important roles in the functional activities of organisms [1]. Before the 1990's radiofrequencies were mainly from a few radio and television transmitters, located in remote areas and/or very high places. Since the introduction of wireless telecommunication in the 1990's the rollout of phone networks has caused a massive increase in electromagnetic pollution in cities and the countryside [2,3].

Multiple sources of mobile communication result in chronic exposure of a significant part of the wildlife (and man) to microwaves at non-thermal levels [4]. In recent years, wildlife has been chronically exposed to microwaves and RFR (Radiofrequency radiation) signals from various sources, including GSM and UMTS/3G wireless phones and base stations, WLAN (Wireless Local Area Networks), WPAN (Wireless Personal Area Networks such as Bluetooth), and DECT (Digital Enhanced (former European) Cordless Telecommunications) that are erected indiscriminately without studies of environmental impact measuring

long-term effects. These exposures are characterized by low intensities, varieties of signals, and long-term durations. The greater portion of this exposure is from mobile telecommunications (geometric mean in Vienna: 73% [5]). In Germany the GSM cellular phone tower radiation is the dominating high frequency source in residential areas [6]. Also GSM is the dominating high frequency source in the wilderness of Spain (personal observation).

Numerous experimental data have provided strong evidence of athermal microwave effects and have also indicated several regularities in these effects: dependence of frequency within specific frequency windows of "resonance-type"; dependence on modulation and polarization; dependence on intensity within specific intensity windows, including super-low power density comparable with intensities from base stations/masts [4,7–9]. Some studies have demonstrated different microwave effects depending on wavelength in the range of mm, cm or m [10,11]. Duration of exposure may be as important as power density. Biological effects resulting from electromagnetic field radiation might depend on dose, which indicates long-term accumulative effects [3,9,12]. Modulated and pulsed radiofrequencies seem to be more effective in producing effects [4,9]. Pulsed waves (in blasts), as well as certain low frequency modulations exert greater

E-mail addresses: abalmori@ono.com, balmoral@jcy1.es.

biological activity [11,13–15]. This observation is important because cell phone radiation is pulsed microwave radiation modulated at low frequencies [8,9].

Most of the attention on possible biological effects of electromagnetic radiation from phone masts has been focused on human health [5,16–21]. The effects of electromagnetic pollution on wildlife, have scarcely been studied [22–25].

The objective of this review is to detail advances in knowledge of radiofrequencies and microwave effects on wildlife. Future research may help provide a better understanding of electromagnetic field (EMF) effects on wildlife and plants and their conservation.

2. Effects on exposed wildlife

2.1. Effects on birds

2.1.1. Effects of phone mast microwaves on white stork

In monitoring a white stork (*Ciconia ciconia*) population in Valladolid (Spain) in vicinity of Cellular Phone Base Stations, the total productivity in nests located within 200 m of antennae, was 0.86 ± 0.16 . For those located further than 300 m, the result was practically doubled, with an average of 1.6 ± 0.14 . Very significant differences among total productivity were found ($U=240$; $P=0.001$, Mann–Whitney test). Twelve nests (40%) located within 200 m of antennae never had chicks, while only one (3.3%) located further than 300 m had no chicks. The electric field intensity was higher on nests within 200 m (2.36 ± 0.82 V/m) than nests further than 300 m (0.53 ± 0.82 V/m). In nesting sites located within 100 m of one or several cellsite antennae with the main beam of radiation impacting directly (Electric field intensity >2 V/m) many young died from unknown causes. Couples frequently fought over nest construction sticks and failed to advance the construction of the nests. Some nests were never completed and the storks remained passively in front of cellsite antennae. These results indicate the possibility that microwaves are interfering with the reproduction of white stork [23]. (Fig. 1)




Fig. 1. Average number of young and electric field intensity (V/m) in 60 nests of white storks (*Ciconia ciconia*) (Hallberg, Ö with data of Balmori, 2005 [23]).

2.1.2. Effects of phone mast microwaves on house sparrows

A possible effect of long-term exposure to low-intensity electromagnetic radiation from mobile phone (GSM) base stations on the number of house sparrows during the breeding season was studied in Belgium. The study was carried out sampling 150 point locations within six areas to examine small-scale geographic variation in the number of house sparrow males and the strength of electromagnetic radiation from base stations. Spatial variation in the number of house sparrow males was negative and highly significantly related to the strength of electric fields from both the 900 and 1800 MHz downlink frequency bands and from the sum of these bands (Chi-square-tests and AIC-criteria, $P < 0.001$). This negative relationship was highly similar within each of the six study areas, despite differences among areas in both the number of birds and radiation levels. Fewer house sparrow males were seen at locations with relatively high electric field strength values of GSM base stations and therefore support the notion that long-term exposure to higher levels of radiation negatively affects the abundance or behavior of house sparrows in the wild [24].

In another study with point transect sampling performed at 30 points visited 40 times in Valladolid (Spain) between 2002 and 2006, counting the sparrows and measuring the mean electric field strength (radiofrequencies and microwaves: 1 MHz to 3 GHz range). Significant declines ($P=0.0037$) were observed in mean bird density over time, and significantly low bird density was observed in areas with high electric field strength. The logarithmic regression of the mean bird density vs. field strength groups (considering field strength in 0.1 V/m increments) was $R = -0.87$; $P = 0.0001$. According to this calculation, no sparrows would be expected to be found in an area with field strength >4 V/m [25]. (Fig. 2)

In the United Kingdom a decline of several species of urban birds, especially sparrows, has recently happened [26]. The sparrow population in England has decreased in the last 30 years from 24 million to less than 14. The more abrupt decline, with 75% descent has taken place from 1994 to 2002. In 2002, the house sparrow was added to the Red List of U.K. endangered species [27]. This coincides with the rollout of mobile telephony and the




Fig. 2. Mean sparrow density as a function of electric field strength grouped in 0.1 V/m. (Balmori and Hallberg, 2007 [25]).




Fig. 3. Annual number of contacts (Mean) for 14 species studied in "Campo Grande" urban park (lack the information of the years 1999–2001).

possible relationship of both circumstances should be investigated.

In Brussels, many sparrows have disappeared recently [28]; similar declines have been reported in Dublin [29]. Van der Poel (cited in Ref. [27]) suggested that sparrows might be declining in Dutch urban centres also.

2.1.3. Effects on the bird community at an urban park

Microwaves may be affecting bird populations in places with high electromagnetic pollution. Since several antennas were installed in proximities of "Campo Grande" urban park (Valladolid, Spain) the bird population has decreased and a reduction of the species and breeding couples has occurred. Between 1997 and 2007, of 14 species, 3 species have disappeared, 4 are in decline and 7 stay stable (Balmori, unpublished data) (Fig. 3). In this time the air pollution (SO₂, NO₂, CO and Benzene) has diminished.

During the research some areas called "silence areas" contaminated with high microwave radiation (>2 V/m), where previously different couples usually bred and later disappeared, have been found. Several anomalies in magpies (*Pica pica*) were detected: plumage deterioration, locomotive problems (limps and deformations in the paws), partial albinism and melanism, especially in flanks [30]. Recently cities have increased cases of partial albinism and melanism in birds (*Passer domesticus*, *Turdus merula* and *P. pica*) (personal observation).

2.1.4. Possible physiological mechanisms of the effects found in birds

Current scientific evidence indicates that prolonged exposure to EMFs, at levels that can be encountered in the environment, may affect immune system function by affecting biological processes [3,31,32]. A stressed immune system may increase the susceptibility of a bird to infectious diseases, bacteria, viruses, and parasites [33].

The plumage of the birds exposed to microwaves looked, in general, discolored and lack of shine. This not only occurred in ornamental birds; such as peacocks, but also in wild birds; such as, tits, great tits, house sparrows, etc (personal observation). We must mention that plumage deterioration is the first sign of weakening or illnesses in birds since damaged feathers are a sure sign of stress.

Physiological conditions during exposure minimize microwave effects. Radical scavengers/antioxidants might be involved in effects of microwaves [4].

Microwaves used in cellphones produce an athermal response in several types of neurons of the birds nervous system [34]. Several studies addressed behavior and teratology in young birds exposed to electromagnetic fields [23,25,35–37]. Most studies indicate that electromagnetic field exposure of birds generally changes, but not always consistently in effect or in direction, their behavior, reproductive success, growth and development, physiology and endocrinology, and oxidative stress [37]. These results can be explained by electromagnetic fields affecting the birds' response to the photoperiod as indicated by altered melatonin levels [38].

Prolonged mobile phone exposure may have negative effects on sperm motility characteristics and male fertility as has been demonstrated in many studies made in man and rats [39–46]. EMF and microwaves can affect reproductive success in birds [23,25,35,36,47]. EMF exposure affected reproductive success of kestrels (*Falco sparverius*), increasing fertility, egg size, embryonic development and fledging success but reducing hatching success [35,36].

The radiofrequency and microwaves from mobile telephony can cause genotoxic effects [48–55]. Increases in cytological abnormalities imply long-term detrimental effects since chromosomal damage is a mechanism relevant to causation of birth defects and cancer [55].

Long-term continuous, or daily repeated EMF exposure can induce cellular stress responses at non-thermal power levels that lead to an accumulation of DNA errors and to inhibition of cell apoptosis and cause increased permeability of blood–brain barrier due to stabilization of endothelial cell stress fibers. Repeated occurrence of these events over a long period of time (years) could become a health hazard due to a possible accumulation of brain tissue damage. These findings have important implications with regards to potential dangers from prolonged and repeated exposure to non-ionizing radiation [56,57].

Pulsed magnetic fields can have a significant influence on the development and incidence of abnormalities in chicken embryos. In five of six laboratories, exposed embryos exhibited more structural anomalies than controls. If the data from all six laboratories are pooled, the difference for the incidence of abnormalities in exposed embryos and controls is highly significant [58]. Malformations in the nervous system and heart, and delayed embryo growth are observed. The embryo is most sensitive to exposure in the first 24 h of incubation [58]. An increase in the mortality [59] and appearance of morphological abnormalities, especially of the neural tube [13,60,61] has been recorded in chicken embryos exposed to pulsed magnetic fields, with different susceptibility among individuals probably for genetic reasons. A statistically significant high mortality rate of chicken embryos subjected to radiation from a cellphone, compared to the control group exists [62,63]. In another study eggs exposed to a magnetic

field intensity of 0.07 T showed embryonic mortality during their incubation was higher. The negative effect of the magnetic field was manifested also by a lower weight of the hatched chicken [64]. Bioelectric fields have long been suspected to play a causal role in embryonic development. Alteration of the electrical field may disrupt the chemical gradient and signals received by embryo cells. It appears that in some manner, cells sense their position in an electrical field and respond appropriately. The disruption of this field alters their response. Endogenous current patterns are often correlated with specific morphogenetic events [65].

Available data suggests dependencies of genotype, gender, physiological and individual factors on athermal microwave effects [4,9]. Genomic differences can influence cellular responses to GSM Microwaves. Data analysis has highlighted a wide inter-individual variability in response, which was replicated in further experiments [4]. It is possible that each species and each individual, show different susceptibility to radiation, since vulnerability depends on genetic tendency, and physiologic and neurological state of the irradiated organism [15,35–37,61,66–68]. Different susceptibility of each species has also been proven in wild birds exposed to electromagnetic fields from high-voltage power lines [47].

2.2. Effects on mammals

2.2.1. Alarm and aversion behavior

Rats spent more time in the halves of shuttle boxes that were shielded from 1.2 GHz. Microwaves irradiation. The average power density was about 0.6 mW/cm². Data revealed that rats avoided the pulsed energy, but not the continuous energy, and less than 0.4 mW/cm² average power density was needed to produce aversion [69]. Navakatikian & Tomashevskaya [70] described a complex series of experiments in which they observed disruption of rat behavior (active avoidance) from radiofrequency radiation. Behavioral disruption was observed at a power density as low as 0.1 mW/cm² (0.027 W/kg). Mice in an experimental group exposed to microwave radiation expressed visible individual panic reaction, disorientation and a greater degree of anxiety. In the sham exposed group these deviations of behavior were not seen and all animals show collective defense reaction [71]. Microwave radiation at 1.5 GHz pulsing 16 ms. At 0.3 mW/cm² power density, in sessions of 30 min/day over one month produced anxiety and alarm in rabbits [72].

Electromagnetic radiation can exert an aversive behavioral response in bats. Bat activity is significantly reduced in habitats exposed to an electromagnetic field strength greater than 2 V/m [73]. During a study in a free-tailed bat colony (*Tadarida teniotis*) the number of bats decreased when several phone masts were placed 80 m from the colony [74].

2.2.2. Deterioration of health

Animals exposed to electromagnetic fields can suffer a deterioration of health and changes in behavior [75,76].

There was proof of frequent death in domestic animals; such as, hamsters and guinea pigs, living near mobile telecommunication base stations (personal observation).

The mice in an experimental group exposed to microwave radiation showed less weight gain compared to control, after two months. The amount of food used was similar in both groups [71]. A link between electromagnetic field exposure and higher levels of oxidative stress appears to be a major contributor to aging, neurodegenerative diseases, immune system disorders, and cancer in mammals [33].

The effects from GSM base transceiver station (BTS) frequency of 945 MHz on oxidative stress in rats were investigated. When EMF at a power density of 3.67 W/m², below current exposure limits, were applied, MDA (malondialdehyde) level was found to increase and GSH (reduced glutathione) concentration was found to decrease significantly ($P < 0.0001$). Additionally, there was a less significant ($P = 0.0190$) increase in SOD (superoxide dismutase) activity under EM exposure [77].

2.2.3. Problems in reproduction

In the town of Casavieja (Ávila, Spain) a telephony antenna was installed that had been in operation for about 5 years. Then some farmers began blaming the antenna for miscarriages in many pigs, 50–100 m from the antenna (on the outskirts of the town). Finally the topic became so bad that the town council decided to disassemble the antenna. It was removed in the spring 2005. From this moment onwards the problems stopped (C. Lumbreras personal communication).

A Greek study reports a progressive drop in the number of rodent births exposed to radiofrequencies. The mice exposed to 0.168 μ W/cm² become sterile after five generations, while those exposed to 1.053 μ W/cm² became sterile after only three generations [22].

In pregnant rats exposed to 27.12 MHz continuous waves at 100 μ W/cm² during different periods of pregnancy, half the pregnancies miscarried before the twentieth day of gestation, compared to only a 6% miscarriage rate in unexposed controls, and 38% of the viable foetuses had incomplete cranial ossification, compared to less than 6% of the controls. Findings included a considerable increase in the percentage of total reabsorptions (post-implantation losses consequent to RF radiation exposure in the first post-implantation stage). Reduced body weight in the exposed dams reflected a negative influence on their health. It seems that the irradiation time plays an important role in inducing specific effects consequent to radiofrequency radiation exposure [78]. There was also a change in the sex ratio, with more males born to rats that had been irradiated from the time of conception [2]. Moorhouse and Macdonald [79] find a substantial decline in female Water Vole numbers in the radio-collared population, apparently resulting from a male skew in the sex ratios of offspring born to this population. Recruits to the radio-tracked population were skewed heavily in favour of males (43:13). This suggests that radio-collaring of females caused male-skewed sex ratios.

Mobile phone exposure may have negative effects on sperm motility characteristics and male fertility in rats [46]. Other studies find a decrease of fertility, increase of deaths after birth and dystrophic changes in their reproductive organs [11]. Intermittent exposure showed a stronger effect than continuous exposure [4]. Brief, intermittent exposure to low-frequency EM fields during the critical prenatal period for neurobehavioral sex differentiation can demasculinize male scent marking behavior and increase accessory sex organ weights in adulthood [80].

In humans, magnetic field exposures above 2.0 mG were positively associated with miscarriage risk [81]. Exposure of pregnant women to mobile phone significantly increased foetal and neonatal heart rate, and significantly decreased the cardiac output [82].

2.2.4. Nervous system

Microwaves may affect the blood brain barrier which lets toxic substances pass through from the blood to the brain [83]. Adang et al. [84] examined the effect of microwave exposure to a GSM-like frequency of 970 MHz pulsed waves on the memory in rats by means of an object recognition task. The rats that have been exposed for 2 months show normal exploratory behavior. The animals that have been exposed for 15 months show derogatory behavior. They do not make the distinction between a familiar and an unfamiliar object. In the area that received radiation directly from "Location Skrunda Radio Station" (Latvia), exposed children had less developed memory and attention, their reaction time was slower and neuromuscular apparatus endurance was decreased [85]. Exposure to cell phones prenatally and, to a lesser degree, postnatally, was associated with behavioral difficulties such as emotional and hyperactivity problems around 7 years of age [86]. Electromagnetic radiation caused modification of sleep and alteration of cerebral electric response (EEG) [87–89]. Microwave radiation from phone masts may cause aggressiveness in people and animals (personal observation).

2.3. Effects on amphibians

Disappearance of amphibians and other organisms is part of the global biodiversity crisis. An associated phenomenon is the appearance of large numbers of deformed amphibians. The problem has become more prevalent, with deformity rates up to 25% in some populations, which is significantly higher than previous decades [90]. Balmori [91] proposed that electromagnetic pollution (in the microwave and radiofrequency range) is a possible cause for deformations and decline of some wild amphibian populations.

Two species of amphibians were exposed to magnetic fields at various stages of development. A brief treatment of early amphibian embryos produced several types of abnormalities [92]. Exposure to a pulsed electromagnetic field produced abnormal limb regeneration in adult Newts [93]. Frog tadpoles (*Rana temporaria*) developed under electro-

magnetic field (50 Hz, 260 A/m) have increased mortality. Exposed tadpoles developed more slowly and less synchronously than control tadpoles and remain at the early stages for longer. Tadpoles developed allergies and EMF caused changes in blood counts [94].

In a current study, exposing eggs and tadpoles ($n=70$) of common frog (*R. temporaria*) for two months, from the phase of eggs until an advanced phase of tadpole, to four telephone base stations located 140 m away: with GSM system 948.0–959.8 MHz; DCS system: 1830.2–1854.8; 1855.2–1879.8 MHz. and UMTS system: 1905–1910; 1950–1965; 2140–2155 MHz. (electric field intensity: 1.847–2.254 V/m). A low coordination of movements, an asynchronous growth, with big and small tadpoles, and a high mortality (90%) was observed. The control group ($n=70$), under the same conditions but inside a Faraday cage (metallic shielding component: EMC-reinforcement fabrics 97442 Marburg Technic), the coordination of movements was normal, the development was synchronously and the mortality rate was only 4.2% [95].

2.4. Effects on insects

The microwaves may affect the insects. Insects are the basis and key species of ecosystems and they are especially sensitive to electromagnetic radiation that poses a threat to nature [96].

Carpenter and Livstone [97] irradiated pupae of *Tenebrio molitor* with 10 GHz microwaves at 80 mW for 20–30 min and 20 mW for 120 min obtained a rise in the proportion of insects with abnormalities or dead. In another study exposing fruit flies (*Drosophila melanogaster*) to mobile phone radiation, elevated stress protein levels (Hsp70) was obtained, which usually means that cells are exposed to adverse environmental conditions ('non-thermal shock') [98]. Panagopoulos et al. [99] exposed fruit flies (*D. melanogaster*) to radiation from a mobile phone (900 MHz) during the 2–5 first days of adulthood. The reproductive capacity of the species reduced by 50–60% in modulated radiation conditions (emission while talking on the phone) and 15–20% with radiation nonmodulated (with the phone silent). The results of this study indicate that this radiation affects the gonadal development of insects in an athermal way. The authors concluded that radio frequencies, specifically GSM, are highly bioactive and provoke significant changes in physiological functions of living organisms. Panagopoulos et al. [100] compare the biological activity between the two systems GSM 900 MHz and DCS 1800 MHz in the reproductive capacity of fruit flies. Both types of radiation were found to decrease significantly and non-thermally the insect's reproductive capacity, but GSM 900 MHz seems to be even more bioactive than DCS 1800 MHz. The difference seems to be dependent mostly on field intensity and less on carrier frequency.

A study in South Africa finds a strong correlation between decrease in ant and beetle diversity with the

electromagnetic radiation exposure (D. MacFadyen, personal communication.). A decrease of insects and arachnids near base stations was detected and corroborated by engineers and antenna's maintenance staff [101]. In houses near antennas an absence of flies, even in summer, was found.

In a recent study carried out with bees in Germany, only a few bees irradiated with DECT radiation returned to the beehive and they needed more time. The honeycomb weight was lower in irradiated bees [102]. In recent years a "colony collapse disorder" is occurring that some authors relate with pesticides and with increasing electromagnetic pollution [96].

The disappearance of insects could have an influence on bird's weakening caused by a lack of food, especially at the first stages in a young bird's life.

2.5. Effects on trees and plants

The microwaves may affect vegetables. In the area that received radiation directly from "Location Skrunda Radio Station" (Latvia), pines (*Pinus sylvestris*) experienced a lower growth radio. This did not occur beyond the area of impact of electromagnetic waves. A statistically significant negative correlation between increase tree growth and intensity of electromagnetic field was found, and was confirmed that the beginning of this growth decline coincided in time with the start of radar emissions. Authors evaluated other possible environmental factors which might have intervened, but none had noticeable effects [103]. In another study investigating cell ultrastructure of pine needles irradiated by the same radar, there was an increase of resin production, and was interpreted as an effect of stress caused by radiation, which would explain the aging and declining growth and viability of trees subjected to pulsed microwaves. They also found a low germination of seeds of pine trees more exposed [104]. The effects of Latvian radar was also felt by aquatic plants. *Spirodela polyrrhiza* exposed to a power density between 0.1 and 1.8 $\mu\text{W}/\text{cm}^2$ had lower longevity, problems in reproduction and morphological and developmental abnormalities compared with a control group who grew up far from the radar [105].

Chlorophylls were quantitatively studied in leaves of black locust (*Robinia pseudoacacia* L.) seedlings exposed to high frequency electromagnetic fields of 400 MHz. It was revealed that the ratio of the two main types of chlorophyll was decreasing logarithmically to the increase of daily exposure time [106].

Exposed tomato plants (*Lycopersicon esculentum*) to low level (900 MHz, 5 V/m) electromagnetic fields for a short period (10 min) measured changes in abundance of three specific mRNA after exposure, strongly suggesting that they are the direct consequence of application of radio-frequency fields and their similarities to wound responses suggests that this radiation is perceived by plants as an injurious stimulus [107]. Non-thermal exposure to radiofrequency fields

induced oxidative stress in duckweed (*Lemna minor*) as well as unespecific stress responses, especially of antioxidative enzymes [108].

For some years progressive deterioration of trees near phone masts have been observed in Valladolid (Spain). Trees located inside the main lobe (beam), look sad and feeble, possibly slow growth and a high susceptibility to illnesses and plagues. In places we have measured higher electric field intensity levels of radiation (>2 V/m) the trees show a more notable deterioration [109]. The tops of trees are dried up where the main beams are directed to, and they seem to be most vulnerable if they have their roots close to water. The trees don't grow above the height of the other ones and, those that stand out far above, have dried tops (Hargreaves, personal communication and personal observation). White and black poplars (*Populus sp.*) and willows (*Salix sp.*) are more sensitive. There may be a special sensitivity of this family exists or it could be due to their ecological characteristics forcing them to live near water, and thus electric conductivity. Other species as *Platanus sp.* and *Lygustrum japonicum*, are more resistant (personal observation). Schorpp [110] presents abundant pictures and explanations of what happens to irradiated trees.

3. Conclusions

This literature review shows that pulsed telephony microwave radiation can produce effects especially on nervous, cardiovascular, immune and reproductive systems [111]:

- Damage to the nervous system by altering electroencephalogram, changes in neural response or changes of the blood-brain barrier.
- Disruption of circadian rhythms (sleep-wake) by interfering with the pineal gland and hormonal imbalances.
- Changes in heart rate and blood pressure.
- Impairment of health and immunity towards pathogens, weakness, exhaustion, deterioration of plumage and growth problems.
- Problems in building the nest or impaired fertility, number of eggs, embryonic development, hatching percentage and survival of chickens.
- Genetic and developmental problems: problems of locomotion, partial albinism and melanism or promotion of tumors.

In the light of current knowledge there is enough evidence of serious effects from this technology to wildlife. For this reason precautionary measures should be developed, alongside environmental impact assessments prior to installation, and a ban on installation of phone masts in protected natural areas and in places where endangered species are present. Surveys should take place to objectively assess the severity of effects.

Acknowledgment

The author is grateful to Denise Ward and Örjan Hallberg.

References

- [1] J.M.R. Delgado, Biological effects of extremely low frequency electromagnetic fields, *J. Bioelectr.* 4 (1985) 75–91.
- [2] A. Pirtenberg, *Microwaving Our Planet: The Environmental Impact of the Wireless Revolution*, 11210, Cellular Phone Taskforce, Brooklyn, NY, 1997.
- [3] A.L. Galeev, The effects of microwave radiation from mobile telephones on humans and animals, *Neurosci. Behav. Physiol.* 30 (2000) 187–194.
- [4] I. Belyaev, Non-thermal biological effects of microwaves, *Microw. Rev.* 11 (2005) 13–29, <http://www.mwr.mediansis.net/pdf/Vol11No2-03-IBelyaev.pdf>.
- [5] H.P. Hunter, H. Moshhammer, P. Wallner, M. Kundi, Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations, *Occup. Environ. Med.* 63 (2006) 307–313.
- [6] T. Haumann, U. Munzenberg, W. Maes, P. Sierck, HF-radiation levels of GSM cellular phone towers in residential areas, in: 2nd International Workshop on Biological effects of EMFS, Rhodes, Greece, 2002.
- [7] W.R. Adey, Tissue interactions with non-ionizing electromagnetic fields, *Physiol. Rev.* 61 (1981) 435–514.
- [8] G.J. Hyland, Physics and biology of mobile telephony, *Lancet* 356 (2000) 1–8.
- [9] H. Lai, Biological effects of radiofrequency electromagnetic field, in: *Encyclopaedia of Biomaterials and Biomedical Engineering*, 2005, doi:10.1081/E-EBBE-120041846, pp. 1–8.
- [10] S. Kemerov, M. Marinkev, D. Getova, Effects of low-intensity electromagnetic fields on behavioral activity of rats, *Folia Med.* 41 (1999) 75–80.
- [11] N. Nikolaevich, A. Igorevna, and G. Vasil, Influence of high-frequency electromagnetic radiation at non-thermal intensities on the human body (A review of work by Russian and Ukrainian researchers), No place to hide, 3 (Supplement), 2001.
- [12] W.R. Adey, Bioeffects of mobile communications fields: possible mechanisms for cumulative dose, in: N. Kuster, Q. Balzano, J.C. Lin, (Eds.), *Mobile communications safety*, New York: Chapman & Hall, 1997, pp. 95–131.
- [13] A. Úbeda, M.A. Trillo, L. Chacón, M.J. Blanco, J. Leal, Chick embryo development can be irreversibly altered by early exposure to weak extremely-low-frequency magnetic fields, *Bioelectromagnetics* 15 (1994) 385–398.
- [14] I.U.G. Grigoriev, Role of modulation in biological effects of electromagnetic radiation, *Radiats. Biol. Radiocool.* 36 (1996) 659–670.
- [15] G.J. Hyland, The physiological and environmental effects of non-ionising electromagnetic radiation, Working document for the STOA Panel, European Parliament, Directorate General for Research, 2001.
- [16] R. Santini, J.M. Santini, P. Danze, M. Leruz, M. Seigne, Enquête sur la santé de riverains de stations relais: I. Incidences de la distance et du sexe, *Pathol. Biol.* 50 (2002) 369–373.
- [17] R. Santini, P. Santini, J.M. Le Ruz, M. Danze, M. Seigne, Survey study of people living in the vicinity of cellular phone base stations, *Electromagn. Biol. Med.* 22 (2003) 41–49.
- [18] R. Santini, P. Santini, J.M. Danze, P. Le Ruz, M. Seigne, Symptoms experienced by people in vicinity of base stations: II/Incidences of age, duration of exposure, location of subjects in relation to the antennas and other electromagnetic factors, *Pathol. Biol.* 51 (2003) 412–415.
- [19] E.A. Navarro, J. Segura, M. Portolés, C. Gómez Perreita, The microwave syndrome: a preliminary study in Spain, *Electromagn. Biol. Med.* 22 (2003) 161–169.
- [20] G. Oberfeld, E. Navarro, M. Portoles, C. Maestu, C. Gomez-Perreita, The microwave syndrome—further aspects of a Spanish study, in: EBBA Congress Kos, Greece, 2004.
- [21] G. Abdel-Rassoul, M.A. Salem, A. Michael, F. Farahat, M. El-Batanouny, E. Salem, Neurobehavioral effects among inhabitants around mobile phone base stations, *Neurotoxicology* 28 (2007) 434–440.
- [22] I.N. Magras, T.D. Xenos, Radiation-induced changes in the prenatal development of mice, *Bioelectromagnetics* 18 (1997) 455–461.
- [23] A. Balmori, Possible effects of electromagnetic fields from phone masts on a population of white stork (*Ciconia ciconia*), *Electromagn. Biol. Med.* 24 (2005) 109–119.
- [24] J. Everaert, D. Bauwens, A possible effect of electromagnetic radiation from mobile phone base stations on the number of breeding House Sparrows (*Passer domesticus*), *Electromagn. Biol. Med.* 26 (2007) 63–72.
- [25] A. Balmori, Ö. Hallberg, The urban decline of the house sparrow (*Passer domesticus*): a possible link with electromagnetic radiation, *Electromagn. Biol. Med.* 26 (2007) 141–151.
- [26] M.J. Raven, D.G. Noble, S.R. Baillie, The breeding bird survey (2002), BTO Research Report 334, British Trust for Ornithology, Theford, 2003.
- [27] J.D. Summers-Smith, The decline of the house sparrow: a review, *Brit. Birds* 96 (2003) 439–446.
- [28] J. De Laet, Ligue Royale Belge pour la Protection des Oiseaux avec l'Université de Gand, 2004, <<http://www.protectiondesoiseaux.be/content/view/full/74>> (Accessed on May 20, 2008).
- [29] A. Prowse, The urban decline of the house sparrow, *Brit. Birds* 95 (2002) 143–146.
- [30] A. Balmori, Aves y telefonía móvil. Resultados preliminares de los efectos de las ondas electromagnéticas sobre la fauna urbana, *El ecologista* 36 (2003) 40–42.
- [31] C.K. Chou, A.W. Guy, L.L. Kunz, R.B. Johnson, J.J. Crowley, J.H. Krupp, Long-term, low-level microwave irradiation of rats, *Bioelectromagnetics* 13 (1992) 469–496.
- [32] E.T. Novoselova, E.E. Fesenko, Stimulation of production of tumour necrosis factor by murine macrophages when exposed in vivo and in vitro to weak electromagnetic waves in the centimeter range, *Biofizika* 43 (1998) 1132–1133.
- [33] K.J. Fernie, D.M. Bird, Evidence of oxidative stress in American kestrels exposed to electromagnetic fields, *Environ. Res.* 86 (2001) 198–207.
- [34] R.C. Beason-Held, P. Semm, Responses of neurons to an amplitude modulated microwave stimulus, *Neurosci. Lett.* 33 (2002) 175–178.
- [35] K.J. Fernie, D.M. Bird, R.D. Dawson, F.C. Lagac, Effects of electromagnetic fields on the reproductive success of American kestrels, *Physiol. Biochem. Zool.* 73 (2000) 60–65.
- [36] K.J. Fernie, N.J. Leonard, D.M. Bird, Behavior of free-ranging and captive American kestrels under electromagnetic fields, *J. Toxicol. Environ. Health, Part A* 59 (2000) 597–603.
- [37] K.J. Fernie, S.J. Reynolds, The effects of electromagnetic fields from power lines on avian reproductive biology and physiology: a review, *J. Toxicol. Environ. Health, Part B* 8 (2005) 127–140.
- [38] K.J. Fernie, D.M. Bird, Effects of electromagnetic fields on body mass and food-intake of American kestrels, *Condor* 101 (1999) 616–621.
- [39] S. Dasdag, M.A. Ketani, Z. Akdag, A.R. Erşay, İ. Sar, Ö.C. Demirtaş, M.S. Celik, Whole body microwave exposure emitted by cellular phones and testicular function of rats, *Urol. Res.* 27 (1999) 219–223.
- [40] M. Davoudi, C. Brösner, W. Kuber, Der Einfluss elektromagnetischer wellen auf die Spermienmobilität, *J. für Urol. Urogynäkol.* 9 (2002) 18–22.
- [41] I. Fejes, Z. Za Vaczki, J. Szollosi, R.S. Kolosza, J. Daru, L. Kova Cs, L.A. Pa, Is there a relationship between cell phone use and semen quality? *Arch. Androl.* 51 (2005) 385–393.
- [42] P. Stefanis, A. Drakeley, R. Gazvani, D.I. Lewis-Jones, Growing concern over the safety of using mobile phones and male fertility, *Arch. Androl.* 52 (2006) 9–14.

- [43] O. Eroglu, E. Oztas, I. Yildirim, T. Kir, B. Aydur, G. Komesli, H.C. Irkilata, M.K. Irmak, A.F. Peker, Effects of electromagnetic radiation from a cellular phone on human sperm motility: an in vitro study, *Arch. Med. Res.* 37 (2006) 840–843.
- [44] A. Agarwal, F. Deepinder, R.K. Sharma, G. Ranga, J. Li, Effect of cell phone usage on semen analysis in men attending infertility clinic: an observational study, *Fertil. Steril.* 89 (2008) 124–128.
- [45] A. Wdowiak, L. Wdowiak, H. Wiktor, Evaluation of the effect of using mobile phones on male fertility, *Ann. Agric. Environ. Med.* 14 (1) (2007) 169–172.
- [46] J.G. Yan, A.M. Grestl, T. Bruce, Y.H. Yan, A. Granlund, H.S. Matlob, Effects of cellular phone emissions on sperm motility in rats, *Fertil. Steril.* 88 (4) (2007) 957–964.
- [47] P.F. Doherty, T.C. Grubb, Effects of high-voltage power lines on birds breeding within the powerlines' electromagnetic fields, *Sialia* 18 (1996) 129–134.
- [48] V. Garaj-Vrhovac, D. Horvat, Z. Koren, The relationship between colony-forming ability, chromosome aberrations and incidence of micronuclei in V79 Chinese hamster cells exposed to microwave radiation, *Mutat. Res.* 263 (1991) 143–149.
- [49] H. Lai, N.P. Singh, Acute low-intensity microwave exposure increases DNA single-strand breaks in rat brain cells, *Bioelectromagnetics* 16 (1995) 207–210.
- [50] S. Balode, Assessment of radio-frequency electromagnetic radiation by the micronucleus test in bovine peripheral erythrocytes, *Sci. Total Environ.* 180 (1996) 81–85.
- [51] I. Belyaev, L. Hillert, E. Markova, R. Sarimov, L. Malmgren, B. Persson, M. Harms-Ringdahl, Microwaves of mobile phones affect human lymphocytes from normal and hypersensitive subjects dependent on frequency, in: 26th Annual Meeting of the Bioelectromagnetics (BEMS), Washington, USA; 2004.
- [52] G. Demisa, D. Vlastos, D.P. Mathopoulos, Effect of 910-MHz electromagnetic field on rat bone marrow, *Sci. World J.* 4 (2004) 48–54.
- [53] Reflex, 2004, <<http://www.verum-foundation.de/cgi-bin/content.cgi?id=cuprojekt01>>.
- [54] E. Diem, C. Schwarz, F. Adlkofer, O. Jahn, H. Rudiger, Non-thermal DNA breakage by mobile-phone radiation (1800 MHz) in human fibroblasts and in transformed GFSH-R17 rat granulosa cells in vitro, *Mut. Res.* 583 (2005) 178–183.
- [55] A.G. Gandhi, P. Singh, Cytogenetic damage in mobile phone users: preliminary data, *Int. J. Hum. Genet.* 5 (2005) 259–265.
- [56] A. Di Carlo, N. Wite, F. Guo, P. Garrett, T. Litovitz, Chronic electromagnetic field exposure decreases HSP70 levels and lowers cytoprotection, *J. Cell. Biochem.* 84 (2002) 447–454.
- [57] D. Leszczynski, S. Joensuu, J. Reivinen, R. Kuokka, Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: molecular mechanism for cancer- and blood-brain barrier-related effects, *Differentiation* 70 (2002) 120–129.
- [58] E. Berman, L. Chacon, D. House, B.A. Koch, W.E. Koch, J. Leal, S. Lovtrup, E. Mantiply, A.H. Martin, G.I. Martucci, K.H. Mild, J.C. Monahan, M. Sandstrom, K. Shamsaifar, R. Tell, M.A. Trillo, A. Ubeda, P. Wagner, Development of chicken embryos in a pulsed magnetic field, *Bioelectromagnetics* 11 (1990) 169–187.
- [59] B.J. Youbicier-Simo, M. Bastide, Pathological effects induced by embryonic and postnatal exposure to EMFs radiation by cellular mobile phones, *Radiat. Protect. I* (1999) 218–223.
- [60] A. Ubeda, J. Leal, M.A. Trillo, M.A. Jimenez, J.M.R. Delgado, Pulse shape of magnetic fields influences chick embryogenesis, *J. Anat.* 137 (1983) 513–536.
- [61] J.M. Farrel, T.L. Litovitz, M. Penafiel, The effect of pulsed and sinusoidal magnetic fields on the morphology of developing chick embryos, *Bioelectromagnetics* 18 (1997) 431–438.
- [62] Ju.G. Grigoriev, Influence of the electromagnetic field of the mobile phones on chickens embryo, to the evaluation of the dangerousness after the criterion of this mortality, *J. Radiat. Biol.* 5 (2003) 541–544.
- [63] F. Batellier, I. Couty, D. Picard, J.P. Brillard, Effects of exposing chicken eggs to a cell phone in "call" position over the entire incubation period, *Theriogenology* 69 (2008) 737–745.
- [64] L. Veterány, A. Veterányová, J. Jedlická, Effect of magnetic field on embryonic mortality, *Cesk. Fyziol.* 50 (2001) 141–143.
- [65] K.B. Hotary, K.R. Robinson, Evidence of a role for endogenous electrical fields in chick embryo development, *Development* 114 (1992) 985–996.
- [66] M. Mevissen, M. Haübler, Acceleration of mammary tumorigenesis by exposure of 7,12-dimethylbenz(a)anthracene-treated female rats in a 50-Hz, 100- μ T magnetic field: replication study, *J. Toxicol. Environ. Health, Part A* 53 (1998) 401–418.
- [67] D. Flipo, M. Fournier, C. Benquet, P. Roux, C. Le Boulaire, Increased apoptosis, changes in intracellular Ca^{2+} , and functional alterations in lymphocytes and macrophages after in vitro exposure to static magnetic field, *J. Toxicol. Environ. Health, Part A* 54 (1998) 63–76.
- [68] M. Fedrowitz, K. Kamino, W. Löscher, Significant differences in the effects of magnetic field exposure on 7,12 dimethylbenz(a)anthracene-induced mammary carcinogenesis in two sub-strains of Sprague-Dawley rats, *Cancer Res.* 64 (2004) 243–251.
- [69] A.H. Frey, S.R. Feld, Avoidance by rats of illumination with low power nonionizing electromagnetic energy, *J. Comp. Physiol. Psychol.* 89 (1975) 183–188.
- [70] M.A. Navakatikian, L.A. Tomashevskaya, Phasic behavioral and endocrine effects of microwaves of nonthermal intensity, in: D.O. Carpenter (Ed.), *Biological Effects of Electric and Magnetic Fields*, J. Academic Press, San Diego, CA, 1994.
- [71] D.D. Krstić, B.J. Dindić, D.T. Sokolović, V.V. Marković, D.M. Petković, S.B. Radić, The results of experimental exposition of mice by mobile telephones, in: *TELSIKS Conference, Serbia and Montenegro, Microw. Rev.* (2005) 34–37.
- [72] I.U.G. Grigoriev, S.N. Luk'ianova, V.P. Makarov, V.V. Ryskov, N.V. Moiseeva, Motor activity of rabbits in conditions of chronic low-intensity pulse microwave irradiation, *Radiat. Biol. Radioecol.* 35 (1995) 29–35.
- [73] B. Nicholls, P.A. Racey, Bats avoid radar installations: Could electromagnetic fields deter bats from colliding with wind turbines? *PLOS One* 3 (2007) e297.
- [74] A. Balmori Murciélago rabudo-*Tadarida teniotis*, En: *Enciclopedia Virtual de los Vertebrados Españoles*, Carrascal, L.M., Salvador, A. (Eds.), Museo Nacional de Ciencias Naturales, Madrid, 2004c, <<http://www.vertebradosibericos.org/>>.
- [75] T.A. Marks, C.C. Ratke, W.O. English, Strain voltage and developmental, reproductive and other toxicology problems in dogs, cats and cows: a discussion, *Vet. Human Toxicol.* 37 (1995) 163–172.
- [76] W. Löscher, G. Käs, Conspicuous behavioural abnormalities in a dairy cow herd near a TV and radio transmitting antenna, *Pract. Vet. Surg.* 29 (1998) 437–444.
- [77] A. Yurekli, M. Ozkan, T. Kalkan, H. Srybatili, H. Tuncel, P. Atukeren, K. Gumustas, S. Seker, GSM Base Station Electromagnetic Radiation and Oxidative Stress in Rats, *Electromagn. Biol. Med.* 25 (2006) 177–188.
- [78] S. Tofani, G. Agnesod, P. Ossola, S. Ferrini, R. Bussi, Effects of continuous low-level exposure to radio-frequency radiation on intrauterine development in rats, *Health Phys.* 51 (1986) 489–499.
- [79] T.P. Moorhouse, D.W. Macdonald, Indirect negative impacts of radio-collaring: sex ratio variation in water voles, *J. Appl. Ecol.* 42 (2005) 91.
- [80] R.F. McGivern, R.Z. Sokol, W.R. Adey, Prenatal exposure to a low-frequency electromagnetic field demasculinizes adult scent marking behavior and increases accessory sex organ weights in rats, *Teratology* 41 (1990) 1–8.
- [81] G.M. Lee, R.R. Neutra, L. Hristova, M. Yost, R.A. Hiatt, A Nested Case-Control Study of Residential and Personal Magnetic Field Measures and Miscarriages, *Epidemiology* 13 (2002) 21–31.

- [82] A.Y. Rezk, K. Abdulqawi, R.M. Mustafa, T.M. Abo El-Azm, H. Al-Inany, Fetal and neonatal responses following maternal exposure to mobile phones, *Saudi Med. J.* 29 (2008) 218–223.
- [83] L.G. Salford, A.E. Brum, J.L. Eberhardt, L. Malmgren, B.R. Persson, Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones, *Environ. Health Perspect.* 111 (2003) 881–893.
- [84] D. Adang, B. Campo, A.V. Vorst, Has a 970 MHz Pulsed exposure an effect on the memory related behaviour of rats? in: *The 9th European Conference on Wireless Technology*, September 2006, 2006, pp.
- [85] A.A. Kolodynski, V.V. Kolodynska, Motor and psychological functions of school children living in the area of the Skrunda Radio Location Station in Latvia, *Sci. Total Environ.* 180 (1996) 87–93.
- [86] H.A. Divan, L. Kheifets, C. Obel, J. Olsen, Prenatal and postnatal exposure to cell phone use and behavioral problems in children, *Epidemiology* 19 (2008) 523–529.
- [87] K. Mann, J. Roschke, Effects of pulsed high-frequency electromagnetic fields on human sleep, *Neuropsychobiology* 33 (1996) 41–47.
- [88] A.V. Kramarenko, U. Tan, Effects of high-frequency electromagnetic fields on human EEG: a brain mapping study, *Int. J. Neurosci.* 113 (2003) 1007–1019.
- [89] A.A. Marino, E. Nilsen, C. Fritel, Nonlinear changes in brain electrical activity due to cell phone radiation, *Bioelectromagnetics* 24 (2003) 339–346.
- [90] A.R. Blaustein, P.T.J. Johnson, Explaining frog deformities, *Sci. Am.* 288 (2003) 60–65.
- [91] A. Balmori, The incidence of electromagnetic pollution on the amphibian decline: is this an important piece of the puzzle? *Toxicol. Environ. Chem.* 88 (2006) 287–299.
- [92] W.C. Levengood, A new teratogenic agent applied to amphibian embryos, *J. Embryol. Exp. Morphol.* 21 (1969) 23–31.
- [93] R.H. Landesman, W. Scott Douglas, Abnormal limb regeneration in adult newts exposed to a pulsed electromagnetic field, *Teratology* 42 (1990) 137–145.
- [94] N.M. Grefner, T.L. Yakovleva, I.K. Boreysha, Effects of electromagnetic radiation on tadpole development in the common frog (*Rana temporaria* L.), *Russ. J. Ecol.* 29 (1998) 133–134.
- [95] A. Balmori, in preparation: Phone masts effects on common frog (*Rana temporaria*) tadpoles: An experiment in the city. See the video clips in: <http://www.hese-project.org/hese-uk/en/issues/nature.php?id=frogs>.
- [96] U. Wamke, Bienen, vögel und menschen, Die Zerstörung der Natur durch "Elektrosmog". Kompetenzinitiative, 2007 46 pp.
- [97] R.L. Carpenter, E.M. Livstone, Evidence for nonthermal effects of microwave radiation: Abnormal development of irradiated insect pupae, *IEEE Trans. Microw. Theor. Tech.* 19 (1971) 173–178.
- [98] D. Weisbrot, H. Lin, L. Ye, M. Blank, R. Goodman, Effects of mobile phone radiation on reproduction and development in *Drosophila melanogaster*, *J. Cell. Biochem.* 89 (2003) 48–55.
- [99] D.J. Panagopoulos, A. Karabourounis, L.H. Margaritis, Effect of GSM 900 MHz Mobile Phone Radiation on the Reproductive Capacity of *Drosophila melanogaster*, *Electromagn. Biol. Med.* 23 (2004) 29–43.
- [100] D.J. Panagopoulos, E.D. Chavdoula, A. Karabourounis, L.H. Margaritis, Comparison of bioactivity between GSM 900 MHz and DCS 1800 MHz mobile telephony radiation, *Electromagn. Biol. Med.* 26 (2007) 33–44.
- [101] A. Balmori, Efectos de las radiaciones electromagnéticas de la telefonía móvil sobre los insectos, *Ecosistemas* (2006).
- [102] H. Stever, J. Kuhn, C. Otten, B. Wunder, W. Harst, Verhaltensänderung unter elektromagnetischer Exposition. Pilotstudie, Institut für mathematik, Arbeitsgruppe, Bildungsinformatik, Universität Koblenz-Landau, 2005.
- [103] V.G. Balodis, K. Brumelis, O. Kalvickis, D. Nikodemus, V.Z. y Tjarve, Does the Skrunda radio location station diminish the radial growth of pine trees? *Sci. Total Environ.* 180 (1996) 57–64.
- [104] T. Selga, M. Selga, Response of *Pinus Sylvestris* L. needles to electromagnetic fields. Cytological and ultrastructural aspects, *Sci. Total Environ.* 180 (1996) 65–73.
- [105] I. Magone, The effect of electromagnetic radiation from the Skrunda Radio Location Station on *Spirodela polyrhiza* (L.) Schleiden cultures, *Sci. Total Environ.* 180 (1996) 75–80.
- [106] D.D. Sandu, C. Goiceanu, A. Ispas, I. Creanga, S. Miclaus, D.E. Creanga, A preliminary study on ultra high frequency electromagnetic fields effect on black locust chlorophylls, *Acta Biol. Hung.* 56 (2005) 109–117.
- [107] D. Roux, A. Vian, S. Girard, P. Bonnet, F. Paladian, E. Davies, G. Ledoigt, High frequency (900 MHz) low amplitude (5 V m^{-1}) electromagnetic field: a genuine environmental stimulus that affects transcription, translation, calcium and energy charge in tomato, *Planta* 227 (2007) 883–891.
- [108] M. Tkalec, K. Malarik, B. Pevalak-Kozlina, Exposure to radiofrequency radiation induces oxidative stress in duckweed *Lemna minor* L., *Sci. Total Environ.* 388 (2007) 78–89.
- [109] A. Balmori, ¿Pueden afectar las microondas pulsadas emitidas por las antenas de telefonía a los árboles y otros vegetales? *Ecosistemas* (2004), http://www.revistaecosistemas.net/articulo.asp?id=29&Id.Categoria=1&tipo=otros_contenidos.
- [110] V. Schorpp, 2007, <<http://www.puls-schlag.org/baumschaeden.htm#linden>>.
- [111] A. Balmori, Posibles efectos de las ondas electromagnéticas utilizadas en la telefonía inalámbrica sobre los seres vivos, *Ardeola* 51 (2004) 477–490.

FM-radio and TV tower signals can cause spontaneous hand movements near moving RF reflector

Paavo Huttunen^{a,*}, Osmo Hänninen^b, Risto Myllylä^a

^a Laboratory of Optoelectronics and Measurement Technology, University of Oulu, P.O. Box 4500, 90014 Oulu, Finland

^b Department of Physiology, University of Kuopio, P.O. Box 1627, 70211 Kuopio, Finland

Received 2 January 2009; received in revised form 23 January 2009; accepted 30 January 2009

Abstract

For testing human sensitivity to radio frequency (RF) standing waves a movable reflecting wall was constructed. Radio waves from the radio-TV tower reflected back and formed a standing wave near the reflector. When the reflector was moved, the position of the maximums of the standing waves changed and the electromagnetic intensity changed in the body of the standing test subject. The computer with an AD-converter registered the signals of the hand movement transducer and the RF-meter with 100 MHz dipole antennas. A total of 29 adults of different ages were tested. There were 9 persons whose hand movement graphs included features like the RF-meter. Six showed responses that did not correlate with the RF-meter. There were also 14 persons who did not react at all. Sensitive persons seem to react to crossing standing waves of the FM-radio or TV broadcasting signals.

© 2009 Elsevier Ireland Ltd. All rights reserved.

Keywords: Sensorimotor responses; Radio frequency standing waves

1. Introduction

Radio frequency radiation (RFR) has been studied intensively in the near GHz region. Subjective symptoms, sleeping problems and cognitive performance have been reported in subjects living near mobile phone base stations [1]. In the recent past, frequencies of FM-radio and television (TV) signals have been much less studied even though these frequencies cause biological and health effects, too. The whole body resonance frequency of an average man and thus the maximum absorption of RF energy occur at 70–80 MHz [2]. This is near the frequencies used in very high frequency (VHF) broadcasting. The head and limbs absorb much more energy than the torso at frequencies above body resonance [3]. Greatest absorption in the head region of man occurs at a frequency of about 375 MHz [4]. Absorption is stronger for wave propagation from head to toe than it is when the electric field is parallel to the long axis. The authors [4] believed that the enhanced absorption in the head region may make

head resonance significant in the study of behavioral effects, blood–brain barrier permeability, cataractogenesis, and other microwave bioeffects. Even increased health risks like cancer, especially melanoma incidence, near FM broadcasting and television transmitters have been reported [5,6].

Nerve impulses initiate muscle contraction by calcium ion release from the sarcoplasmic reticulum, which takes place when electric nerve signals reach the plasma membrane and T-tubules of muscle fibers [7]. Voltage dependent Ca-channels open. Acetylcholine esterase (AChE) breaks down the acetylcholine, and Na-channels close [7]. It has been reported that the number of Ca²⁺ ions liberated from hen's frontal brain depends on the modulation frequency of the weak VHF radiation; with a maximum at a frequency of 16 Hz, while an unmodulated field causes no ion release [2,8]. Multiple RF power-density windows in calcium ion release from brain tissue have presented [9]. A significant decrease in AChE activity has been found in rats exposed to radio frequency radiation of 147 MHz and its sub-harmonics 73.5 and 36.75 MHz amplitude modulated at 16 and 76 Hz. A decrease in AChE activity was independent of carrier wave frequencies [10].

* Corresponding author.

E-mail address: paavo.huttunen@elisanet.fi (P. Huttunen).




Fig. 1. Testing human radio wave sensitivity. Radio waves from the TV tower reflect back from the reflector and form a standing wave. When the reflector moves, the position of the maximums of the standing wave change, and the electromagnetic intensity changes in the body of the test subject. The computer with an AD-converter registers the signals of hand movement transducer and the RF-meter with the dipole antennas.

As there is previous evidence from human and animal studies that electromagnetic irradiation has effects in the brain, the aim of the present study was to find out, if the motor responses are generated in sensitive persons, when they move across a set of standing waves caused by radiation of a FM-radio and TV tower. The connection between the hand movements and the integrated intensity of electromagnetic field of FM-radio broadcasting were recorded.

2. Methods

The wavelength of a 100-MHz radio wave is 3 m. For testing human sensitivity to moving standing waves a movable reflecting wall with wooden frame 3 m height and 5 m wide was constructed (Fig. 1). Steel net of 20 mm × 20 mm mesh was used. Five horizontal net slices of 60 cm wide were bound together with steel wire forming a radio waves reflecting surface. The test place was 5 km from the FM-radio tower. The frame was placed in an open field perpendicular to the incoming wave. The test subject was standing back towards the frame, and he had the hand movement transducer in his hands. The RF-meter with horizontal dipole antenna was close behind him. When started, the frame was 2 m from his back and it was moved 20 m forth and back. The computer registered both signals. The method and the aim of the test were at first presented, in brief, to the test persons. All together 29 adult persons of different ages were tested. They were participants in a seminar relating to effects of electric fields, and thus they possibly do not represent a normal population.

The broadband (30–300 MHz) RF-meter and the hand movement transducer were constructed for this study by the authors. The signals were digitised by Pico high resolution




Fig. 2. Hand movements near the moving RF reflector. The standing waves moved slowly with the reflector. Intensity of the electric field was measured with the broadband RF-meter with horizontal dipole antennas. Variation of the field intensity is presented in the upper curve and the hand movements of the standing test person are in the lower curve.

data logger (ADC16). The radio frequency spectrum was measured using a spectrum analyser (GW instek GSP-827, 2.7 GHz) with 1.5 m horizontal dipole antennas. When measured, the antenna was fastened to a wooden frame 1 m from the ground.

3. Results and discussion

Results on the movable frame showed different hand movement reactions of the test subjects. There were 9 persons who reacted like the RF-meter (Fig. 2), 6 persons whose graphs, though obvious, showed no correlation to the RF-meter and 14 persons who did not react or showed only small noise like changes in their graphs (Table 1). Spectrum at the test place contains mainly the FM-radio broadcasting signals and four digital TV signals (Fig. 3). Most prominent (85 dB μ V, approximately 50 mV/m) are the 6 horizontally polarized FM-radio signals (Fig. 4).

Resonances in body parts affects the power absorption. Theoretically, the optimal length of a thin antenna in radio-frequency reception is nearly half of the wavelength of the

Table 1
Reactions to standing waves of FM-radio signals. Classification of results of 29 tested persons. Test subject was standing and the radio wave reflector was moved behind him/her. The hand movement graphs were compared to the graphs of the broadband radio frequency (RF) meter.

Reactions to standing waves	9 persons	Hand movement graphs include features like graphs of RF-meter.
Possible reaction	6 persons	Changes in the graphs but no correlation to RF-meter.
No reaction	14 persons	Only small noise like changes in the graphs.




Fig. 3. Spectrum 1–1000 MHz at the test place. The highest peaks at the left are FM-radio broadcasting signals and the four lower peaks in the middle are the digital TV signals. Because the measurement was made with 1.5 m dipoles, signals near 100 MHz are more prominent because of antenna resonance.

incoming radio wave. The experimental maximum whole body resonance frequency is lower than the resonance frequency for an ideal half wave dipole antenna [11]. The whole body resonance length of a human at the frequencies of 80–108 MHz applied to FM broadcasting is about 1.1–1.5 m. Because in this experiment the test subjects were standing and the 100 MHz FM-radio signals and TV signals at higher frequencies are horizontally polarized, the absorption is obviously higher in the shoulder area. The distance between two maximums of the 100 MHz standing wave is 1.5 m. The half




Fig. 4. Spectrum of the FM-radio broadcasting at the test place. Six channels were sending and the maximum electric field intensity was 85 dB μ V.

waves of local digital TV signals (500–700 MHz) are only about 20–30 cm. This means that there can be many maximums of standing waves of TV signals in the body at the same time, even near the reflector.

The biggest variation in the local field intensity was caused by the FM broadcasting. There were 6 channels in the tower. Because of different wave lengths, the standing waves near the reflector are at the same phase and they amplify each other, but further away, the phases are mixed and so the amplitude of the summed standing waves is smaller.

With this experiment, we cannot exactly say where the reaction occurs, in limbs, muscles or in the head. It is possible that a change of intensity in standing radiowaves causes a small change in the nerve-muscle permeability of the nerve signal. The person feels it like a spontaneous muscle contraction. His hands are moving away and closer when the standing waves are passing. By some persons, the distance from hand to hand varied 0–60 cm. That means that some of muscles in arms and shoulders should react.

The spectrum contains many frequencies of electromagnetic radiation. The radiation is not only coming from the nearest tower, and it is impossible to clean the test area from other waves. This experiment was made at rural area, but even there, the private hand held telephone signals cause interferences to RF-instruments.

4. Conclusions

Sensitive persons seem to react to crossing standing waves of the FM-radio or TV broadcasting signals. The reactions were apparently initiated by RFR near reflecting objects, but they became more random in very weak variations of total field intensity. In any case, individuals are different, and in natural situations many sources interfere with each other.

References

- [1] H.-P. Hunter, H. Moshammer, P. Wallner, M. Kundl, Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations, *Occup. Environ. Med.* 63 (2006) 307–313.
- [2] R.V. Adey, Frequency and power windowing in tissue interactions with weak electromagnetic fields, *Proc. IEEE* 68 (1) (1980) 119–125.
- [3] R.G. Olsen, T.A. Griner, Specific absorption rate in models of man and monkey at 225 and 2,000 MHz, *Bioelectromagnetics* 8 (1987) 377–384.
- [4] M.J. Hagmann, O.P. Gandhi, J.A. D'andrea, I. Chatterjee, Head resonance: numerical solutions and experimental results, *IEEE Trans. Microw. Theory Tech.* MTT-27 (9) (1979) 809–813.
- [5] H.G. Dolk, P. Shaddik, Walls, et al., Cancer incidence near radio and television transmitters in Great Britain I and II, *Am.J. Epidemiol.* 145 (1997) 1–17.
- [6] O. Hallberg, O. Johansson, Melanoma incidence and frequency modulation (FM) broadcasting, *Arch. Environ. Health* 57 (1) (2002) 32–40.
- [7] B. Alberts, D. Bray, J. Lewis, M. Raff, K. Roberts, J.D. Watson (Eds.), *Molecular Biology of the Cell, The Nervous System*, Garland Publishing Inc., New York/London, 1983, pp. 1013–1098 (Chapter 18).

- [8] C.F. Blackman, J.A. Elder, C.M. Weil, S.G. Benane, D.C. Eichinger, D.E. House, Induction of calcium-ion efflux from brain tissue by radio-frequency radiation: effects of modulation frequency and field strength, *Radio Sci.* 14 (1979) 93–98.
- [9] C. Blackman, L.S. Kinney, D.E. House, W.T. Joines, Multiple power-density windows and their possible origin, *Bioelectromagnetics* 10 (2) (1989) 115–128.
- [10] K.K. Kunjilwar, J. Behari, Effects of amplitude-modulated radio frequency radiation on cholinergic system of developing rats, *Brain Res.* 601 (1–2) (1993) 321–324.
- [11] O.P. Gandhi, State of the knowledge for electromagnetic absorbed dose in man and animals (1980), *Proc. IEEE* 68 (1980) 24–32.

Cell phone radiation: Evidence from ELF and RF studies supporting more inclusive risk identification and assessment[☆]

Carl Blackman

Raleigh, NC 27607, USA

Received 21 January 2009; received in revised form 3 February 2009; accepted 3 February 2009

Abstract

Many national and international exposure standards for maximum radiation exposure from the use of cell phone and other similar portable devices are ultimately based on the production of heat particularly in regions of the head, that is, thermal effects (TE). The recent elevation in some countries of the allowable exposure, that is, averaging the exposure that occurs in a 6 min period over 10 g of tissue rather than over 1 g allows for greater heating in small portions of the 10-g volume compared to the exposure that would be allowed averaged over 1-g volume. There is concern that 'hot' spots, that is, momentary higher intensities, could occur in portions of the 10-g tissue piece, might have adverse consequences, particularly in brain tissue.

There is another concern about exposure to cell phone radiation that has been virtually ignored except for the National Council of Radiation Protection and Measurements (NCRP) advice given in a publication in 1986 [National Council for Radiation Protection and Measurements, Biological Effects and Exposure Criteria for Radiofrequency Electromagnetic Fields, National Council for Radiation Protection and Measurements, 1986, 400 pp.]. This NCRP review and guidance explicitly acknowledge the existence of non-thermal effects (NTE), and included provisions for reduced maximum-allowable limits should certain radiation characteristics occur during the exposure.

If we are to take most current national and international exposure standards as completely protective of thermal injury for acute exposure only (6 min time period) then the recent evidence from epidemiological studies associating increases in brain and head cancers with increased cell phone use per day and per year over 8–12 years, raises concerns about the possible health consequences on NTE first acknowledged in the NCRP 1986 report [National Council for Radiation Protection and Measurements, Biological Effects and Exposure Criteria for Radiofrequency Electromagnetic Fields, National Council for Radiation Protection and Measurements, 1986, 400 pp.].

This paper will review some of the salient evidence that demonstrates the existence of NTE and the exposure complexities that must be considered and understood to provide appropriate, more thorough evaluation and guidance for future studies and for assessment of potential health consequences. Unfortunately, this paper is necessary because most national and international reviews of the research area since the 1986 report [National Council for Radiation Protection and Measurements, Biological Effects and Exposure Criteria for Radiofrequency Electromagnetic Fields, National Council for Radiation Protection and Measurements, 1986, 400 pp.] have not included scientists with expertise in NTE, or given appropriate attention to their requests to include NTE in the establishment of public-health-based radiation exposure standards. Thus, those standards are limited because they are not comprehensive.

© 2009 Elsevier Ireland Ltd. All rights reserved.

Keywords: Non-thermal effects; Electromagnetic fields; Exposure standards

1. Introduction

1.1. The current approach to exposure limits (based on heating and electric current flow in tissues)

It is universally accepted that radiofrequency radiation (RFR) can cause tissue heating (thermal effects, TE) and that extremely low-frequency (ELF) fields, e.g., 50

[☆] Disclaimer: The opinions expressed in this text are those of its author, and are not necessarily those of his employer, the U.S. Environmental Protection Agency.

E-mail address: Carl.Blackman@gmail.com.

and 60 Hz, can cause electrical current flows that shock and even damage or destroy tissues. These factors alone are the underlying bases for present exposure standards. EMF exposures that cause biological effects at intensities that do not cause obvious thermal changes, that is, non-thermal effects (NTE), have been widely reported in the scientific literature since the 1970s including beneficial applications in development and repair processes. The current public safety limits do not take modulation into account and thus are no longer sufficiently protective of public health where chronic exposure to pulsed or pulse-modulated signal is involved, and where sub-populations of more susceptible individuals may be at risk from such exposures.

1.2. Modulation as a critical element

Modulation signals are one important component in the delivery of EMF signals to which cells, tissues, organs and individuals can respond biologically. At the most basic level, modulation can be considered a pattern of pulses or repeating signals which have specific meaning in defining that signal apart from all others. Modulated signals have a specific 'beat' defined by how the signal varies periodically or aperiodically over time. Pulsed signals occur in an on-off pattern, which can be either smooth and rhythmic, or sharply pulsed in quick bursts. Amplitude and frequency modulation involves two very different processes where the high-frequency signal, called the carrier wave, has a lower frequency signal that is superimposed on or 'rides' on the carrier frequency. In amplitude modulation, the lower frequency signal is embedded on the carrier wave as changes in its amplitude as a function of time, whereas in frequency modulation, the lower frequency signal is embedded as slight changes in the frequency of the carrier wave. Each type of low-frequency modulation conveys specific 'information', and some modulation patterns are more effective (more bioactive) than others depending on the biological reactivity of the exposed material. This enhanced interaction can be a good thing for therapeutic purposes in medicine, but can be deleterious to health where such signals could stimulate disease-related processes, such as increased cell proliferation in precancerous lesions. Modulation signals may interfere with normal, non-linear biological functions. More recent studies of modulated RF signals report changes in human cognition, reaction time, brain-wave activity, sleep disruption and immune function. These studies have tested the RF and ELF-modulated RF signals from emerging wireless technologies (cell phones) that rely on pulse-modulated RF to transmit signals. Thus modulation can be considered as information content embedded in the higher frequency carrier wave that may have biological consequences beyond any effect from the carrier wave directly.

In mobile telephony, for example, modulation is one of the underlying ways to categorize the radiofrequency signal

of one telecom carrier from another (TDMA from CDMA from GSM). Modulation is likely a key factor in determining whether and when biological reactivity might be occurring, for example in the new technologies which make use of modulated signals, some modulation (the packaging for delivery for an EMF 'message') may be bioactive, for example, when frequencies are similar to those found in brain wave patterns. If a new technology happens to use brain wave frequencies, the chances are higher that it will have effects, in comparison, for example, to choosing some lower or higher modulation frequency to carry the same EMF information to its target.

This chapter will show that other EMF factors may also be involved in determining if a given low-frequency signal directly, or as a modulation of a radiofrequency wave, can be bioactive. Such is the evolving nature of information about modulation. It argues for great care in defining standards that are intended to be protective of public health and well-being. This chapter will also describe some features of exposure and physiological conditions that are required in general for non-thermal effects to be produced, and specifically to illustrate how modulation is a fundamental factor which should be taken into account in public safety standards.

2. Laboratory evidence

Published laboratory studies have provided evidence for more than 40 years on bioeffects at much lower intensities than cited in the various widely publicized guidelines for limits to prevent harmful effects. Many of these reports show EMF-caused changes in processes associated with cell growth control, differentiation and proliferation, that are biological processes of considerable interest to physicians for potential therapeutic applications and for scientists who study the molecular and cellular basis of cancer. EMF effects have been reported in gene induction, transmembrane signaling cascades, gap junction communication, immune system action, rates of cell transformation, breast cancer cell growth, regeneration of damaged nerves and recalcitrant bone-fracture healing. These reports have cell growth control as a common theme. Other more recent studies on brainwave activity, cognition and human reaction time lend credence to modulation (pulsed RF and ELF-modulated RF) as a concern for wireless technologies, most prominently from cell phone use.

In the process of studying non-thermal biological effects, various exposure parameters have been shown to influence whether or not a specific EMF can cause a biological effect, including intensity, frequency, the co-occurrence of the static magnetic field (both the natural earth's magnetic field and anthropogenic fields), the presence of the electrical field, the magnetic field, or their combination, and whether EMF is sinusoidal, pulsed or in more com-

plex wave forms. These parameters will be discussed below.

Experimental results will be used to illustrate the influence of each EMF parameter, while also demonstrating that it is highly unlikely the effects are due to EMF-caused current flow or heating.

2.1. Initial studies that drew attention to NTE

Several papers in the 1960s and early 1970s reported that ELF fields could alter circadian rhythms in laboratory animals and humans. In the latter 1960s, a paper by Hamer [2] reported that the EMF environment in planned space capsules could cause human response time changes, i.e., the interval between a signal and the human response. Subsequent experiments by a research group led by Adey were conducted with monkeys, and showed similar response time changes and also EEG pattern changes [3,4]. The investigators shifted the research subject to cats and decided they needed to use a radiofrequency field to carry the ELF signal into the cat brain, and observed EEG pattern changes, ability to sense and behaviorally respond to the ELF component of RFR, and the ability of minor electric current to stimulate the release of an inhibitory neurotransmitter, GABA, and simultaneous release of a surrogate measure, calcium ions, from the cortex [5,6]. At this time Bawin, a member of the research group, adopted newly hatch chickens as sources of brain tissue and observed changes in the release of calcium ions from in vitro specimens as a function of ELF frequency directly or as amplitude modulation ('am') of RFR (RFRam) [7–11]. Tests of both EMF frequency and intensity dependences demonstrated a single sensitive region (termed 'window') over the range of frequency and intensity examined. This series of papers showed that EMF-induced changes could occur in several species (human, monkey, cat and chicken), that calcium ions could be used as surrogate measures for a neurotransmitter, that ELF fields could produce effects similar to RFRam (note: without the 'am', there was no effect although the RFR intensity was the same), and that the dose and frequency response consisted of a single sensitivity window.

Subsequent, independent research groups published a series of papers replicating and extending this earlier work. Initial studies by Blackman, Joines and colleagues [12–25] used the same chick brain assay system as Bawin and colleagues. These papers reported multiple windows in intensity and in frequency within which calcium changes were observed in the chick brain experimental systems under EMF exposure. Three other independent groups offered confirmation of these results by reporting intensity and frequency windows for calcium, neurotransmitter or enolase release under EMF exposure of human and animal nervous system-derived cells in vitro by Dutta et al. [26–29], of rat pancreatic tissue slices by Albert et al. [30], and of frog heart by Schwartz et al. [31] but not frog-heart

atrial strips in vitro [32]. This series of papers showed that multiple frequency and intensity windows were a common phenomenon that required the development of new theoretical concepts to provide a mechanism of action paradigm.

2.2. Refined laboratory studies reveal more details

Additional aspects of the EMF experiments with the chick brain described by Blackman and colleagues, above, also revealed critical co-factors that influenced the action of EMF to cause changes in calcium release, including the influence of the local static magnetic field, and the influence of physico-chemical parameters, such as pH, temperature and the ionic strength of the bathing solution surrounding the brain tissue during exposure. This information provides clues for and constraints on any theoretical mechanism that is to be developed to explain the phenomenon. Most current theories ignore these parameters that need to be monitored and controlled for EMF exposure to produce NTE. These factors demonstrate that the current risk assessment paradigms, which ignore them, are incomplete and thus may not provide the level of protection currently assumed.

2.3. Sensitivity of developing organisms

An additional study was also conducted to determine if EMF exposure of chicken eggs while the embryo was developing could influence the response of brain tissue from the newly hatched chickens. The detailed set of frequency and intensity combinations under which effects were observed, were all obtained from hatched chickens whose eggs were incubated for 21 days in an electrically heated chamber containing 60-Hz fields. Thus tests were performed to determine if the 60-Hz frequency of ELF fields (10 V/m in air) during incubation, i.e., during embryogenesis and organogenesis, would alter the subsequent calcium release responses of the brain tissue to EMF exposure. The reports of Blackman et al. [19] and Joines et al. [25] showed that the brain tissue response was changed when the field during the incubation period was 50 Hz rather than 60 Hz. This result is consistent with an anecdotal report of adult humans, institutionalized because of chemical sensitivities, who were also responsive to the frequency of power-line EM fields that were present in the countries where they were born and raised [33]. This information indicates there may be animal and human exposure situations where EMF imprinting during development could be an important factor in laboratory and epidemiological situations. EMF imprinting, which may only become manifest when a human is subjected to chemical or biological stresses, could reduce ability to fight disease and toxic insult from environmental pollution, resulting in a population in need of more medical services, with resulting lost days at work.

3. Fundamental exposure parameters—to be considered when establishing a mode (or mechanism) of action for non-thermal EMF-induced biological effects

3.1. Intensity

There are numerous reports of biological effects that show intensity “windows”, that is, regions of intensity that cause changes surrounded by higher and lower intensities that show no effects from exposure. One very clear effect by Blackman and colleagues is 16-Hz, sine wave-induced changes in calcium efflux from brain tissue in a test tube because it shows two very distinct and clearly separated intensity windows of effects surrounded by regions of intensities that caused no effects [17]. There are other reports for similar multiple windows of intensity in the radiofrequency range [22,26,29,31]. Note that calcium ions are a secondary signal transduction agent active in many cellular pathways. These results show that intensity windows exist, they display an unusual and unanticipated “non-linear” (non-linear and non-monotonic) phenomenon that has been ignored in all risk assessment and standard setting exercises, save the NCRP 1986 publication [1]. Protection from multiple intensity windows has never been incorporated into any risk assessment; to do so would call for a major change in thinking. These results mean that lower intensity is not necessarily less bioactive, or less harmful.

Multiple intensity windows appeared as an unexpected phenomenon in the late 1970s and 1980s. There has been one limited attempt to specifically model this phenomenon by Thompson et al. [34], which was reasonably successful. This modeling effort should be extended because there are publications from two independent research groups showing multiple intensity windows for 50, 147, and 450 MHz fields when amplitude modulated at 16 Hz using the calcium ion release endpoint in chicken brains, *in vitro*. The incident intensities (measured in air) for the windows at the different carrier frequencies do not align at the same values. However, Joines et al. [23,24] and Blackman et al. [20] noted the windows of intensity align across different carrier frequencies if one converts the incident intensity to the intensity expected within the sample at the brain surface. This conversion was accomplished by correcting for the different dielectric constants of the sample materials due to the different carrier frequencies. The uniqueness of this response provides a substantial clue to theoreticians but it is interesting and disappointing that no publications have appeared attempting to address this relationship. It is obvious that this phenomenon is one that needs further study.

3.2. Frequency

Frequency-dependent phenomena are common occurrences in nature. For example, the human ear only hears a portion of the sound that is in the environment, typically from

20 to 20,000 Hz, which is a frequency “window”. Another biological frequency window can be observed for plants grown indoors. Given normal indoor lighting the plants may grow to produce lush vegetation but not produce flowers unless illuminated with a lamp that emits a different spectrum of light partially mimicking the light from the sun. Thus, frequency windows of response to various agents exist in biological systems from plants to homo sapiens.

In a similar manner, there are examples of EMF-caused biological effects that occur in a frequency-dependent manner that cannot be explained by current flow or heating. The examples include reports of calcium ion efflux from brain tissue *in vitro* by Blackman and Joines and colleagues at low frequency [15,19] and at high frequency modulated at low frequency [20,35,24]. An additional example of an unexpected result is by Liboff [36].

In addition, two apparently contradictory multiple-frequency exposure results provide examples of the unique and varied non-thermal interactions of EMF with biological systems. Litovitz and colleagues showed that an ELF sinusoidal signal could induce a biological response in a cell culture preparation, and that the addition of a noise signal of equal average intensity could block the effect caused by the sinusoidal signal, thereby negating the influence of the sinusoidal signal [37]. Similar noise canceling effects were observed using chick embryo preparations [38,39]. It was also shown that the biological effects caused by microwave exposures imitating cell phone signals could be mitigated by ELF noise [40]. However, this observation should not be generalized; a noise signal is not always benign. Milham and Morgan [41] showed that a sinusoidal ELF (60-Hz) signal was not associated with the induction of cancer in humans, but when that sinusoidal signal was augmented by a noise signal, basically transients that added higher frequencies, an increase in cancer was noted in humans exposed over the long-term. Thus, the addition of noise in this case was associated with the appearance of a health issue. Havas [42–44] has described other potential health problems associated with these higher frequency transients, termed “dirty power.” The bioactive frequency regions observed in these studies have never been explicitly considered for use in any EMF risk assessments, thus demonstrating the incomplete nature of current exposure guideline limits.

There are also EMF frequency-dependent alterations in the action of nerve growth factor (NGF) to stimulate neurite outgrowth (growth of primitive axons or dendrites) from a peripheral-nerve-derived cell (PC-12) in culture shown by Blackman et al. [45,46] and by Trillo et al. [47]. The combined effect of frequency and intensity is also a common occurrence in both the analogous sound and the light examples given above. Too much or too little of either frequency or intensity show either no or undesirable effects. Similarly, Blackman et al. [15] has reported EMF responses composed of effect “islands” of intensity and frequency combinations, surrounded by a “sea” of intensity and frequency combinations of null effects. Although the mechanisms responsible

for these effects have not been established, the effects represent a here-to-fore unknown phenomenon that may have complex ramifications for risk assessment and standard setting. Nerve growth and neurotransmitter release that can be altered by different combinations of EMF frequencies and intensities, especially in developing organisms like children, could conceivably produce over time a subsequent altered ability to successfully or fully respond behaviorally to natural stressors in the adult environment; research is urgently needed to test this possibility in animal systems.

Nevertheless, this phenomenon of frequency dependence is ignored in the development of present exposure standards. These standards rely primarily on biological responses to intensities within an arbitrarily defined engineering-based frequency bands, not biologically based response bands, and are solely based on an energy deposition determinations.

4. Static magnetic field—a completely unexpected complexity

The magnetic field of the earth at any given location has a relatively constant intensity as a function of time. However, the intensity value, and the inclination of the field with respect to the gravity vector, varies considerable over the face of the earth. More locally, these features of the earth's magnetic field can also vary by more than 20% inside manufactured structures, particularly those with steel support structures.

At the Bioelectromagnetics Society annual meeting in 1984 [48], Blackman revealed his group's discovery that the intensity of the static magnetic field could establish and define those oscillatory frequencies that would cause changes in calcium ion release in his chick brain preparation. This result was further discussed at a NATO Advanced Research workshop in Erice, Italy in the fall of 1984 and by publications from that meeting and subsequent research: Blackman et al. [14,18] and Liboff et al. [36,49,50]. Substantial additional research on this feature was reported by Liboff and colleagues [51,52,50]. Blackman et al. also reported on the importance of the relative orientation of the static magnetic field vector to the oscillating magnetic field vector [21] and demonstrated a reverse biological response could occur depending on parallel or perpendicular orientations of the static and oscillating magnetic fields [53].

There have been many attempts to explain this phenomenon by a number of research teams led by Smith [49], Blackman [15], Liboff [36,54], Lednev [55], Blanchard [56], Zhadin [57], del Giudice [58], Binhi [59–62], and Matronchik [63] but none has been universally accepted. Nevertheless, experimental results continued to report static and oscillating field dependencies for non-thermally induced biological effects in studies led by Zhadin [64,65], Vorobyov [66], Bau-reus Koch [67], Sarimov [68], Prato [69,70], Comisso [71], and Novikov [72].

With this accumulation of reports from independent, international researchers, it is now clear that if a biological

response depends on the static magnetic field intensity, and even its orientation with respect to an oscillating field, then the conditions necessary to reproduce the phenomenon are very specific and might easily escape detection (see for example, Blackman and Most [73]). The consequences of these results are that there may be exposure situations that are truly detrimental (or beneficial) to organisms, but that are insufficiently common on a large scale that they would not be observed in epidemiological studies; they need to be studied under controlled laboratory conditions to determine impact on health and wellbeing.

5. Electric and magnetic components—both biological active with different consequences

Both the electric and the magnetic components have been shown to directly and independently cause biological changes. There is one report that clearly distinguishes the distinct biological responses caused by the electric field and by the magnetic field. Marron et al. [74] show that electric field exposure can increase the negative surface charge density of an amoeba, *Physarum polycephalum*, and that magnetic field exposure of the same organism causes changes in the surface of the organism to reduce its hydrophobic character. Other scientists have used concentric growth surfaces of different radii and vertical magnetic fields perpendicular to the growth surface to determine if the magnetic or the induced electric component is the agent causing biological change. Liburdy et al. [75], examining calcium influx in lymphocytes, and Greene et al. [76], monitoring ornithine decarboxylase (ODC) activity in cell culture, showed that the induced electric component was responsible for their results. In contrast, Blackman et al. [77,78] monitoring neurite outgrowth from two different clones of PC-12 cells and using the same exposure technique used by Liburdy and by Greene showed the magnetic component was the critical agent in their experiments. EMF-induced changes on the cell surface, where it interacts with its environment, can dramatically alter the homeostatic mechanisms in tissues, whereas changes in ODC activity are associated with the induction of cell proliferation, a desirable outcome if one is concerned about wound healing, but undesirable if the concern is tumor cell growth. This information demonstrates the multiple, different ways that EMF can affect biological systems. Present analyses for risk assessment and standard setting have ignored this information, thus making their conclusions of limited value.

6. Sine and pulsed waves—like different programs on a radio broadcast station

Important characteristics of pulsed waves that have been reported to influence biological processes include the following: (1) frequency, (2) pulse width, (3) intensity, (4) rise and fall time, and (5) the frequency, if any, within the pulse ON

time. Chiabrera et al. [79] showed that pulsed fields caused de-differentiation of amphibian red blood cells. Scarfi et al. [80] showed enhanced micronuclei formation in lymphocytes of patients with Turner's syndrome (only one X chromosome) but no change in micronuclei formation when the lymphocytes were exposed to sine waves (Scarfi et al. [81]). Takahashi et al. [82] monitored thymidine incorporation in Chinese hamster cells and explored the influence of pulse frequency (two windows of enhancement reported), pulse width (one window of enhancement reported) and intensity (two windows of enhancement reported followed by a reduction in incorporation). Ubeda et al. [83] showed the influence of difference rise and fall times of pulsed waves on chick embryo development.

6.1. Importance for risk assessment

It is important to note that the frequency spectrum of pulsed waves can be represented by a sum of sine waves which, to borrow a chemical analogy, would represent a mixture of chemicals, any one of which could be biologically active. Risk assessment and exposure limits have been established for specific chemicals or chemical classes of compounds that have been shown to cause undesirable biological effects. Risk assessors and the general public are sophisticated enough to recognize that it is impossible to declare all chemicals safe or hazardous; consider the difference between food and poisons, both of which are chemicals. A similar situation occurs for EMF; it is critical to determine which combinations of EMF conditions have the potential to cause biological harm and which do not.

Obviously, pulse wave exposures represent an entire genre of exposure conditions, with additional difficulty for exact independent replication of exposures, and thus of results, but with increased opportunities for the production of biological effects. Current standards were not developed with explicit knowledge of these additional consequences for biological responses.

7. Mechanisms

Two papers have the possibility of advancing understanding in this research area. Chiabrera et al. [84] created a theoretical model for EMF effects on an ion's interaction with protein that includes the influence of thermal energy and of metabolism. Before this publication, theoreticians assumed that biological effects in living systems could not occur if the electric signal is below the signal caused by thermal noise, in spite of experimental evidence to the contrary. In this paper, the authors show that this limitation is not absolute, and that different amounts of metabolic energy can influence the amount and parametric response of biological systems to EMF. The second paper, by Marino et al. [85], presents a new analytical approach to examine endpoints in systems exposed to EMF. The authors, focusing on exposure-induced lymphoid phenotypes, report that EMF may not cause changes in the mean values of endpoints, but by using recurrence analysis, they capture exposure-induced, statistically significant, non-linear movements of the endpoints to either side of the mean endpoint value. They provide further evidence using immunological endpoints from exposed and sham treated mice [86–88]. Additional research has emerged from this laboratory on EMF-induced animal and human brain activity changes that provides more evidence for the value of their research approach (Marino et al. [89–92], Kolomytkin et al. [93] and Carrubba et al. [94–98]). Further advanced theoretical and experimental studies of relevance to non-thermal biological effects are emerging; see for example reports by Binhi et al. [59–62], Zhadin et al. [64,99,65], and Novikov et al. [72]. *It is apparent that much remains to be examined and explained in EMF biological effects research through more creative methods of analysis than have been used before. The models described above need to be incorporated into risk assessment determinations.*

phoid phenotypes, report that EMF may not cause changes in the mean values of endpoints, but by using recurrence analysis, they capture exposure-induced, statistically significant, non-linear movements of the endpoints to either side of the mean endpoint value. They provide further evidence using immunological endpoints from exposed and sham treated mice [86–88]. Additional research has emerged from this laboratory on EMF-induced animal and human brain activity changes that provides more evidence for the value of their research approach (Marino et al. [89–92], Kolomytkin et al. [93] and Carrubba et al. [94–98]). Further advanced theoretical and experimental studies of relevance to non-thermal biological effects are emerging; see for example reports by Binhi et al. [59–62], Zhadin et al. [64,99,65], and Novikov et al. [72]. *It is apparent that much remains to be examined and explained in EMF biological effects research through more creative methods of analysis than have been used before. The models described above need to be incorporated into risk assessment determinations.*

8. Problems with current risk assessments—observations of effects are segregated by artificial frequency bands that ignore modulation

One fundamental limitation of most reviews of EMF biological effects is that exposures are segregated by the physical (engineering/technical) concept of frequency bands favored by the engineering community. This is a default approach that follows the historical context established by the incremental addition of newer technologies that generate increasingly higher frequencies. However, this approach fails to consider unique responses from biological systems that are widely reported at various combinations of frequencies, modulations and intensities.

When common biological responses are observed without regard for the particular, engineering-defined EMF frequency band in which the effects occur, this reorganization of the results can highlight the commonalities in biological responses caused by exposures to EMF across the different engineering-defined frequency bands. An attempt to introduce this concept to escape the limitations of the engineering-defined structure occurred with the development of the 1986 NCRP radiofrequency exposure guidelines because published papers from the early 1970s to the mid 1980s (to be discussed below) demonstrated the need to include amplitude modulation as a factor in setting of maximum exposure limits. The 1986 NCRP guideline [1] was the one and only risk evaluation that included an exception for modulated fields.

The current research and risk assessment attempts are no longer tenable. The 3-year delay in the expected report of the 7-year Interphone study results has made this epidemiological approach a 10-year long effort, and the specific exposure conditions, due to improved technology, have changed so that the results may no longer be applicable to the current

exposure situation. It is unproductive to continue to fund epidemiological studies of people who are exposed to a wide variety of diversified, uncontrolled, and poorly characterized EMF in their natural and work environments. In place of the funding of more epidemiological studies should be funding to support controlled laboratory studies to focus on the underlying processes responsible for the NTE described above, so that mechanisms or modes of action can be developed to provide a theoretical framework to further identify, characterize and unify the action of the heretofore ignored exposure parameters shown to be important.

8.1. Potential explanation for the failure to optimize research in EMF biological effects

Unfortunately, risk evaluations following the 1986 NCRP example [1], returned to the former engineering-defined analysis conditions, in part because scientists who reported non-thermal effects were not placed on the review committees, and in the terms of Slovic [100] “Risk assessment is inherently subjective and represent a blend of science and judgment with important psychological, social, cultural, and political factors. . . . Whoever controls the definition of risk controls the rational solution to the problem at hand. . . . Defining risk is thus an exercise in power.” It appears that by excluding scientists experienced with producing non-thermal biological effects, the usually sound judgment by the selected committees was severely limited in its breadth-of-experience, thereby causing the members to retreat to their own limited areas of expertise when forced to make judgments, as described by Slovic [100], “Public views are also influenced by worldviews, ideologies, and values; so are scientists’ views, particularly when they are working at limits of their expertise.” The current practice of segregating scientific investigations (and resulting public health limits) by artificial divisions of frequency dramatically dilutes the impact of the basic science results, thereby reducing and distorting the weight of evidence in any evaluation process (see evaluations of bias by Havas [101], referring to NRC 1997 [102] compared to NIEHS 1998 [103] and NIEHS 1999 [104]).

9. Suggested research

Are there substitute approaches that would improve on the health-effects evaluation situation? As mentioned above, it may be useful in certain cases to develop a biologically based clustering of the data to focus on and enrich understanding of certain aspects of biological responses. Some examples to consider for biological clustering include: (1) EMF features, such as frequency and intensity inter-dependencies, (2) common co-factors, such as the earth’s magnetic field or co-incident application of chemical agents to perturb and perhaps sensitize the biological system to EMF, or (3) physiological state of the biological specimen, such as age or sensitive sub-populations, including genetic predisposition

as described by Fedrowitz et al. [105,106], and for human populations, recently reported by Yang et al. [107].

To determine if this approach has merit, one could combine reports of biological effects found in the ELF (including sub-ELF) band with effects found in the RF band when the RF exposures are amplitude modulated (AM) using frequencies in the ELF band. The following data should be used: (a) human response time changes under ELF exposure [2], (b) monkey response time and EEG changes under ELF exposure [3,4], (c) cat brain EEG, GABA and calcium ion changes induced by ELF and AM-RF [8,9,7,10,6,11,108,5], (d) calcium ion changes in chick brain tissue under ELF and AM-RF [8,9,7,10,13–15,21,16–18,12,19,20,22,35,23–25,11], and (e) calcium changes under AM-RF in brain cells in culture [26–28] and in frog heart under AM-RF [31]. The potential usefulness of applying biological clustering in the example given above even though AM is used, is that the results may have relevance to assist in the examination of some of the effects reportedly caused by cellular phone exposures which include more complex types of modulation of RF. This suggestion is reasonable because three groups later reported human responses to cell phone emissions that include changes in reaction times – Preece et al. [109,110], Koivisto et al. [111,112] and Krause et al. [113,114] – or to brain wave potentials that may be associated with reaction time changes—Freude et al. [115,116].

Subsequently, Preece et al. [117] tested cognitive function in children and found a trend, but not a statistically significant change in simple reaction time under exposure, perhaps because he applied a Bonferroni correction to his data (alpha for significance was required to be less than 0.0023). It would appear that a change in the experimental protocol might provide a more definitive test of the influence of exposure on simple reaction time because it is known that a Bonferroni correction is a particularly severe test of statistical significance, or as the author observed, “a particularly conservative criterion.”

Krause et al. [118] examined cognitive activity by observing oscillatory EEG activity in children exposed to cell phone radiation while performing an auditory memory task and reported exposure related changes in the ~4–8 Hz EEG frequencies during memory encoding, and changes in that range and also ~15 Hz during recognition. The investigators also examined cognitive processing, an auditory memory task or a visual working memory task, in adults exposed to CW or pulsed cell phone radiation on either the right or left side of the head, and reported modest changes in brain EEG activity in the ~4–8 Hz region, compared to CW exposure, but with caveats that no behavior changes were observed, and that the data were varying, unsystematic and inconsistent with previous reports (Krause et al. [119]). Haarala and colleagues conducted an extensive series of experiments, examining reaction time [120], short-term memory [121], short-term memory in children [122], and right versus left hemisphere exposure [123]. Although these studies did not

support the positive effects from exposure reported by others, they provided possible explanations for the apparent lack of agreement.

Other research groups have also examined the effects of cell phone radiation on the central nervous system, including Borbely et al. [124], Huber et al. [125], Loughran et al. [126], and D'Costa et al. [127], who found changes in sleep EEG patterns and other measures during or after short-term exposures, while others, such as Fritzer et al. [128] exposed for longer time periods found no changes in sleep parameters, EEG power spectra, correlation dimension nor cognitive function. The work of Pritchard [129] served as the basis to examining correlation dimensions, which is opening a potentially fertile avenue for investigation. Although this approach provides more indepth information on ongoing processes and function, it has not yet been used to address potential consequences associated with long-term cell phone use.

The papers published in the 1960s through 1991, described in earlier sections of this paper, foreshadowed the more recent publications in 1999 through 2008 showing response time changes; or associated measures; in human subjects during exposure to cell phone-generated radiation. It is unfortunate that essentially none of the earlier studies was acknowledged in these recent reports on cognition, reaction time and other measures of central nervous system processes. Without guidance from this extensive earlier work, particularly those demonstrating the variety of exposure parameter spaces that must be controlled to produce repeatable experiments, the development of the mechanistic bases for non-thermal effects from EMF exposures will be substantially delayed. The omission of the recognition of the exposure conditions that affect the biological outcomes continues as recently as the National Academy of Science 2009 publication [130] of future directions for research, which emphasizes the modest perspective in the results from committee members working at the limits of expertise, as anticipated by Slovic [100].

Let us hope that subsequent national and international committees that consider future directions for EMF research include members who have performed and reported non-thermal effects, in order to provide a broader perspective to develop programs that will more expeditiously address potential health problems as well as to provide guidance to industry on prudent procedures to establish for their technologies.

At present, we are left with a recommendation voiced in 1989 by Abelson [131] in an editorial in Science Magazine that addressed electric power-specific EMF, but is applicable to higher frequency EMF as well, to "adopt a prudent avoidance strategy" by "adopting those which look to be 'prudent' investments given their cost and our current level of scientific understanding about possible risks."

10. Conclusions

There is substantial scientific evidence that some modulated fields (pulsed or repeated signals) are bioactive, which

increases the likelihood that they could have health impacts with chronic exposure even at very low exposure levels. Modulation signals may interfere with normal, non-linear biological processes. Modulation is a fundamental factor that should be taken into account in new public safety standards; at present it is not even a contributing factor. To properly evaluate the biological and health impacts of exposure to modulated RFR (carrier waves), it is also essential to study the impact of the modulating signal (lower frequency fields or ELF-modulated RF). Current standards have ignored modulation as a factor in human health impacts, and thus are inadequate in the protection of the public in terms of chronic exposure to some forms of ELF-modulated RF signals. The current IEEE and ICNIRP standards are not sufficiently protective of public health with respect to chronic exposure to modulated fields (particularly new technologies that are pulse-modulated and heavily used in cellular telephony). The collective papers on modulation appear to be omitted from consideration in the recent WHO and IEEE science reviews. This body of research has been ignored by current standard setting bodies that rely only on traditional energy-based (thermal) concepts. More laboratory as opposed to epidemiological research is needed to determine which modulation factors, and combinations are bioactive and deleterious at low intensities, and are likely to result in disease-related processes and/or health risks; however this should not delay preventative actions supporting public health and wellness. If signals need to be modulated in the development of new wireless technologies, for example, it makes sense to use what existing scientific information is available to avoid the most obviously deleterious exposure parameters and select others that may be less likely to interfere with normal biological processes in life. The current membership on Risk Assessment committees needs to be made more inclusive, by adding scientists experienced with producing non-thermal biological effects. The current practice of segregating scientific investigations (and resulting public health limits) by artificial, engineering-based divisions of frequency needs to be changed because this approach dramatically dilutes the impact of the basic science results and eliminates consideration of modulation signals, thereby reducing and distorting the weight of evidence in any evaluation process.

References

- [1] National Council for Radiation Protection and Measurements. Biological Effects and Exposure Criteria for Radiofrequency Electromagnetic Fields, National Council for Radiation Protection and Measurements, 1986, 400 pp.
- [2] J. Hamer, Effects of low level, low frequency electric fields on human reaction time, Communications in Behavioral Biology 2 (5 part A) (1968) 217–222.
- [3] R.J. Gavalas, D.O. Walter, J. Hamer, W.R. Adey, Effect of low-level, low-frequency electric fields on eeg and behavior in macaca nemestrina, Brain Research 18 (3) (1970) 491–501.

- [4] R. Cavalas-Medici, S.R. Day-Magdalena, Extremely low frequency, weak electric fields affect schedule-controlled behaviour of monkeys, *Nature* 261 (5557) (1976) 256–259.
- [5] L.K. Kaczmarek, W.R. Adey, The efflux of $^{45}\text{Ca}^{2+}$ and (3h)gamma-aminobutyric acid from cate cerebral cortex, *Brain Research* 63 (1973) 331–342.
- [6] L.K. Kaczmarek, W.R. Adey, Weak electric gradients change ionic and transmitter fluxes in cortex, *Brain Research* 66 (3) (1974) 537–540.
- [7] S.M. Bawin, L.K. Kaczmarek, W.R. Adey, Effects of modulated vhf fields on the central nervous system, *Annals of the New York Academy of Sciences* 247 (1975) 74–81.
- [8] S.M. Bawin, W.R. Adey, Sensitivity of calcium binding in cerebral tissue to weak environmental electric fields oscillating at low frequency, *Proceedings of the National Academy of Sciences of the United States of America* 73 (6) (1976) 1999–2003.
- [9] S.M. Bawin, W.R. Adey, I.M. Sabbot, Ionic factors in release of $^{45}\text{Ca}^{2+}$ from chicken cerebral tissue by electromagnetic fields, *Proceedings of the National Academy of Sciences of the United States of America* 75 (12) (1978) 6314–6318.
- [10] S.M. Bawin, A.R. Sheppard, W.R. Adey, Possible mechanism of weak electromagnetic field coupling in brain tissue, *Bioelectrochemistry & Bioenergetics* 5 (1978) 67–76.
- [11] A.R. Sheppard, S.M. Bawin, W.R. Adey, Models of long-range order in cerebral macromolecules: effects of sub-elf and of modulated vhf and uhf fields, *Radio Science* 14 (6S) (1979) 141–145.
- [12] C.F. Blackman, J.A. Elder, C.M. Weil, S.G. Benane, D.C. Eichinger, D.E. House, Induction of calcium ion efflux from brain tissue by radio-frequency radiation: effects of modulation-frequency and field strength, *Radio Science* 14 (6S) (1979) 93–98.
- [13] C.F. Blackman, S.G. Benane, J.A. Elder, D.E. House, J.A. Lampe, J.M. Faulk, Induction of calcium-ion efflux from brain tissue by radiofrequency radiation: effect of sample number and modulation frequency on the power-density window, *Bioelectromagnetics* 1 (1) (1980) 35–43.
- [14] C.F. Blackman, The biological influences of low-frequency sinusoidal electromagnetic signals alone and superimposed on rf carrier waves; in: A. Chiabrera, C. Nicolini, H.P. Schwan (Eds.), *Interaction between Electromagnetic Fields and Cells*, Erice, Italy, Plenum, New York, 1984, NATO ASI Series A97, pp. 521–535.
- [15] C.F. Blackman, S.G. Benane, D.J. Elliott, D.E. House, M.M. Pollock, Influence of electromagnetic fields on the efflux of calcium ions from brain tissue in vitro: a three-model analysis consistent with the frequency response up to 510 hz, *Bioelectromagnetics* 9 (3) (1988) 215–227.
- [16] C.F. Blackman, S.G. Benane, W.T. Joines, M.A. Hollis, D.E. House, Calcium-ion efflux from brain tissue: power-density versus internal field-intensity dependencies at 50-mhz rf radiation, *Bioelectromagnetics* 1 (3) (1980) 277–283.
- [17] C.F. Blackman, S.G. Benane, L.S. Kinney, W.T. Joines, D.E. House, Effects of elf fields on calcium-ion efflux from brain tissue in vitro, *Radiation Research* 92 (3) (1982) 510–520.
- [18] C.F. Blackman, S.G. Benane, J.R. Rabinowitz, D.E. House, W.T. Joines, A role for the magnetic field in the radiation-induced efflux of calcium ions from brain tissue in vitro, *Bioelectromagnetics* 6 (4) (1985) 327–337.
- [19] C.F. Blackman, D.E. House, S.G. Benane, W.T. Joines, R.J. Spiegel, Effect of ambient levels of power-line-frequency electric fields on a developing vertebrate, *Bioelectromagnetics* 9 (2) (1988) 129–140.
- [20] C.F. Blackman, W.T. Joines, J.A. Elder, Calcium-ion efflux in brain tissue by radiofrequency radiation, in: K.H. Illinger (Ed.), *Biological Effects of Nonionizing Radiation*, vol. 157, American Chemical Society, Washington, DC, 1981, pp. 299–314.
- [21] C.F. Blackman, S.G. Benane, D.E. House, D.J. Elliott, Importance of alignment between local dc magnetic field and an oscillating magnetic field in responses of brain tissue in vitro and in vivo, *Bioelectromagnetics* 11 (2) (1990) 159–167.
- [22] C.F. Blackman, L.S. Kinney, D.E. House, W.T. Joines, Multiple power-density windows and their possible origin, *Bioelectromagnetics* 10 (2) (1989) 115–128.
- [23] W.T. Joines, C.F. Blackman, Equalizing the electric field intensity within chick brain immersed in buffer solution at different carrier frequencies, *Bioelectromagnetics* 2 (4) (1981) 411–413.
- [24] W.T. Joines, C.F. Blackman, M.A. Hollis, Broadening of the rf power-density window for calcium-ion efflux from brain tissue, *IEEE Transactions on Bio-Medical Engineering* 28 (8) (1981) 568–573.
- [25] W.T. Joines, C.F. Blackman, R.J. Spiegel, Specific absorption rate in electrically coupled biological samples between metal plates, *Bioelectromagnetics* 7 (2) (1986) 163–176.
- [26] S.K. Dutta, K. Das, B. Ghosh, C.F. Blackman, Dose dependence of acetylcholinesterase activity in neuroblastoma cells exposed to modulated radio-frequency electromagnetic radiation, *Bioelectromagnetics* 13 (4) (1992) 317–322.
- [27] S.K. Dutta, B. Ghosh, C.F. Blackman, Radiofrequency radiation-induced calcium ion efflux enhancement from human and other neuroblastoma cells in culture, *Bioelectromagnetics* 10 (2) (1989) 197–202.
- [28] S.K. Dutta, A. Subramoniam, B. Ghosh, R. Parshad, Microwave radiation-induced calcium ion efflux from human neuroblastoma cells in culture, *Bioelectromagnetics* 5 (1) (1984) 71–78.
- [29] S.K. Dutta, M. Verma, C.F. Blackman, Frequency-dependent alterations in enolase activity in escherichia coli caused by exposure to electric and magnetic fields, *Bioelectromagnetics* 15 (5) (1994) 377–383.
- [30] E. Albert, C. Blackman, F. Slaby, Calcium dependent secretory protein release and calcium efflux during rf irradiation of rat pancreatic tissue slices, in: A.J. Berteaud, B. Servantie (Eds.), *Ondes Electromagnetiques et Biologie*, URSI International Symposium on Electromagnetic Waves and Biology, June 30–July 4, Jouy-en-Josas, France, Centre National de la Recherche Scientifique, 2 rue Henry Dunant, 94320 Thiais, France: A.J. Berteaud, 1980, pp. 325–329.
- [31] J.L. Schwartz, D.E. House, G.A. Mealing, Exposure of frog hearts to cw or amplitude-modulated vhf fields: selective efflux of calcium ions at 16 hz, *Bioelectromagnetics* 11 (4) (1990) 349–358.
- [32] J.L. Schwartz, G.A. Mealing, Calcium-ion movement and contractility in atrial strips of frog heart are not affected by low-frequency-modulated, 1 ghz electromagnetic radiation, *Bioelectromagnetics* 14 (6) (1993) 521–533.
- [33] C.F. Blackman, Can EMF exposure during development leave an imprint later in life? *Electromagnetic Biology and Medicine* 25 (4) (2006) 217–225.
- [34] C.J. Thompson, Y.S. Yang, V. Anderson, A.W. Wood, A cooperative model for Ca^{++} efflux windowing from cell membranes exposed to electromagnetic radiation, *Bioelectromagnetics* 21 (6) (2000) 455–464.
- [35] W.T. Joines, C.F. Blackman, Power density, field intensity, and carrier frequency determinants of rf-energy-induced calcium-ion efflux from brain tissue, *Bioelectromagnetics* 1 (3) (1980) 271–275.
- [36] A.R. Liboff, Cyclotron resonance in membrane transport, in: A. Chiabrera, C. Nicolini, H.P. Schwan (Eds.), *Interaction Between Electromagnetic Fields and Cells*, Erice, Italy, Plenum, New York, 1984, NATO ASI Series A97, pp. 281–296.
- [37] T.A. Litovitz, D. Krause, C.J. Montrose, J.M. Mullins, Temporally incoherent magnetic fields mitigate the response of biological systems to temporally coherent magnetic fields, *Bioelectromagnetics* 15 (5) (1994) 399–409.
- [38] J.M. Farrell, M. Barber, D. Krause, T.A. Litovitz, The superposition of a temporally incoherent magnetic field inhibits 60 hz-induced changes in the odc activity of developing chick embryos, *Bioelectromagnetics* 19 (1) (1998) 53–56.
- [39] T.A. Litovitz, C.J. Montrose, P. Doinov, K.M. Brown, M. Barber, Superimposing spatially coherent electromagnetic noise inhibits field-induced abnormalities in developing chick embryos, *Bioelectromagnetics* 15 (2) (1994) 105–113.

- [40] T.A. Litovitz, L.M. Penafiel, J.M. Farrel, D. Krause, R. Meister, J.M. Mullins, Bioeffects induced by exposure to microwaves are mitigated by superposition of elf noise, *Bioelectromagnetics* 18 (6) (1997) 422–430.
- [41] S. Milham, L.L. Morgan, A new electromagnetic exposure metric: high frequency voltage transients associated with increased cancer incidence in teachers in a California school, *American Journal of Industrial Medicine* 51 (8) (2008) 579–586.
- [42] M. Havas, Electromagnetic hypersensitivity: biological effects of dirty electricity with emphasis on diabetes and multiple sclerosis, *Electromagnetic Biology and Medicine* 25 (4) (2006) 259–268.
- [43] M. Havas, Dirty electricity elevates blood sugar among electrically sensitive diabetics and may explain brittle diabetes, *Electromagnetic Biology and Medicine* 27 (2) (2008) 135–146.
- [44] M. Havas, A. Olsiad, Power quality affects teacher wellbeing and student behavior in three Minnesota schools, *The Science of the Total Environment* 402 (2–3) (2008) 157–162.
- [45] C.F. Blackman, S.G. Benane, D.E. House, Frequency-dependent interference by magnetic fields of nerve growth factor-induced neurite outgrowth in pc-12 cells, *Bioelectromagnetics* 16 (6) (1995) 387–395.
- [46] C.F. Blackman, J.P. Blanchard, S.G. Benane, D.E. House, Experimental determination of hydrogen bandwidth for the ion parametric resonance model, *Bioelectromagnetics* 20 (1) (1999) 5–12.
- [47] M.A. Trillo, A. Ubeda, J.P. Blanchard, D.E. House, C.F. Blackman, Magnetic fields at resonant conditions for the hydrogen ion affect neurite outgrowth in pc-12 cells: a test of the ion parametric resonance model, *Bioelectromagnetics* 17 (1) (1996) 10–20.
- [48] L. Slesin, Highlights: Elf bioeffects studies at beams, *Microwave News* IV (7, Sept.) (1984) 2.
- [49] S.D. Smith, B.R. McLeod, A.R. Liboff, K. Cooksey, Calcium cyclotron resonance and diamon mobility, *Bioelectromagnetics* 8 (3) (1987) 215–227.
- [50] J.R. Thomas, J. Schrot, A.R. Liboff, Low-intensity magnetic fields alter operant behavior in rats, *Bioelectromagnetics* 7 (4) (1986) 349–357.
- [51] A.R. Liboff, B.R. McLeod, Kinetics of channelized membrane ions in magnetic fields, *Bioelectromagnetics* 9 (1) (1988) 39–51.
- [52] A.R. Liboff, W.C. Parkinson, Search for ion-cyclotron resonance in an na^{+} -transport system, *Bioelectromagnetics* 12 (2) (1991) 77–83.
- [53] C.F. Blackman, J.P. Blanchard, S.G. Benane, D.E. House, Effect of ac and dc magnetic field orientation on nerve cells, *Biochemical and Biophysical Research Communications* 220 (3) (1996) 807–811.
- [54] A.R. Liboff, Electric-field ion cyclotron resonance, *Bioelectromagnetics* 18 (1) (1997) 85–87.
- [55] V.V. Lednev, Possible mechanism for the influence of weak magnetic fields on biological systems, *Bioelectromagnetics* 12 (2) (1991) 71–75.
- [56] J.P. Blanchard, C.F. Blackman, Clarification and application of an ion parametric resonance model for magnetic field interactions with biological systems, *Bioelectromagnetics* 15 (3) (1994) 217–238.
- [57] M.N. Zhadin, E.E. Fesenko, Ionic cyclotron resonance in biomolecules, *Biomedical Science* 1 (3) (1990) 245–250.
- [58] E. Del Giudice, M. Fleischmann, G. Preparata, G. Talpo, On the “Unreasonable” Effects of elf magnetic fields upon a system of ions, *Bioelectromagnetics* 23 (7) (2002) 522–530.
- [59] V.N. Binhi, Stochastic dynamics of magnetosomes and a mechanism of biological orientation in the geomagnetic field, *Bioelectromagnetics* 27 (1) (2006) 58–63.
- [60] V.N. Binhi, A few remarks on ‘combined action of dc and ac magnetic fields on ion motion in a macromolecule’, *Bioelectromagnetics* 28 (5) (2007) 409–412, discussion 412–404.
- [61] V.N. Binhi, A.B. Rubin, Magnetobiology: the kt paradox and possible solutions, *Electromagnetic Biology and Medicine* 26 (1) (2007) 45–62.
- [62] V.N. Binhi, A.V. Savin, Molecular gyroscopes and biological effects of weak extremely low-frequency magnetic fields, *Physical Review* 65 (5 Pt 1) (2002) 051912.
- [63] A.Y. Matronchik, I.Y. Belyaev, Mechanism for combined action of microwaves and static magnetic field: slow non uniform rotation of charged nucleoid, *Electromagnetic Biology and Medicine* 27 (4) (2008) 340–354.
- [64] M.N. Zhadin, Combined action of static and alternating magnetic fields on ion motion in a macromolecule: theoretical aspects, *Bioelectromagnetics* 19 (5) (1998) 279–292.
- [65] M.N. Zhadin, V.V. Novikov, F.S. Barnes, N.F. Pergola, Combined action of static and alternating magnetic fields on ionic current in aqueous glutamic acid solution, *Bioelectromagnetics* 19 (1) (1998) 41–45.
- [66] V.V. Vorobyov, E.A. Sosunov, N.I. Kukushkin, V.V. Lednev, Weak combined magnetic field affects basic and morphine-induced rat's eeg, *Brain Research* 781 (1–2) (1998) 182–187.
- [67] C.L. Baureus Koch, M. Sommarin, B.R. Persson, L.G. Salford, J.L. Eberhardt, Interaction between weak low frequency magnetic fields and cell membranes, *Bioelectromagnetics* 24 (6) (2003) 395–402.
- [68] R. Sarimov, E. Markova, F. Johansson, D. Janssen, I. Belyaev, Exposure to elf magnetic field tuned to zn inhibits growth of cancer cells, *Bioelectromagnetics* 26 (8) (2005) 631–638.
- [69] F.S. Prato, M. Kavaliers, J.J. Carson, Behavioural evidence that magnetic field effects in the land snail, *Cepaea nemoralis*, might not depend on magnetite or induced electric currents, *Bioelectromagnetics* 17 (2) (1996) 123–130.
- [70] F.S. Prato, M. Kavaliers, A.P. Cullen, A.W. Thomas, Light-dependent and -independent behavioral effects of extremely low frequency magnetic fields in a land snail are consistent with a parametric resonance mechanism, *Bioelectromagnetics* 18 (3) (1997) 284–291.
- [71] N. Comisso, E. Del Giudice, A. De Nino, M. Fleischmann, L. Giuliani, G. Mengoli, F. Merlo, G. Talpo, Dynamics of the ion cyclotron resonance effect on amino acids adsorbed at the interfaces, *Bioelectromagnetics* 27 (1) (2006) 16–25.
- [72] V.V. Novikov, I.M. Sheiman, E.E. Fesenko, Effect of weak static and low-frequency alternating magnetic fields on the fission and regeneration of the planarian *dugesia (girardia) tigrina*, *Bioelectromagnetics* 29 (5) (2008) 387–393.
- [73] C.F. Blackman, B. Most, A scheme for incorporating dc magnetic fields into epidemiological studies of EMP exposure, *Bioelectromagnetics* 14 (5) (1993) 413–431.
- [74] M.T. Marron, E.M. Goodman, P.T. Sharpe, B. Greenbaum, Low frequency electric and magnetic fields have different effects on the cell surface, *FEBS Letters* 230 (1–2) (1988) 13–16.
- [75] R.P. Liburdy, Calcium signaling in lymphocytes and elf fields. Evidence for an electric field metric and a site of interaction involving the calcium ion channel, *FEBS Letters* 301 (1) (1992) 53–59.
- [76] J.J. Greene, W.J. Skowronski, J.M. Mullins, R.M. Nardone, M. Penafiel, R. Meister, Delineation of electric and magnetic field effects of extremely low frequency electromagnetic radiation on transcription, *Biochemical and Biophysical Research Communications* 174 (2) (1991) 742–749.
- [77] C.F. Blackman, S.G. Benane, D.E. House, Evidence for direct effect of magnetic fields on neurite outgrowth, *FASEB J* 7 (9) (1993) 801–806.
- [78] C.F. Blackman, S.G. Benane, D.E. House, M.M. Pollock, Action of 50 hz magnetic fields on neurite outgrowth in pheochromocytoma cells, *Bioelectromagnetics* 14 (3) (1993) 273–286.
- [79] A. Chiabrera, M. Hinsenkamp, A.A. Pilla, J. Ryaby, D. Ponta, A. Belmont, F. Beltrame, M. Gratarola, C. Nicolini, Cytofluorometry of electromagnetically controlled cell dedifferentiation, *The Journal of Histochemistry and Cytochemistry* 27 (1) (1979) 375–381.
- [80] M.R. Scarfi, F. Prisco, M.B. Lioi, O. Zeni, M. Della Noce, R. Di Pietro, C. Franceschi, D. Iafusco, M. Motta, B. F., Cytogenetic effects induced by extremely low frequency pulsed magnetic fields in lymphocytes from Turner's syndrome subjects, *Bioelectrochemistry & Bioenergetics* 43 (1997) 221–226.
- [81] M.R. Scarfi, M.B. Lioi, O. Zeni, G. Franceschetti, C. Franceschi, F. Bersani, Lack of chromosomal aberration and micronucleus induction

- in human lymphocytes exposed to pulsed magnetic fields, *Mutation Research* 306 (2) (1994) 129–133.
- [82] K. Takahashi, I. Kaneko, M. Date, E. Fukada, Effect of pulsing electromagnetic fields on DNA synthesis in mammalian cells in culture, *Experientia* 42 (2) (1986) 185–186.
- [83] A. Uboda, J. Leal, M.A. Trillo, M.A. Jimenez, J.M. Delgado, Pulse shape of magnetic fields influences chick embryogenesis, *Journal of Anatomy* 137 (Pt 3) (1983) 513–536.
- [84] A. Chiabrera, B. Bianco, E. Moggia, J.J. Kaufman, Zeeman-stark modeling of the rf EMF interaction with ligand binding, *Bioelectromagnetics* 21 (4) (2000) 312–324.
- [85] A.A. Marino, R.M. Wolcott, R. Chervenak, F. Jour'd'Heuil, E. Nilsen, C. Frilot 2nd, Nonlinear response of the immune system to power-frequency magnetic fields, *American Journal of Physiology* 279 (3) (2000) R761–768.
- [86] A.A. Marino, R.M. Wolcott, R. Chervenak, F. Jour'd'Heuil, E. Nilsen, C. Frilot 2nd, Nonlinear determinism in the immune system. In vivo influence of electromagnetic fields on different functions of murine lymphocyte subpopulations, *Immunological Investigations* 30 (4) (2001) 313–334.
- [87] A.A. Marino, R.M. Wolcott, R. Chervenak, F. Jour'd'Heuil, E. Nilsen, C. Frilot 2nd, Nonlinear dynamical law governs magnetic field induced changes in lymphoid phenotype, *Bioelectromagnetics* 22 (8) (2001) 529–546.
- [88] A.A. Marino, R.M. Wolcott, R. Chervenak, F. Jour'd'Heuil, E. Nilsen, C. Frilot 2nd, S.B. Prwet, Coincident nonlinear changes in the endocrine and immune systems due to low-frequency magnetic fields, *Neuroimmunomodulation* 9 (2) (2001) 65–77.
- [89] A.A. Marino, E. Nilsen, A.L. Chesson Jr., C. Frilot, Effect of low-frequency magnetic fields on brain electrical activity in human subjects, *Clinical Neurophysiology* 115 (5) (2004) 1193–1201.
- [90] A.A. Marino, E. Nilsen, C. Frilot, Localization of electroreceptive function in rabbits, *Physiology & Behavior* 79 (4–5) (2003) 803–810.
- [91] A.A. Marino, E. Nilsen, C. Frilot, Nonlinear changes in brain electrical activity due to cell phone radiation, *Bioelectromagnetics* 24 (5) (2003) 339–346.
- [92] A.A. Marino, E. Nilsen, C. Frilot, Consistent magnetic-field induced dynamical changes in rabbit brain activity detected by recurrence quantification analysis, *Brain Research* 964 (2) (2003) 317–326.
- [93] O.V. Kolomytkin, S. Dunn, F.X. Hart, C. Frilot 2nd, D. Kolomytkin, A.A. Marino, Glycoproteins bound to ion channels mediate detection of electric fields: a proposed mechanism and supporting evidence, *Bioelectromagnetics* 28 (3) (2007) 379–385.
- [94] S. Carrubba, C. Frilot, A. Chesson, A.A. Marino, Detection of nonlinear event-related potentials, *Journal of Neuroscience Methods* 157 (1) (2006) 39–47.
- [95] S. Carrubba, C. Frilot, A.L. Chesson, A.A. Marino, Nonlinear eeg activation evoked by low-strength low-frequency magnetic fields, *Neuroscience Letters* 417 (2) (2007) 212–216.
- [96] S. Carrubba, C. Frilot 2nd, A.L. Chesson Jr., A.A. Marino, Evidence of a nonlinear human magnetic sense, *Neuroscience* 144 (1) (2007) 356–367.
- [97] S. Carrubba, C. Frilot, A.L. Chesson Jr., A.A. Marino, Method for detection of changes in the eeg induced by the presence of sensory stimuli, *Journal of Neuroscience Methods* 173 (1) (2008) 41–46.
- [98] S. Carrubba, C. Frilot, A.L. Chesson Jr., C.L. Webber Jr., J.P. Zbilut, A.A. Marino, Magnetosensory evoked potentials: consistent nonlinear phenomena, *Neuroscience Research* 60 (1) (2008) 95–105.
- [99] M.N. Zhadin, O.N. Deryugina, T.M. Pisachenko, Influence of combined dc and ac magnetic fields on rat behavior, *Bioelectromagnetics* 20 (6) (1999) 378–386.
- [100] P. Slovic, Trust, emotion, sex, politics, and science: surveying the risk-assessment battlefield, *Risk Analysis* 19 (4) (1999) 689–701.
- [101] M. Havas, Biological effects of non-ionizing electromagnetic energy: a critical review of the reports by the US national research council and the us national institute of environmental health sciences as they relate to the broad realm of EMF bioeffects, *Environmental Reviews* 8 (2000) 173–253.
- [102] National Research Council (U.S.), Committee on the Possible Effects of Electromagnetic Fields on Biologic Systems, National Academy Press, Washington, DC, 1997, 356 pp.
- [103] National Institute of Environmental Health Science Working Group Report, Assessment of health effects from exposure to power-line frequency electric and magnetic fields, 1998, NIH Pub 98-3981, 508 pp.
- [104] National Institute of Environmental Health Science, Report on health effects from exposure to power-line frequency electric and magnetic fields, NIH Pub No 99-4493, 1999, 67 pp.
- [105] M. Fedrowitz, K. Kamino, W. Loscher, Significant differences in the effects of magnetic field exposure on 7, 12-dimethylbenz(a)anthracene-induced mammary carcinogenesis in two substrains of sprague-dawley rats, *Cancer Research* 64 (1) (2004) 243–251.
- [106] M. Fedrowitz, W. Loscher, Power frequency magnetic fields increase cell proliferation in the mammary gland of female fischer 344 rats but not various other rat strains or substrains, *Oncology* 69 (6) (2005) 486–498.
- [107] Y. Yang, X. Jin, C. Yan, Y. Tian, J. Tang, X. Shen, Case-only study of interactions between DNA repair genes (hmlh1, apex1, mgmt, xrc1 and xpd) and low-frequency electromagnetic fields in childhood acute leukemia, *Leukemia & Lymphoma* 49 (12) (2008) 2344–2350.
- [108] S.M. Bawin, R.J. Gavalas-Medici, W.R. Adey, Effects of modulated very high frequency fields on specific brain rhythms in cats, *Brain Research* 58 (2) (1973) 365–384.
- [109] A.W. Preece, G. Iwi, A. Davies-Smith, K. Wesnes, S. Butler, E. Lim, A. Vary, Effect of a 915-mhz simulated mobile phone signal on cognitive function in man, *International Journal of Radiation Biology* 75 (4) (1999) 447–456.
- [110] A.W. Preece, K.A. Wesnes, G.R. Iwi, The effect of a 50 hz magnetic field on cognitive function in humans, *International Journal of Radiation Biology* 74 (4) (1998) 463–470.
- [111] M. Koivisto, C.M. Krause, A. Revonsuo, M. Laine, H. Hamalainen, The effects of electromagnetic field emitted by gsm phones on working memory, *Neuroreport* 11 (8) (2000) 1641–1643.
- [112] M. Koivisto, A. Revonsuo, C. Krause, C. Haara, L. Sillanmaki, M. Laine, H. Hamalainen, Effects of 902 mhz electromagnetic field emitted by cellular telephones on response times in humans, *Neuroreport* 11 (2) (2000) 413–415.
- [113] C.M. Krause, L. Sillanmaki, M. Koivisto, A. Haggqvist, C. Saarela, A. Revonsuo, M. Laine, H. Hamalainen, Effects of electromagnetic field emitted by cellular phones on the eeg during a memory task, *Neuroreport* 11 (4) (2000) 761–764.
- [114] C.M. Krause, L. Sillanmaki, M. Koivisto, A. Haggqvist, C. Saarela, A. Revonsuo, M. Laine, H. Hamalainen, Effects of electromagnetic fields emitted by cellular phones on the electroencephalogram during a visual working memory task, *International Journal of Radiation Biology* 76 (12) (2000) 1659–1667.
- [115] G. Freude, P. Ullsperger, S. Eggert, I. Ruppe, Effects of microwaves emitted by cellular phones on human slow brain potentials, *Bioelectromagnetics* 19 (6) (1998) 384–387.
- [116] G. Freude, P. Ullsperger, S. Eggert, I. Ruppe, Microwaves emitted by cellular telephones affect human slow brain potentials, *European Journal of Applied Physiology* 81 (1–2) (2000) 18–27.
- [117] A.W. Preece, S. Goodfellow, M.G. Wright, S.R. Butler, E.J. Dunn, Y. Johnson, T.C. Manktelow, K. Wesnes, Effect of 902 mhz mobile phone transmission on cognitive function in children, *Bioelectromagnetics* (Suppl. 7) (2005) S138–143.
- [118] C.M. Krause, C.H. Björnberg, M. Pesonen, A. Hulten, T. Liesivuori, M. Koivisto, A. Revonsuo, M. Laine, H. Hamalainen, Mobile phone effects on children's event-related oscillatory eeg during an auditory memory task, *International Journal of Radiation Biology* 82 (6) (2006) 443–450.

- [119] C.M. Krause, M. Pesonen, C. Haarala Bjornberg, H. Hamalainen. Effects of pulsed and continuous wave 902 mhz mobile phone exposure on brain oscillatory activity during cognitive processing, *Bioelectromagnetics* 28 (4) (2007) 296–308.
- [120] C. Haarala, L. Bjornberg, M. Ek, M. Laine, A. Revonsuo, M. Koivisto, H. Hamalainen. Effect of a 902 mhz electromagnetic field emitted by mobile phones on human cognitive function: a replication study, *Bioelectromagnetics* 24 (4) (2003) 283–288.
- [121] C. Haarala, M. Ek, L. Bjornberg, M. Laine, A. Revonsuo, M. Koivisto, H. Hamalainen. 902 mhz mobile phone does not affect short term memory in humans, *Bioelectromagnetics* 25 (6) (2004) 452–456.
- [122] C. Haarala, M. Bergman, M. Laine, A. Revonsuo, M. Koivisto, H. Hamalainen. Electromagnetic field emitted by 902 mhz mobile phones shows no effects on children's cognitive function, *Bioelectromagnetics Suppl. 7* (2005) S144–150.
- [123] C. Haarala, F. Takio, T. Rintee, M. Laine, M. Koivisto, A. Revonsuo, H. Hamalainen. Pulsed and continuous wave mobile phone exposure over left versus right hemisphere: effects on human cognitive function, *Bioelectromagnetics* 28 (4) (2007) 289–295.
- [124] A.A. Borbely, R. Huber, T. Graf, B. Fuchs, E. Gallmann, P. Achermann. Pulsed high-frequency electromagnetic field affects human sleep and sleep electroencephalogram, *Neuroscience Letters* 275 (3) (1999) 207–210.
- [125] R. Huber, J. Schuderer, T. Graf, K. Jutz, A.A. Borbely, N. Kuster, P. Achermann. Radio frequency electromagnetic field exposure in humans: estimation of sar distribution in the brain, effects on sleep and heart rate, *Bioelectromagnetics* 24 (4) (2003) 262–276.
- [126] S.P. Loughran, A.W. Wood, J.M. Barton, R.J. Croft, B. Thompson, C. Stough. The effect of electromagnetic fields emitted by mobile phones on human sleep, *Neuroreport* 16 (17) (2005) 1973–1976.
- [127] H. D'Costa, G. Trueman, L. Tang, U. Abdel-rahman, W. Abdel-rahman, K. Ong, I. Cosic. Human brain wave activity during exposure to radiofrequency field emissions from mobile phones, *Australasian Physical & Engineering Sciences in Medicine* 26 (2003) 162–167.
- [128] G. Fritzer, R. Goder, L. Friege, J. Wachter, V. Hansen, D. Hinze-Selch, J.B. Aldenhoff. Effects of short- and long-term pulsed radiofrequency electromagnetic fields on night sleep and cognitive functions in healthy subjects, *Bioelectromagnetics* 28 (4) (2007) 316–325.
- [129] W.S. Pritchard, D.W. Duke. Measuring chaos in the brain: a tutorial review of nonlinear dynamical eeg analysis, *The International Journal of Neuroscience* 67 (1–4) (1992) 31–80.
- [130] National Academy of Science. Identification of Research Needs Relating to Potential Biological or Adverse Health Effects of Wireless Communication, Washington, DC, 2009. <http://www.nap.edu/catalog/12036.html>.
- [131] P.H. Abelson. Effects of electric and magnetic fields, *Science* (New York, N.Y.) 245 (4915) (1989) 241.

Public health implications of wireless technologies

Cindy Sage^{a,*}, David O. Carpenter^b

^a Sage Associates, 1396 Danielson Road, Santa Barbara, CA 93108, USA

^b Institute for Health and the Environment, University at Albany, Rensselaer, NY, USA

Received 18 January 2008; accepted 30 January 2009

Abstract

Global exposures to emerging wireless technologies from applications including mobile phones, cordless phones, DECT phones, WI-FI, WLAN, WiMAX, wireless internet, baby monitors, and others may present serious public health consequences. Evidence supporting a public health risk is documented in the BioInitiative Report. New, biologically based public exposure standards for chronic exposure to low-intensity exposures are warranted. Existing safety standards are obsolete because they are based solely on thermal effects from acute exposures. The rapidly expanding development of new wireless technologies and the long latency for the development of such serious diseases as brain cancers means that failure to take immediate action to reduce risks may result in an epidemic of potentially fatal diseases in the future. Regardless of whether or not the associations are causal, the strengths of the associations are sufficiently strong that in the opinion of the authors, taking action to reduce exposures is imperative, especially for the fetus and children. Such action is fully compatible with the precautionary principle, as enunciated by the Rio Declaration, the European Constitution Principle on Health (Section 3.1) and the European Union Treaties Article 174. © 2009 Elsevier Ireland Ltd. All rights reserved.

Keywords: Wireless technology; Brain cancer; Radiofrequency; Cell phones; Wireless antenna facilities; Children's health

1. Introduction and background

Exposure to electromagnetic fields (EMF) has been linked to a variety of adverse health outcomes that may have significant public health consequences [1–13]. The most serious health endpoints that have been reported to be associated with extremely low frequency (ELF) and/or RF include childhood and adult leukemia, childhood and adult brain tumors, and increased risk of the neurodegenerative diseases, Alzheimer's and amyotrophic lateral sclerosis (ALS). In addition, there are reports of increased risk of breast cancer in both men and women, genotoxic effects (DNA damage and micronucleation), pathological leakage of the blood–brain barrier, altered immune function including increased allergic and inflammatory responses, miscarriage and some cardiovascular effects [1–13]. Insomnia (sleep disruption) is reported in studies of people living in very low-intensity RF environments with WI-FI and cell tower-level exposures [85–93]. Short-term effects on cognition, memory and learning, behavior, reaction time, attention and concentration, and altered

brainwave activity (altered EEG) are also reported in the scientific literature [94–107]. Biophysical mechanisms that may account for such effects can be found in various articles and reviews [136–144].

The public health implications of emerging wireless technologies are enormous because there has been a very rapid global deployment of both old and new forms in the last 15 years. In the United States, the deployment of wireless infrastructure has accelerated greatly in the last few years with 220,500 cell sites in 2008 [14–16]. Eighty-four percent of the population of the US own cell phones [16]. Annualized wireless revenues in 2008 will reach \$144 billion and US spending on wireless communications will reach \$212 billion by 2008. Based on the current 15% annual growth rate enjoyed by the wireless industry, in the next 5 years wireless will become a larger sector of the US economy than both the agriculture and automobile sectors. The annualized use of cell phones in the US is estimated to be 2.23 trillion minutes in 2008 [16]. There are 2.2 billion users of cell phones worldwide in 2008 [17] and many million more users of cordless phones.

Over 75 billion text messages were sent in the United States, compared with 7.2 billion in June 2005, according to

* Corresponding author. Tel.: +1 805 969 0557; fax: +1 805 969 5003.
E-mail address: sage@silcom.com (C. Sage).

Behavior
to Brain
effect

economy

CTIA, the Wireless Association, the leading industry trade group [16]. The consumer research company Nielsen Mobile, which tracked 50,000 individual customer accounts in the second quarter of this year, found that Americans each sent or received 357 text messages a month then, compared with 204 phone calls. That was the second consecutive quarter in which mobile texting significantly surpassed the number of voice calls [17].

The Electronics Industries Alliance (EIA) represents 80% of the \$550 billion US electronics industry "that provides two million jobs for American workers." Its members include companies from the consumer electronics and telecommunications industries, among others [17].

There is intense industry competition for market share. Telecom taxes form an immense revenue generator for the government sector. Sale of the airwaves (auctions selling off wireless bandwidth) is a multi-million dollar industry for governments, and multi-billion dollar global advertising budgets are common. Lobbying dollars from the telecom-related industries are estimated to be \$300 million annually. The media is nearly silent on health issues, perhaps in part because of global advertising revenues that compromise journalistic independence and discourage balanced coverage of health, equity and economic issues.

2. Evidence supporting a public health risk

Even if there is only a small risk to health from chronic use of and exposure to wireless technologies, there is the potential for a profound public health impact. RF radiation now saturates the airwaves, resulting in exposure to both users and non-users. The effects are both short-term (sleep disruption, hormone disruption, impairment of cognitive function, concentration, attention, behavior, and well-being) and they are almost certainly long-term (generational impacts on health secondary to DNA damage, physiological stress, altered immune function, electrosensitivity, miscarriage risks, effects on sperm quality and motility leading to infertility, increased rates of cancer, and neurological diseases including Alzheimer's disease and ALS—at least for ELF exposures). (Chapters 5–12 of the BioInitiative Report [1] and papers in this Supplement.)

There is credible scientific evidence that RF exposures cause changes in cell membrane function, metabolism and cellular signal communication, as well as activation of proto-oncogenes and triggering of the production of stress proteins at exposure levels below current regulatory limits. There is also generation of reactive oxygen species, which cause DNA damage, chromosomal aberrations and nerve cell death. A number of different effects on the central nervous system have also been documented, including activation of the endogenous opioid systems, changes in brain function including memory loss, slowed learning, motor dysfunction and performance impairment in children, and increased frequency of headaches, fatigue and sleep disorders. Melatonin secretion

is reduced, resulting in altered circadian rhythms and disruption of several physiological functions. (Chapters 5–12 of the BioInitiative Report [1] and papers in this Supplement.)

These effects can reasonably be presumed to result in adverse health effects and disease with chronic and uncontrolled exposures, and children may be particularly vulnerable [1,19]. The young are also largely unable to remove themselves from such environments. Second-hand non-ionizing radiation, like second-hand smoke may be considered of public health concern based on the evidence at hand.

2.1. Malignant brain tumors

At present, the most persuasive evidence for cancer resulting from RF exposure is that there is a significantly increased risk of malignant glioma in individuals that have used a mobile phone for 10 or more years, with the risk being elevated only on the side of the head on which the phone is used regularly (ipsilateral use) [1,3,4,6–8,18]. While the risk for adults after 10 or more years of use is reported to be more than doubled, there is some evidence beginning to appear that indicates that the risk is greater if the individual begins to use a mobile phone at younger ages. Hardell et al. [18] reported higher odds ratios in the 20–29-year-old group than other age ranges after more than 5 years of use of either analog or cordless phones. Recently in a London symposium Hardell reported that after even just 1 or more years of use there is a 5.2-fold elevated risk in children who begin use of mobile phones before the age of 20 years, whereas for all ages the odds ratio was 1.4. Studies from Israel have found that the risk of parotid gland tumors (a salivary gland in the cheek) is increased with heavy cell phone use [7]. The risk of acoustic neuroma (a benign but space-occupying tumor on the auditory nerve) is also significantly increased on the ipsilateral side of the head after 10 or more years of mobile phone use [1,3]. This relationship has also been documented in some of the published reports of the WHO Interphone Study, a decade-long 13-country international assessment of cell phone risks and cancer [6,8].

Kundi reports that "(E)pidemiological evidence compiled in the last 10 years starts to indicate an increased risk, in particular for brain tumors (glioma, meningioma, acoustic neuroma), from mobile phone use. Considering biases that may have been operating in most studies the risk estimates are rather too low, although recall bias could have increased risk estimates. The net result, when considering the different errors and their impact is still an elevated risk" [19].

The latency for (most brain tumors is 20 years) or more when related to other environmental agents, for example, to X-ray exposure. Yet, for cell phone use the increased risks are occurring much sooner than twenty years, as early as 10 years for brain tumors in adults and with even shorter latencies in children. This suggests that we may currently be significantly underestimating the impact of current levels of

use of RF technology, since we do not know how long the average latency period really is. If it is 20 years, then the risk rate will likely be much higher than an overall doubling of risk for cell phone users if the peak comes later than 10 years. It may also signal very troubling risks for those who start using cell phones, and perhaps all wireless devices, in early childhood. We may not have proof of effect for decades until many hundreds of thousands of new cases of malignant gliomas are set in motion by long-term cell phone use.

The preliminary evidence that mobile phone use at younger ages may lead to greater risk than for older persons is of particular concern. There is a large body of evidence that childhood exposure to environmental agents poses greater risk to health than comparable exposure during adulthood [20,21]. There is reason to expect that children would be more susceptible to the effects of EMP exposure since they are growing, their rate of cellular activity and division is more rapid, and they may be more at risk for DNA damage and subsequent cancers. Growth and development of the central nervous system is still occurring well into the teenage years so that neurological changes may be of great importance to normal development, cognition, learning, and behavior.

A greater vulnerability of children to developing brain cancer from mobile phone use may be the consequence of a combination of patterns of use, stage of development and physical characteristics related to exposure. In addition to the fact that the brain continues to develop through the teen years, many young children and teenagers now spend very large periods of time using mobile phones. The brain is the main target organ of cell phones and cordless phones, with highest exposure to the same side as the phone is used. Further, due to anatomical reasons, the brain of a child is more exposed to RF radiation than the brain of an adult [22,23]. This is caused by the smaller brain size, a thinner pinna of the ear, thinner skin and thinner skull bone permitting deeper penetration into the child's brain. A recent French study showed that children absorb twice the RF from cell phone use as do adults [24].

In addition to concerns about cancer, there is evidence for short-term effects of RF exposure on cognition, memory and learning, behavior, reaction time, attention and concentration, altered brainwave activity (altered EEG) [95–108], and all of these effects argue for extreme caution with regard to exposure of children. The development of children into adults is characterized by faster cell division during growth, the long period needed to fully develop and mature all organ systems, and the need for properly synchronized neural development until early adulthood. Chronic, cumulative RF exposures may alter the normal growth and development of children and adversely affect their development and capacity for normal learning, nervous system development, behavior and judgment [1,97,102].

Prenatal exposure to EMP has been identified as a possible risk factor for childhood leukemia (1). Maternal use of cell phones has been reported to adversely affect fetal brain development, resulting in behavioral problems in those children by

the time they reach school age [25]. Their exposure is involuntary in all cases. Children are largely unable to remove themselves from exposures to harmful substances in their environments.

2.2. Plausible biological mechanisms for a relationship between RF exposure and cancer

2.2.1. DNA damage and oxidative stress

Damage to DNA from ELF and from RF cell phone frequencies at very low intensities (far below FCC and ICNIRP safety limits) has been demonstrated in many studies [1,2,26–35]. Both single- and double-strand DNA damage have been reported by various researchers in different laboratories. This is damage to the human genome, and can lead to mutations which can be inherited, or which can cause cancer, or both.

Non-ionizing radiation is assumed to be of too low energy to cause direct DNA damage. However both ELF and RF radiation induce reactive oxygen species, free radicals that react with cellular molecules including DNA. Free-radical production and/or the failure to repair DNA damage (secondary to damage to the enzymes that repair damage) created by such exposures can lead to mutations. Whether it is greater free-radical production, reduction in anti-oxidant protection or reduced repair capacity, the result will be altered DNA, increased risk of cancer, impaired or delayed healing, and premature aging [36–54]. Exposures have also been linked to decreased melatonin production, which is a plausible biological mechanism for decreased cancer surveillance in the body, and increased cancer risk [34,39,44,46,47,49,50,54]. An increased risk of cancers and a decrease in survival has been reported in numerous studies of ELF and RF [55–69].

2.2.2. Stress proteins (heat shock proteins or HSP)

Another well-documented effect of exposure to low-intensity ELF and RF is the creation of stress proteins (heat shock proteins) that signal a cell is being placed under physiological stress [70–80]. The HSP response is generally associated with heat shock, exposure to toxic chemicals and heavy metals, and other environmental insults. HSP is a signal of cells in distress. Plants, animals and bacteria all produce stress proteins to survive environmental stressors like high temperatures, lack of oxygen, heavy metal poisoning, and oxidative stress.

We can now add ELF and RF exposures to this list of environmental stressors that cause a physiological stress response. Very low-level ELF and RF exposures can cause cells to produce stress proteins, meaning that the cell recognizes ELF and RF exposures as harmful. This is another important way in which scientists have documented that ELF and RF exposures can be harmful, and it happens at levels far below the existing public safety standards. An additional concern is that if the stress goes on too long, the protective effect is diminished. The reduced response with prolonged exposure means the cell is less protected against

Children
7 am

damage, and this is why prolonged or chronic exposures may be harmful, even at very low intensities.

2.2.3. RF-induced gene expression changes

Many environment agents cause diseases, including cancer, not by direct damage to DNA but rather by up- or down-regulation of genes that regulate cell growth and function. Usually there are many genes whose expression is changed, and it is difficult to determine the exact changes responsible for the disease. Both ELF and RF exposures have been shown to result in altered gene expression. Olivares-Banuelos et al. [81] found that ELF exposure of chromaffin cells resulted in changed expression of 53 transcripts. Zhao et al. [82] investigated the gene expression profile of rat neurons exposed to 1800 MHz RF fields (2 W/kg) and found 24 up-regulated genes and 10 down-regulated genes after a 24-h exposure. The altered genes were involved in multiple cellular functions including cytoskeleton, signal transduction pathways and metabolism. Kariene et al. [83] exposed human skin to mobile phone radiation, and found by punch biopsy that 8 proteins were significantly altered in expression, consistent with gene induction. Several other studies have found altered gene expression following RF exposure, although none have been found that explain specific disease states [84].

DNA activation at very low ELF and RF levels, as in the stress response, and DNA damage (strand breaks and micronuclei) at higher levels, are molecular precursors to changes that are believed to lead to cancer. These, along with gene induction, provide plausible biological mechanisms linking exposure to cancer.

The biochemical pathways that are activated are the same for ELF and for RF exposures, and are non-thermal (do not require heating or induced electrical currents). This is true for the stress response. DNA damage, generation of reactive oxygen species as well as gene induction. Thus it is not surprising that the major cancers resulting from exposure to ELF and RF are the same, namely leukemia and brain cancer. The safety standards for both ELF and RF, based on protection from heating, are irrelevant and not protective. ELF exposure levels of only 5–10 mG have been shown to activate the stress response genes (<http://www.bioinitiative.org>, Sections 1 and 7 [1]).

3. Sleep, cognitive function and performance

The relationship of good sleep to cognition, performance and healing is well recognized. Sleep is a profoundly important factor in proper healing, anti-inflammatory benefits, reduction in physical symptoms of such as tendonitis, over-use syndrome, fatigue-induced lethargy, cognition and learning. Incomplete or slowed physiological recovery is common when sleep is impaired. Circadian rhythms that normalize stress hormone production (cortisol, for example) depend on synchronized sleep patterns.

People who are chronically exposed to low-level wireless antenna emissions report symptoms such as problems in sleeping (insomnia), as well as other symptoms that include fatigue, headache, dizziness, grogginess, lack of concentration, memory problems, ringing in the ears (tinnitus), problems with balance and orientation, and difficulty in multi-tasking [85–93,99]. In children, exposures to cell phone radiation have resulted in changes in brain oscillatory activity during some memory tasks [97,102]. Cognitive impairment, loss of mental concentration, distraction, speeded mental function but lowered accuracy, impaired judgment, delayed reaction time, spatial disorientation, dizziness, fatigue, headache, slower motor skills and reduced learning ability in children and adults have all been reported [85–108].

These symptoms are more common among “electrosensitive” individuals, although electrosensitivity has not been documented in double-blind tests of individual identifying themselves as being electrosensitive as compared to controls [109,110]. However people traveling to laboratories for testing are pre-exposed to a multitude of RF and ELF exposures, so they may already be symptomatic prior to actual testing. There is also evidence that RF exposures testing behavioral changes show delayed results; effects are observed after termination of RF exposure. This suggests a persistent change in the nervous system that may be evident only after time has passed, so is not observed during a short testing period.

3.1. Plausible biological mechanisms for neurobehavioral effects

3.1.1. The melatonin hypothesis

While there remains controversy as to the degree that RF and ELF fields alter neurobehavioral function, emerging evidence provides a plausible mechanism for both effects on sleep and cognition. Sleep is controlled by the central circadian oscillator in the suprachiasmatic nucleus, located in the hypothalamus. The activity of this central circadian oscillator is, in turn, controlled by the hormone, melatonin, which is released from the pineal gland [111]. There is considerable evidence that ELF exposure reduces the release of melatonin from the pineal gland—see Section 12 of the Bioinitiative Report [1]. There has been less study of the effects of RF exposure on melatonin release, but investigations have demonstrated a reduced excretion of the urinary metabolite of melatonin among persons using a mobile phone for more than 25 min per day [112]. In a study of women living near to radio and television transmitters, Clark et al. [113] found no effect on urinary melatonin metabolite excretion among pre-menopausal women, but a strong effect in post-menopausal women.

The “melatonin hypothesis” also provides a possible basis for other reported effects of EMFs. Melatonin has important actions on learning and memory, and inhibits electrophysiological components of learning in some but not all areas of the brain [114,115]. Melatonin has properties as a free-radical scavenger and anti-oxidant [116], and consequently,

24h
time
24h

non-thermal
*
*

RF

all
Dopamine
Blind
Datura

menopausal
♀

*Melatonin
7/2007/04
7/Alzheimer*

a reduction in melatonin levels would be expected to increase susceptibility to cancer and cellular damage. Melatonin could also be the key to understanding the relationship between EMF exposure and Alzheimer's disease. Noonan et al. [117] reported that there was an inverse relationship between excretion of the melatonin metabolite and the 1–42 amino acid form of amyloid beta in electric utility workers. This form of amyloid beta has been found to be elevated in Alzheimer's patients.

3.1.2. Blood–brain barrier alterations

Central nervous system effects of EMFs may also be secondary to damage to the blood–brain barrier (BBB). The blood–brain barrier is a critical structure that prevents toxins and other large molecules that are in peripheral blood from having access to the brain matter itself. Salford et al. [118] have reported that a 2-h exposure of rats to GSM-900 radiation with a SAR of 2–200 mW/kg resulted in nerve cell damage. In a follow-up study, Eberhardt et al. report that 2-h exposures to cell phone GSM microwave RF resulted in leakage of albumin across the blood–brain barrier and neuronal death [119]. Neuronal albumin uptake was significantly correlated to occurrence of damaged neurons when measured at 28 days post-exposure. The lowest exposure level was 0.12 mW/kg (0.00012 W/kg) for 2 h. The highest exposure level was 120 mW/kg (0.12 W/kg). The weakest exposure level showed the greatest effect in opening the BBB [118]. Earlier blood–brain studies by Salford and Schirrmacher [120,121] report similar effects.

4. What are sources of wireless radiation?

There are many overlapping sources of radiofrequency and microwave emissions in daily life, both from industrial sources (like cell towers) and from personal items (cell and cordless phones, personal digital assistants (PDAs), wireless routers, etc.). Published data on typical levels found in some cities and from some sources are available at <http://www.bioinitiative.org> [1,122–124].

Cell phones are the single most important source of radiofrequency radiation to which we are exposed because of the relatively high exposure that results from the phone being held right against the head. Cell phones produce two types of emissions that should be considered. First, the radiofrequency radiation (typically microwave frequency radiation) is present. However, there is also the contribution of the switching battery pack that produces very high levels of extremely low frequency electromagnetic field [125–127].

Cordless telephones have not been widely recognized as similar in emissions to cell phones, but they can and do produce significant RF exposures. Since people tend to use them as substitutes for in-home and in-office corded or traditional telephones, they are often used for long periods of time. As the range of cordless phones has increased (the distance away that you can carry on a conversation is related to the power

output of the phone), the more powerful the RF signal will be. Hence, newer cordless phones may in some cases be similar to the power output of cell phones. The cumulative emissions from cell and cordless phones taken together should be recognized when considering the relative risks of wireless communication exposures.

PDAs such as the BlackBerry, Treo and iPhone units are 'souped-up' versions of the original voice communication devices (cell phones). The often produce far higher ELF emissions than do cell phones because they use energy from the battery very intensively for powering color displays and during data transmission functions (email, sending and receiving large files, photos, etc.) [125–127]. ELF emissions have been reported from PDAs at several tens to several hundreds of milligauss. Evidence of significantly elevated ELF fields during normal use of the PDA has public health relevance and has been reported in at least three scientific papers [125,128,129]. In the context of repetitive, chronic exposure to significantly elevated ELF pulses from PDAs worn on the body, relevant health studies point to a possible relationship between ELF exposure and cancer and pregnancy outcomes [130–133].

We include discussion of the ELF literature for two reasons. As mentioned above ELF activates the same biology as RF, it contributes to the total EMF burden of the body. In addition, PDAs and cell phones emit both radiofrequency/microwave radiation (RF) and extremely low frequency ELF from the battery switching of the device (the power source). Studies show that some devices produce excessively high ELF exposures during voice and data transmission. ELF is already classified as a 2B (Possible) Carcinogen by IARC, which means that ELF is indisputably an issue to consider in the wireless technology debate. ELF has been classified as a Group 2B carcinogen for all humans, not just children. The strongest evidence came from epidemiological studies on childhood leukemia, but the designation applies to all humans, both adults and children [1,25].

Wireless headsets that allow for conversations with cell phones at a distance from the head itself reduce the emissions. Depending on the type of wireless device, they may operate (transmit signal) only during conversations or they may be operational continuously. The cumulative dose of wireless headsets has not been well characterized under either form of use. Substantial cumulative RF exposure would be expected if the user wears a wireless headset that transmits a signal continuously during the day. However a critical factor is where the cell phone is placed. If worn on a belt with a headset, the exposure to the brain is reduced but the exposure to the pelvis may be significant.

Cell towers (called "masts" in Europe and Scandinavian countries) are wireless antenna facilities that transmit the cell phone signals within communities. They are another major source of RF exposures for the public. They differ from RF exposures from wireless devices like cell phones in that they produce much lower RF levels (generally 0.05 to 1–2 μ W/cm² in the first several hundred feet around them) in comparison to several hundred microwatts per centimeter

souped up iPhone

xc #3

squared for a cell phone held at the head. However they create a constant zone of elevated RF for up to 24 h per day, many hours per day, and the exposure is whole body rather than localized at the head. These facilities are the distribution system for wireless voice communications, internet connections and data transmission within communities. They are often erected on free-standing towers. They may be constructed on telephone poles or electrical poles. They may be built into the façade or rooftops of buildings behind wood screening. These are called stealth installations for wireless antenna facilities. Some installations are camouflaged to resemble 'false trees or rocks'. They emit RF to provide cell service to specific "cells" or locations that receive the signal.

Other forms of wireless transmission that are common in areas providing cell service are wireless land area networks (WLAN), (WiMAX) and WIFI networks. Some cities are installing city-wide WIFI service to allow any user on the street to log into the internet (without cables or wire connections). WIFI installations may have a signal reach for a few hundred feet where WiMAX installations may transmit signal more than 10 miles, so produce a stronger RF emission for those in close proximity. Each type has its particular signal strength and intended coverage area, but what they have in common is the production of continuous RF exposure for those within the area. We do not know what the cumulative exposure (dose) might be for people living, working or going to school in continuously elevated RF fields, nor are the possible health implications yet known. However, based on studies of populations near cell sites in general, there is a constellation of generally observed health symptoms that are reported to occur [85–107]. In this regard it is important to note that children living near to AM radio transmitters have been found to elevated risks of leukemia [134,135]. While AM radio RF fields are lower in frequency than that common in mobile phones, this is a total body irradiation with RF. The fact that leukemia, not brain cancer, is apparent in these studies suggests that leukemia is the cancer seen at the lowest levels of both ELF and RF fields under the circumstances of whole-body exposure.

Commercial surveillance systems or security gates pose an additional source of strong RF exposures. They are ubiquitous in department stores, markets and shops at the entry and exit points to discourage shoplifting and theft of goods. Security gates can produce excessively high RF exposures (although transitory) and have been associated with interference with pacemakers in heart patients. The exposure levels may approach thermal public safety limits in intensity, although no one expects a person to stand between the security gate bars for more than 6 min (safety limits for uncontrolled public access are variable depending on the frequency, but are all averaged over a 6-min exposure period).

RFID chips (radiofrequency identification chips) are being widely used to track purchases and for security of pets, and in some cases to keep track of patients with Alzheimer's disease and of children. RFID chips are implanted in fabrics, inserted in many types of commercial goods, and can be implanted

under the skin. They create a detectable signal to track the location of people and goods.

5. Problems with existing public health standards (safety limits)

If the existing standards were adequate none of the effects documented above should occur at levels to which people are regularly exposed. The fact that these effects are seen with our current ambient levels of exposure means that our existing public safety standards are obsolete. It also means that new, biologically based public exposure standards for wireless technologies are urgently needed. Whether it is feasible to achieve low enough levels that still work and also protect health against effects of chronic RF exposure – for all age groups – is uncertain. Whether we can protect the public and still allow the kinds of wireless technology uses we see today is unknown.

The nature of electromagnetic field interactions with biological systems has been well studied [136–144]. For purposes of standard-setting processes for both ELF and RF, the hypothesis that tissue damage can result only from heating is the fundamental flaw in the misguided efforts to understand the basic biological mechanisms leading to health effects.

The thermal standard is clearly untenable as a measure of dose when EMF stimuli that differ by many orders of magnitude in energy can stimulate the same biological response. In the ELF range, the same biological changes occur as in the RF, and no change in temperature can even be detected. With DNA interactions the same biological responses are stimulated in ELF and RF ranges even though the frequencies of the stimuli differ by many orders of magnitude. The effects of EMF on DNA to initiate the stress response or to cause molecular damage reflect the same biology in different frequency ranges. For this reason it should be possible to develop a scale based on DNA biology, and use it to define EMF dose in different parts of the EM spectrum. We also see a continuous scale in DNA experiments that focus on molecular damage where single and double strand breaks have long been known to occur in the ionizing range, and recent studies have shown similar effects in both ELF and RF ranges [144].

Existing standard-setting bodies that regulate wireless technologies, assume that there are no bioeffects of concern at exposure levels that do not cause measurable heating. However, it has been established beyond any reasonable doubt that bioeffects and some adverse health effects occur at far lower levels of RF and ELF exposure where no heating (or induced current) occurs; some effects are shown to occur a thousand times or more below the existing public safety limits. New, biologically based public exposure limits are urgently needed. New wireless technologies for cell and cordless phones, other wireless communication and data transmission systems affect living organisms in new ways that our antiquated safety limits have not foreseen, nor protected against.

The exposure of children to electromagnetic fields has not been studied extensively; in fact, the Federal Communications Commission (FCC) standards for exposure to radiofrequency radiation are based on the height, weight and stature of a 6-foot tall man, not scaled to children or adults of smaller stature. They do not take into account the unique susceptibility of growing children to exposures, nor are there studies of particular relevance to children.

In addition there is a problem in the consideration of the level of evidence taken into consideration by these bodies. There have not been adequate animal models shown to have cancer as an endpoint, and a perception that no single mechanism is proven to explain these associations. Thus these committees have tended to ignore or minimize the evidence for direct hazard to humans, and believe there is no proof of cause and effect. These bodies assume from the beginning that only conclusive scientific evidence (absolute proof) will be sufficient to warrant change, and refuse to take action on the basis of a growing body of evidence which provides early but consequential warning of risks.

The Radiofrequency Interagency Working Group of the US governmental agencies involved in RF matters (RFLAWG) issued a Guidelines Statement in June of 1999 that concluded the present RF standard "may not adequately protect the public" [145]. The RFLAWG identified fourteen (14) issues that they believe are needed in the planned revisions of ANSI/IEEE RF exposure guidelines including "to provide a strong and credible rationale to support RF exposure guidelines". In particular, the RFLAWG criticized the existing standards as not taking into account chronic, as opposed to acute exposures, modulated or pulsed radiation (digital or pulsed RF is proposed at this site), time-averaged measurements that may erase the unique characteristics of an intensity-modulated RF radiation that may be responsible for reported biologic effects, and stated the need for a comprehensive review of long-term, low-level exposure studies, neurological-behavioral effects and micronucleus assay studies (showing genetic damage from low-level RF) [145]. This important document from relevant US agencies questions existing standards in the following ways: (a) selection of an adverse effect level for chronic exposures not based on tissue heating and considering modulation effects; (b) recognition of different safety criteria for acute and chronic exposures at non-thermal or low-intensity levels; (c) recognition of deficiencies in using time-averaged measurements of RF that does not differentiate between intensity-modulated RF and continuous wave (CW) exposure, and *therefore may not adequately protect the public*; (d) having standards based on adult males rather than considering children to be the most vulnerable group.

6. Prudent public health responses

Emerging environmental health problems require preventative public health responses even where scientific and

medical uncertainties still exist, but where policy decisions today may greatly reduce human disease and societal costs tomorrow.

Policy decisions in public health must address some amount of uncertainty when balancing likely benefits and estimated costs. Although new insight will allow better appreciation of difficult issues, such as those occurring in environmental and occupational health, an expanded perspective may also enlarge the list of problems that need to be managed. Ignoring the problems carries its own costs (as deferring a decision is a decision in itself). With environmental and other public health problems becoming increasingly complex and international in scope, scientific documentation alone rarely justifies simple solutions [146].

Social issues regarding the controversy over public and occupational exposures to ELF and RF center on the resolute adherence to existing ICNIRP and FCC/IEEE standards by many countries, in the face of growing scientific evidence of health risks at far lower levels [10]. The composition of these committees, usually with excessive representation of the physics and engineering communities rather than public health professionals, results in a refusal to adopt biologically based exposure standards. Furthermore, there is widespread belief that governments are ignoring this evidence and there is widespread distrust of and lack of confidence in governments and their health agencies. The basis on which most review bodies and standard-setting agencies have avoided the conclusion that the science is strong enough to warrant new safety limits for ELF and RF is to require a demonstration of absolute proof before taking action. A causal level of evidence, or scientific certainty standard is implicit in nearly all reviews of the ELF and RF science, although this runs counter to good public health protection policies.

There is no question that global implementation of the safety standards proposed in the Bioinitiative Report, if implemented abruptly and without careful planning, have the potential to not only be very expensive but also disruptive of life and the economy as we know it. Action must be a balance of risk to cost to benefit. The major risk from maintaining the status quo is an increasing number of cancer cases, especially in young people, as well as neurobehavioral problems at increasing frequencies. The benefits of the status quo are expansion and continued development of communication technologies. But we suspect that the true costs of even existing technologies will only become much more apparent with time. Whether the costs of remedial action are worth the societal benefits is a formula that should reward precautionary behavior. Prudent corporate policies should be expected to address and avoid future risks and liabilities, otherwise, there is no market incentive to produce safe (and safer) products.

The deployment of new technologies is running ahead of any reasonable estimation of possible health impacts and estimates of probabilities, let alone a solid assessment of risk. However, what has been missing with regard to EMF has been an acknowledgement of the risk that is demonstrated by

the scientific studies. There is clear evidence of risk, although the magnitude of the risk is uncertain, and the magnitude of doing nothing on the health effects cost to society is similarly uncertain. This situation is very similar to our history of dealing with the hazards of smoking decades ago, where the power of the industry to influence governments and even conflicts of interest within the public health community delayed action for more than a generation, with consequent loss of life and enormous extra health care costs to society. New standards are warranted now, based on the totality of scientific evidence; the risks of taking no-action, the large population at risk, costs associated with ignoring the problem in new and upgraded site selection and construction, and the loss of public trust by ignoring the problem.

Direct medical and rehabilitative health costs associated with treatment for diseases that are reasonably related to wireless technologies may be very large. Although there is uncertainty involved in how much disease is related to wireless exposures, the mere scale of the problem with several billion users of cell phones and even larger impacts on bystander populations (from cell site exposures, from other WI-FI and wireless exposures in-home and commercial use, etc.) the associated public health costs will likely be monumental. Furthermore the costs to families with cancers, neurological diseases or learning disabilities in children related in part or in whole to wireless technologies extend beyond medical costs. They may reasonably extend to family disruption and family psychological problems, losses in job productivity and income loss.

The history of governments and their official health agencies to deal with emerging and newly identified risks to health is not good [147–149]. This is particularly true where industry investments in new products and technologies occur without full recognition, disclosure or even knowledge of possible health consequences. Large economic investments in polluting industries often make for perilously slow regulatory action, and the public health consequences may be very great as a result [150,151].

Free markets do not internalize the costs to society of “guessing wrong”: Unexpected or hidden health costs of new technologies may not be seen for many years; when the ability to recall or to identify the precise exposures related to disease outcomes is difficult or impossible. The penalty nearly always falls to the individual, the family or the taxpayer and not to the industry that benefits economically,—at least in free-market economies. Thus, the profits go to industry but the costs may go to the individual who can suffer both diminished quality of life and health and economic disadvantage. If all disease endpoints that may be reasonably related to chronic exposure to electromagnetic fields are considered even a small attributable fraction for one or more industries, it will have enormous global impact on public health. The public health implications are immense. But they can be reduced by strong government and public health interventions providing information on alternatives to wireless technologies, public education campaigns, health advisories,

Table 1

Public health implications of wireless technologies argue for change in governmental and health agency actions.

Secure US and EU legislative mandates for safer technologies for communication and data transmission, for security and surveillance needs.
Promote wired alternatives for voice and data communication (cable, fiber-optic)
Discourage or ban use of cell phones by children and young teenagers
Provide permanent (unremovable) labels on cell phones “Not for use by children under the age of 16”
Implement national public education campaigns on health issues (cell phones, cordless phones, PDAs, wireless internet, city-wide WI-FI, WLAN and WIMAX exposures)
Promote industry redesign for safer products: support innovation for alternatives and solutions
Slow or stop deployment of wireless technologies to discourage reliance on wireless technologies for communication and security needs
Put the burden of proof on industry to show “new wireless tech” is safe before deployment
Adopt and enforce restricted use areas for sensitive or more vulnerable segments of society including low-EMF environments in public areas and “No Cell” zones in airports, hospitals, schools
Acknowledge FCC and ICNIRP thermal safety standards are obsolete for wireless technologies
Appoint new standard-setting bodies familiar with biological effects to develop new guidelines for public safety limits.
Develop new biologically based standards that address low-intensity, chronic exposures
Require standard of evidence and level of proof = public health
Reject “causal” standard of evidence for taking action on science
Make industry financially liable for “guessing wrong” and ignoring health risks

requirements for redesign of wireless devices, proscription of use of wireless devices by children and teenagers; strong and independent research programs on causes and prevention of EMF-related diseases, and consultation with all stakeholders on issues relating to involuntary exposures (bystander or second-hand radiation exposures from wireless technologies) (Table 1).

The scientific information contained in this Supplement argues for thresholds or guidelines that are substantially below current FCC and ICNIRP standards for localized exposures to wireless devices and for whole-body exposure. Uncertainty about how low such standards might have to go to be prudent from a public health standpoint should not prevent reasonable efforts to respond to the information at hand. No lower limit for bioeffects and adverse health effects from RF has been established, so the possible health risks of wireless WLAN and WI-FI systems, for example, will require further research. No assertion of safety at any level of wireless exposure (chronic exposure) can be made at this time. The lower limit for reported human health effects has dropped 100-fold below the safety standard (for mobile phones and PDAs); 1000–10,000-fold for other wireless (cell towers at distance; WI-FI and WLAN devices). The entire basis for safety standards is called into question, and it is not unreasonable to question the safety of RF at any level.

It is likely that for both ELF and RF, as for other carcinogens, there is no threshold of exposure that is without risk, but the magnitude of the risk increases linearly with the level of exposure. Our society will not go back to the pre-electric and pre-wireless age, but the clear evidence of health hazards to the human population from exposure mandates that we develop ways in which to reduce exposure through education, new technologies and the establishment of biomedically based standards.

7. Conclusions and recommended actions

New ELF limits are warranted based on a public health analysis of the overall existing scientific evidence. These limits should reflect environmental levels of ELF that have been demonstrated to increase risk for childhood leukemia, and possibly other cancers and neurological diseases. ELF limits should be set below those exposure levels that have been linked in childhood leukemia studies to increased risk of disease, plus an additional safety factor. It is no longer acceptable to build new power lines and electrical facilities that place people in ELF environments that have been determined to be risky. These levels are in the 2–4 milligauss (mG) range (0.2–0.4 μT), not in the 10 s of mG or 100 s of mG. The existing ICNIRP limit is 1000 mG (100 μT) and 904 mG (90.4 μT) in the US for ELF is outdated and based on faulty assumptions. These limits are can no longer be said to be protective of public health and they should be replaced. A safety buffer or safety factor should also be applied to a new, biologically based ELF limit, and the conventional approach is to add a safety factor lower than the risk level.

While new ELF limits are being developed and implemented, a reasonable approach would be a 1 mG (0.1 μT) planning limit for habitable space adjacent to all new or upgraded power lines and a 2 mG (0.2 μT) limit for all other new construction. It is also recommended that a 1 mG (0.1 μT) limit be established for existing habitable space for children and/or women who are pregnant (because of the possible link between childhood leukemia and *in utero* exposure to ELF). This recommendation is based on the assumption that a higher burden of protection is required for children who cannot protect themselves, and who are at risk for childhood leukemia at rates that are traditionally high enough to trigger regulatory action. This situation in particular warrants extending the 1 mG (0.1 μT) limit to existing occupied space. "Establish" in this case probably means formal public advisories from relevant health agencies. While it is not realistic to reconstruct all existing electrical distribution systems, in the short-term, steps to reduce exposure from these existing systems need to be initiated, especially in places where children spend time, and should be encouraged. These limits should reflect the exposures that are commonly associated with increased risk of childhood leukemia (in the 2–5 mG (0.2–0.5 μT) range for all children, and over 1.4 mG (0.14 μT) for children age 6 and younger). Nearly all of

the occupational studies for adult cancers and neurological diseases report their highest exposure category is 4 mG (0.4 μT) and above, so that new ELF limits should target the exposure ranges of interest, and not necessarily higher ranges.

Avoiding chronic ELF exposure in schools, homes and the workplace above levels associated with increased risk of disease will also avoid most of the possible bioactive parameters of ELF discussed in the relevant literature.

It is not prudent public health policy to wait any longer to adopt new public safety limits for ELF. These limits should reflect the exposures that are commonly associated with increased risk of childhood leukemia (in the 2–5 mG (0.2–0.5 μT) range for all children, and over 1.4 mG (0.14 μT) for children age 6 and younger). Avoiding chronic ELF exposure in schools, homes and the workplace above levels associated with increased risk of disease will also avoid most of the possible bioactive parameters of ELF discussed in the relevant literature.

The rapid deployment of new wireless technologies that chronically expose people to pulsed RF at levels reported to cause bioeffects, which in turn, could reasonably be presumed to lead to serious health impacts, is a public health concern. There is suggestive to strongly suggestive evidence that RF exposures may cause changes in cell membrane function, cell communication, metabolism, activation of proto-oncogenes and can trigger the production of stress proteins at exposure levels below current regulatory limits. Resulting effects can include DNA breaks and chromosome aberrations, cell death including death of brain neurons, increased free-radical production, activation of the endogenous opioid system, cell stress and premature aging, changes in brain function including memory loss, retarded learning, performance impairment in children, headaches and fatigue, sleep disorders, neurodegenerative conditions, reduction in melatonin secretion and cancers (BioInitiative Report Chapters 5–10, 12) [1].

This information now argues for thresholds or guidelines that are substantially below current FCC and ICNIRP standards for whole-body exposure. Uncertainty about how low such standards might have to go to be prudent from a public health standpoint should not prevent reasonable efforts to respond to the information at hand. No lower limit for bioeffects and adverse health effects from RF has been established, so the possible health risks of wireless WLAN and WI-FI systems, for example, will require further research and no assertion of safety at any level of wireless exposure (chronic exposure) can be made at this time. The lower limit for reported human health effects has dropped 100-fold below the safety standard (for mobile phones and PDAs); 1000–10,000-fold for other wireless (cell towers at distance; WI-FI and WLAN devices). The entire basis for safety standards is called into question, and it is not unreasonable to question the safety of RF at any level.

A cautionary target level for pulsed RF exposures for ambient wireless that could be applied to RF sources from cell tower antennas, WI-FI, WI-MAX and other similar sources

is proposed. The recommended cautionary target level is 0.1 microwatts per centimeter squared ($\mu\text{W}/\text{cm}^2$) (or 0.614 V per meter or V/m) for pulsed RF where these exposures affect the general public; this advisory is proportionate to the evidence and in accord with prudent public health policy. A precautionary limit of 0.1 $\mu\text{W}/\text{cm}^2$ should be adopted for outdoor, cumulative RF exposure. This reflects the current RF science and prudent public health response that would reasonably be set for pulsed RF (ambient) exposures where people live, work and go to school. This level of RF is experienced as whole-body exposure, and can be a chronic exposure where there is wireless coverage present for voice and data transmission for cell phones, pagers and PDAs and other sources of radiofrequency radiation. An outdoor precautionary limit of 0.1 $\mu\text{W}/\text{cm}^2$ would mean an even lower exposure level inside buildings, perhaps as low as 0.01 $\mu\text{W}/\text{cm}^2$. Some studies and many anecdotal reports on ill health have been reported at lower levels than this; however, for the present time, it could prevent some of the most disproportionate burdens placed on the public nearest to such installations. Although this RF target level does not preclude further rollout of WI-FI technologies, we also recommend that wired alternatives to WI-FI be implemented, particularly in schools and libraries so that children are not subjected to elevated RF levels until more is understood about possible health impacts. This recommendation should be seen as an interim precautionary limit that is intended to guide preventative actions; and more conservative limits may be needed in the future.

Broadcast facilities that chronically expose nearby residents to elevated RF levels from AM, FM and television antenna transmission are also of public health concern given the potential for very high RF exposures near these facilities (antenna farms). RF levels can be in the 10s to several 100s of $\mu\text{W}/\text{cm}^2$ in residential areas within half a mile of some broadcast sites (for example, Lookout Mountain, Colorado and Awbrey Butte, Bend, Oregon). Like wireless communication facilities, RF emissions from broadcast facilities that are located in, or expose residential populations and schools to elevated levels of RF will very likely need to be re-evaluated for safety.

For emissions from wireless devices (cell phones; personal digital assistant or PDA devices, etc.) there is enough evidence for increased risk of brain tumors and acoustic neuromas now to warrant intervention with respect to their use. Redesign of cell phones and PDAs could prevent direct head and eye exposure, for example, by designing new units so that they work only with a wired headset or on speakerphone mode.

These effects can reasonably be presumed to result in adverse health effects and disease with chronic and uncontrolled exposures, and children may be particularly vulnerable. The young are also largely unable to remove themselves from such environments. Second-hand radiation, like second-hand smoke is an issue of public health concern based on the evidence at hand.

In summary, the following recommendations are made:

- ELF limits should be set below those exposure levels that have been linked in childhood leukemia studies to increased risk of disease, plus an additional safety factor. It is no longer acceptable to build new power lines and electrical facilities that place people in ELF environments that have been determined to be risky (at levels generally at 2 mG (0.2 μT) and above).
- While new ELF limits are being developed and implemented, a reasonable approach would be a 1 mG (0.1 μT) planning limit for habitable space adjacent to all new or upgraded power lines and a 2 mG (0.2 μT) limit for all other new construction. It is also recommended for that a 1 mG (0.1 μT) limit be established for existing habitable space for children and/or women who are pregnant. This recommendation is based on the assumption that a higher burden of protection is required for children who cannot protect themselves, and who are at risk for childhood leukemia at rates that are traditionally high enough to trigger regulatory action. This situation in particular warrants extending the 1 mG (0.1 μT) limit to existing occupied space. "Establish", in this case probably means formal public advisories from relevant health agencies.
- While it is not realistic to reconstruct all existing electrical distributions systems, in the short-term; steps to reduce exposure from these existing systems need to be initiated and should be encouraged, especially in places where children spend time.
- A precautionary limit of 0.1 $\mu\text{W}/\text{cm}^2$ (which is also 0.614 V per meter) should be adopted for outdoor, cumulative RF exposure. This reflects the current RF science and prudent public health response that would reasonably be set for pulsed RF (ambient) exposures where people live, work and go to school. This level of RF is experienced as whole-body exposure, and can be a chronic exposure where there is wireless coverage present for voice and data transmission for cell phones, pagers and PDAs and other sources of radiofrequency radiation. Some studies and many anecdotal reports on ill health have been reported at lower levels than this; however, for the present time, it could prevent some of the most disproportionate burdens placed on the public nearest to such installations. Although this RF target level does not preclude further rollout of WI-FI technologies, we also recommend that wired alternatives to WI-FI be implemented, particularly in schools and libraries so that children are not subjected to elevated RF levels until more is understood about possible health impacts. This recommendation should be seen as an interim precautionary limit that is intended to guide preventative actions; and more conservative limits may be needed in the future.

References

- [1] C. Sage, D.O. Carpenter (Eds.), *BioInitiative Working Group BioInitiative Report: A Rationale for a Biologically-based Public*

- [39] J.B. Burch, J.S. Reif, C.W. Noonan, T. Ichinose, A.M. Bachand, T.L. Koleber, M.G. Yost, Melatonin metabolite excretion among cellular telephone users, *Int. J. Rad. Biol.* 78 (2002) 1029–1036.
- [40] Girgert, et al., Induction of tamoxifen resistance in breast cancer cells by ELF electromagnetic fields, *Biochem. Biophys. Res. Commun.* 336 (2005) 1144–1149.
- [41] Harland, et al., Environmental magnetic fields inhibit the antiproliferative action of tamoxifen and melatonin in a human breast cancer cell line, *Bioelectromagnetics* 18 (1997) 555–562.
- [42] Harland, et al., Evidence for a slow time-scale of interaction for magnetic fields inhibiting tamoxifen's antiproliferative action in human breast cancer cells, *Cell Biochem. Biophys.* 31 (3) (1999) 295–306.
- [43] A. Ilhan, A. Gurel, F. Armutcu, S. Kamisli, M. Iraz, O. Akyol, S. Ozen, Ginkgo biloba prevents mobile phone-induced oxidative stress in rat brain, *Clin. Chim. Acta* 340 (1–2) (2004) 153–162.
- [44] Ishido, et al., Magnetic fields (MF) of 50 Hz at 1.2 μ T as well as 100 μ T cause uncoupling of inhibitory pathways of adenylyl cyclase mediated by melatonin 1a receptor in MF-sensitive MCF-7 cells, *Carcinogenesis* 22 (7) (2001) 1043–1048.
- [45] H. Koylu, H. Mollaoglu, F.N. Ozguner, Melatonin modulates 900 Mhz microwave-induced lipid peroxidation changes in rat brain, *Toxicol. Ind. Health* 22 (5) (2006) 211–216.
- [46] R.P. Liburdy, T.R. Sloma, et al., ELF magnetic fields, breast cancer, and melatonin: 60 Hz fields block melatonin's oncostatic action on ER+ breast cancer cell proliferation, *J. Pineal Res.* 14 (1993) 89–97.
- [47] R.P. Liburdy, et al., Magnetic Fields, melatonin, tamoxifen and human breast cancer cell growth, in: R.G. Stevens, B.W. Wilson, L.E. Anderson (Eds.), *The Melatonin Hypothesis—Breast Cancer and Use of Electric Power*, Battelle Press, Columbus, Richland, 1997, pp. 669–700.
- [48] L.I. Loberg, Gene expression in human breast epithelial cells exposed to 60 Hz magnetic fields, *Carcinogenesis* 20 (1999) 1633–1636.
- [49] Luben, et al., Replication of 12 mG EMF effects on melatonin responses of MCF-7 breast cancer cells in vitro, in: Abstract A-1 of the 1996 Annual Review of Research on Biological Effects of Electric and Magnetic Fields from the Generation, Delivery and Use of Electricity, San Antonio, TX, November 17–21, 1996, p. 1.
- [50] Luben, et al., Independent replication of 60-Hz 1.2 μ T EMF effects on melatonin and tamoxifen responses of MCF-7 cells in vitro, in: Abstract A-3.4, *Bioelectromagnetics Society Annual Meeting*, St. Pete Beach, FL, June 7–11, 1998, pp. 17–18.
- [51] Morris, In vitro exposure of MCF-7 human breast cancer cells to 60-Hz magnetic fields, in: Abstract p-125A, *Bioelectromagnetics Society Annual Meeting*, St. Pete Beach, FL, June 7–11, 1998, pp. 204–205.
- [52] F. Oktem, F. Ozguner, H. Mollaoglu, A. Koyu, E. Uz, Oxidative damage in the kidney induced by 900-MHz-emitted mobile phone: protection by melatonin, *Arch. Med. Res.* 36 (4) (2005) 350–355.
- [53] F. Ozguner, G. Aydin, H. Mollaoglu, O. Gokalp, A. Koyu, G. Cesur, Prevention of mobile phone induced skin tissue changes by melatonin in rat: an experimental study, *Toxicol. Ind. Health* 20 (6–10) (2004) 133–139.
- [54] F. Ozguner, A. Altunbas, M. Ozaydin, A. Dogan, H. Vural, A.N. Kisioglu, G. Cesur, N.G. Yikirim, Mobile phone-induced myocardial oxidative stress: protection by a novel antioxidant agent caffeic acid phenethyl ester, *Toxicol. Ind. Health* 21 (9) (2005) 223–230.
- [55] A.W. Guy, C.K. Chou, L.L. Kunz, J. Crowley, J. Krupp, Effects of long-term low-level radiofrequency radiation exposure on rats. US Air Force School of Aerospace Medicine Brooks Air Force Base, Texas TR-85-64 Final Report August 1985, Approved for public release: distribution is unlimited.
- [56] C.K. Chou, Long-term low level microwave irradiation of rats, *Bioelectromagnetics* 13 (1992) 469–496.
- [57] H. Dotk, et al., Cancer incidence near radio and television transmitters in Great Britain, *Am. J. Epidemiol.* 145 (1) (1997) 1–9.
- [58] V. Garaj-Vrhovac, D. Horvat, Z. Koren, The relationship between colony-forming ability, chromosome aberrations and incidence of micronuclei in V79 Chinese hamster cells exposed to microwave radiation, *Mutat. Res.* 263 (3) (1991) 143–149.
- [59] V. Garaj-Vrhovac, A. Fucic, D. Horvat, The correlation between the frequency of micronuclei and specific chromosome aberrations in human lymphocytes exposed to microwave radiation in vitro, *Mutat. Res.* 281 (3) (1992) 181–186.
- [60] V. Garaj-Vrhovac, Micronucleus assay and lymphocyte mitotic activity in risk assessment of occupational exposure to microwave radiation, *Chemosphere* 39 (13) (1999) 2301–2312.
- [61] M. Ha, H. Im, M. Lee, H.J. Kim, B.C. Kim, Y.M. Gimmi, et al., Radio-frequency radiation exposure from AM radio transmitters and childhood leukemia and brain cancer, *Am. J. Epidemiol.* 166 (2007) 270–279.
- [62] B. Hocking, et al., Cancer incidence and mortality and proximity to TV towers, *Med. J. Aust.* 165 (11–12) (1996) 601–605.
- [63] B. Hocking, Decreased survival for childhood leukemia in proximity to TV towers, in: Poster Presented at the Annual Scientific Meeting of the Royal Australian College of Physicians in Adelaide, SA, Australia, May, 2000.
- [64] D.E. Foliart, B.H. Pollock, G. Mezell, R. Iriye, J.M. Silva, K.L. Epi, L. Kheifets, M.P. Lind, R. Kavet, Magnetic field exposure and long-term survival among children with leukemia, *Brit. J. Cancer* 94 (2006) 161–164.
- [65] A. Huss, A. Spoerri, M. Egger, Rööslü for the Swiss national cohort study, residence near power lines and mortality from neurodegenerative diseases: longitudinal study of the Swiss population, *Am. J. Epidemiol.* (November) (2008) (Epub ahead of print).
- [66] F. Marinelli, D. La Sala, G. Ciccioiti, L. Cattini, C. Trimarchi, S. Putti, A. Zamparelli, L. Giuliani, G. Tomassetti, C. Cinti, Exposure to 900 MHz electromagnetic field induces an unbalance between pro-apoptotic and pro-survival signals in T-lymphoblastoid leukemia CCRF-CEM cells, *J. Cell Physiol.* 198 (2) (2004) 324–332.
- [67] P. Michelozzi, C. Ancona, D. Fusco, F. Forastiere, C.A. Perucci, Risk of leukemia and residence near a radio transmitter in Italy, *Epidemiology* 9 (Suppl.) (1998) 354.
- [68] F. Repacholi, et al., Lymphomas in E μ -Pim1 transgenic mice exposed to pulsed 900 MHz electromagnetic fields, *Radiat. Res.* 147 (1997) 31–40.
- [69] A. Stang, et al., The possible role of radiofrequency radiation in the development of uveal melanoma, *Epidemiology* 12 (1) (2001) 7–12.
- [70] M. Blank (2007), Section 7, pp. 1–40, Evidence for Stress Response (Stress Proteins). In *BioInitiative Report: A Scientific Perspective on Health Risk of Electromagnetic Fields*. Published Online 31 August 2007. <http://www.bioinitiative.org/report/index.htm>.
- [71] C. Daniels, I. Duce, D. Thomas, P. Sewell, J. Tattersall, D. de Pomerai, Transgenic nematodes as biomonitors of microwave-induced stress, *Mutat. Res.* 399 (1998) 55–64.
- [72] D. de Pomerai, et al., Non-thermal heat-shock response to microwaves, *Nature* 405 (2000) 417–418.
- [73] S. Kwee, et al., The biological effects of microwave radiation, in: *Proceedings of the Second World Congress for Electricity and Magnetism in Biology and Medicine*, Bologna, Italy, June, 1997.
- [74] S. Kwee, et al., Changes in cellular proteins due to environmental non-ionizing radiation. I. Heat-shock proteins, *Electro Magnetobiol.* 20 (2001) 141–152.
- [75] D. Leszczynski, S. Oenaväri, J. Reivinen, R. Kuokka, Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: molecular mechanism for cancer- and blood-brain barrier-related effects, *Differentiation* 70 (2002) 120–129.
- [76] D. Leszczynski, R. Nylund, S. Joenvaara, J. Reivinen, Applicability of discovery science approach to determine biological effects of mobile phone radiation, *Proteomics* 4 (2) (2004) 426–431.
- [77] S. Lixia, Y. Ke, W. Kaijun, L. Dequiang, H. Huajun, G. Xiangwei, W. Baohong, Z. Wei, L. Jianling, W. Wei, Effects of 1.8 GHz

- Exposure Standard for Electromagnetic Fields (ELF and RF), 2007. <http://www.bioinitiative.org>.
- [2] REFLEX Program. Risk evaluation of potential environmental hazards from low frequency electromagnetic field exposure using sensitive *in vitro* methods. A project funded by the European Union under the programme Quality of Life and Management of Living Resources, Key Action 4 (2004).
 - [3] L. Hardell, C. Sage, Biological effect from electromagnetic field exposure and public exposure standards, *Biomed. Pharmacother.* 62 (2008) 104–109, doi:10.1016/j.biopha. (2007)12.004.
 - [4] L. Hardell, M. Carlberg, P. Söderqvist, K. Hansson Mild, Meta-analysis of long-term mobile phone use and the association with brain tumours, *Int. J. Oncol.* 32 (2008) 1097–1103.
 - [5] P. Kan, S.E. Simonsen, J.L. Lyon, J.R.W. Kestle, Cellular phone use and brain tumor: a meta-analysis, *J. Neurooncol.* (2007), doi:10.1007/s11060-007-9432-1.
 - [6] E. Cardis, Interphone Study Memo, International Agency for Cancer Research, October 2008.
 - [7] S. Sadetzki, A. Chetrit, A. Jarus-Hakak, E. Cardis, Y. Deutch, S. Dvenderani, A. Zultan, I. Novikov, L. Freedman, M. Wolf, Cellular phone use and risk of benign and malignant parotid gland tumors—a nationwide case-control study, *Am. J. Epidemiol.* (2008), doi:10.1093/aje/kwn325.
 - [8] A. Lahkola, T. Salminen, J. Raitanen, S. Heimavaara, M.J. Schoemaker, H.C. Christensen, M. Feychting, C. Johansen, L. Klæboe, S. Lonn, A.J. Swerdlow, T. Tynes, A. Auvinen, Meningioma and mobile phone use—a collaborative case-control study in five North European countries, *Int. J. Epidemiol. Adv. Access* (August) (2008), doi:10.1093/ije/dyn155.
 - [9] T. Takebayashi, S. Akiba, Y. Kikuchi, et al., Mobile phone use and acoustic neuroma risk in Japan, *Occup. Environ. Med.* 63 (2007) 802–807.
 - [10] D.O. Carpenter, C.L. Sage, Setting prudent public health policy for electromagnetic field exposures, *Rev. Environ. Health* 23 (2) (2008) 91–117.
 - [11] D.O. Carpenter, C.L. Sage, BioInitiative Working Group, Key scientific evidence and public health policy options, BioInitiative Report at <http://www.bioinitiative.org>, 17 (2007).
 - [12] European Commission, Health and Consumer Protection, Scientific Committee on SCENIHR Report on Emerging and Newly Identified Health Risks—Possible Effects of Electromagnetic Fields (EMF) on Human Health, 2007.
 - [13] World Health Organization, Extremely Low Frequency Fields Environmental Health Criteria Monograph 238, 2007. <http://www.who.int/peh-erf/project/en> and <http://www.who.int/peh-erf/meetings/elf.emf.workshop.2007/en/index.html>.
 - [14] <http://news.cnet.com/Emerging-markets-fuel-cell-phone-growth/2100-1039-3-6159491.html>.
 - [15] <http://news.foxpda.com/news/2-14-billion-cell-phone-subscribers-in-2005-2120.shtml>.
 - [16] http://www.ctia.org/media/industry_info/index.cfm/AID/10323.
 - [17] <http://www.nytimes.com/2008/09/20/us/20messaging.html> and Members Info, Electronics Industries Alliance Website, November, 2000; Members Info, Electronics Industries Alliance Website, November 2000.
 - [18] L. Hardell, M. Carlberg, K. Hansson Mild, Pooled analysis of two case-control studies on use of cellular and cordless telephones and the risk for malignant brain tumours diagnosed in 1997–2003, *Int. Arch. Environ. Health* 79 (2006) 630–639, doi:10.1007/s00420-006-0088-5.
 - [19] M. Kundi, The controversy about a possible relationship between mobile phone use and cancer, *Environ. Health Persp.* (September) (2008), doi: 10.1289/ehp.11902.
 - [20] J. Pronczuk-Garbino (Ed.), *Children's Health and the Environment: A Global Perspective*, World Health Organization, Geneva, Switzerland, 2005, p. 367.
 - [21] World Health Organization, *Children's Health and Environment: A Review of Evidence: A Joint Report from the European Environmental Agency and The World Health Organization*, 2002. <http://www.who.int/peh-cmf>.
 - [22] O.P. Gandhi, G. Lazzi, C.M. Furse, Electromagnetic absorption in the human head and neck for mobile telephones at 835 and 1900 MHz, *IEEE Trans. Microw. Theory Tech.* 44 (1996) 1884–1896.
 - [23] O.P. Gandhi, G. Kang, Some present problems and a proposed experimental phantom for SAR compliance testing of cellular telephones at 835 and 1900 MHz, *Phys. Med. Biol.* 47 (2002) 1501–1518.
 - [24] J. Wiart, A. Hadjem, M.F. Wong, I. Bloch, Analysis of RF exposure in the head tissues of children and adults, *Phys. Med. Biol.* 53 (2008) 3681–3695; J. Wiart, A. Hadjem, M.F. Wong, I. Bloch, Analysis of RF exposure in the head tissues of children and adults, *Wiart Phys. Med. Biol.* 53 (June) (2008) 2771–2783.
 - [25] M.A. Divan, L. Kheifets, C. Obel, J. Olsen, Prenatal and postnatal exposure to cell phone use and behavioral problems in children, *Epidemiology* 19 (4) (2008).
 - [26] International Commission on Non-ionizing Radiation Protection, Guidelines for limiting exposure to time-varying electric, magnetic and electromagnetic fields (up to 300 GHz), *Health Phys.* 74(4) (1998) 494–522. <http://www.icnirp.de>.
 - [27] Institute of Electrical and Electronics Engineers, Inc (IEEE), Section 4.2 of "IEEE Standard for Safety Levels with Respect to Human Exposure to Radio Frequency Electromagnetic Fields, 3 kHz to 300 GHz," ANSI/IEEE C95.1-1992. New York, NY, 1992.
 - [28] H. Lai, N.P. Singh, Single and double strand DNA breaks in rat brain cells after acute exposure to radiofrequency electromagnetic radiation, *Int. J. Radiat. Biol.* 69 (1996) 513–521.
 - [29] S. Ivancsis, A. Pilger, E. Diem, O. Jahn, H.W. Rudinger, Cell type-specific genotoxic effects of intermittent extremely-low frequency electromagnetic fields, *Mutat. Res.* 583 (2005) 184–188.
 - [30] J. Phillips, et al., DNA damage in multi-4 lymphoblastoid cells exposed to cellular telephone radiofrequency fields *in vitro*, *Bioelectrochem. Bioenerg.* 45 (1998) 103–110.
 - [31] R.J. Aitken, L.E. Bennetts, D. Sawyer, A.M. Wiklundt, B.V. King, Impact of radio frequency electromagnetic radiation on DNA integrity in the male germline, *Int. J. Androl.* 28 (2005) 171.
 - [32] J.Y. Kim, S.Y. Hong, Y.M. Lee, S.A. Yu, W.S. Koh, J.R. Hong, T. Son, S.K. Chang, M. Lee, *In vitro* assessment of clastogenicity of mobile-phone radiation (835 MHz) using the alkaline comet assay and chromosomal aberration test, *Environ. Toxicol.* 23 (2008) 319.
 - [33] S. Lixia, K. Yao, W. Kaijun, I. Deglang, H. Huanjun, G. Xiangwei, W. Baohong, Z. Wei, L. Jianling, W. Wei, Effects of 1.8 GHz radiofrequency field on DNA damage and expression of heat shock protein 70 in human lens epithelial cells, *Mutat. Res.* 602 (2006) 135.
 - [34] R. Paulraj, J. Behari, Single strand DNA breaks in rat brain cells exposed to microwave radiation, *Mutat. Res.* 596 (2006) 76.
 - [35] M. Mashevich, D. Folkman, A. Kesar, A. Barbul, R. Korenstein, E. Jerby, L. Avivi, Exposure of human peripheral blood lymphocytes to electromagnetic fields associated with cellular phones leads to chromosomal instability, *Bioelectromagnetics* 24 (2003) 82.
 - [36] I.G. Akoev, M.S. Pashovkina, L.P. Dolgacheva, T.P. Semenova, V.L. Kalmykov, Enzymatic activity of some tissues and blood serum from animals and humans exposed to microwaves and hypothesis on the possible role of free radical processes in the nonlinear effects and modification of emotional behavior of animals, *Radiat. Biol. Radioecol.* 42 (3) (2002) 32–330.
 - [37] C. Blackman, et al., The influence of 1.2 μ T, 60 Hz magnetic fields on melatonin and tamoxifen-induced inhibition of MCF-7 cell growth, *Bioelectromagnetics* 22 (2) (2001) 122–128.
 - [38] D.E. Blask, S.M. Hill, Effects of melatonin on cancer: studies on MCF-7 human breast cancer cells in culture, *J. Neural Transm. Suppl.* 21 (1986) 433–449.

- radiofrequency field on DNA damage and expression of heat shock protein 70 in human lens epithelial cells, *Mutat. Res.* (2006). doi:10.1016/j.mrfmmm.2006.08.010.
- [78] M. Simko, C. Hartwig, M. Lantow, M. Lupke, M.O. Mattsson, Q. Rahman, J. Rollwitz, Hsp 70 expression and free radical release after exposure to non-thermal radiofrequency electromagnetic fields and ultrafine particles in human Mono Mac 6 cells, *Toxicol. Lett.* 161 (2006) 73–82 (Elsevier Science Direct).
- [79] S. Velizarov, The effects of radiofrequency fields on cell proliferation are non-thermal, *Bioelectrochem. Bioenerg.* 48 (1999) 177–180.
- [80] D. Weisbrodt, H. Lin, L. Ye, M. Blank, R. Goodman, Effects of mobile phone radiation on reproduction and development in *Drosophila melanogaster*, *J. Cell Biochem.* 89 (1) (2003) 48–55.
- [81] T. Olivares-Banuelos, L. Navarro, A. Gonzalez, R. Drucker-Colin, Differentiation of chromaffin cells elicited by ELF MF modifies gene expression pattern, *Cell Biol. Int.* 28 (2004) 273–279.
- [82] R. Zhao, S. Zhang, Z. Xu, L. Ju, D. Lu, G. Yao, Studying gene expression profile of rat neuron exposed to 1800 MHz radiofrequency electromagnetic fields with cDNA microarray, *Toxicology* 235 (2007) 167–175.
- [83] A. Karinen, S. Hejnavara, R. Nylund, D. Leszczynski, Mobile phone radiation might alter protein expression in human skin, *BMC Genomics* 9 (2008) 77, doi:10.1186/1471-2164-9-77.
- [84] J. Vanderstraeten, L. Verschaeve, Gene and protein expression following exposure to radiofrequency fields from mobile phones, *Environ. Health Persp.* 116 (2008) 1131–1135.
- [85] G. Abdel-Rassoul, O.A. El-Fateh, M.A. Salem, A. Michael, F. Farahat, M. El-Batanouny, E. Salem, Neurobehavioral effects among inhabitants around mobile phone base stations, *Neurotoxicology* 28 (2007) 434–440.
- [86] P. Achermann, Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep EEG, *Neuroreport* 11 (15) (2000) 3321–3325.
- [87] E.S. Altpeter, T.H. Krebs, Study on health effects of the shortwave transmitter station of Schwarzenburg, Bern, Switzerland, University of Bern BEW Publications Study No. 56, The Federal Office of Energy, 1995.
- [88] A.A. Borbely, et al., Pulsed high-frequency electromagnetic field affects human sleep and sleep electroencephalogram, *Neurosci. Lett.* 275 (3) (1999) 207–210.
- [89] R. Huber, T. Graf, K.A. Cote, L. Wittmann, E. Gallmann, D. Matter, J. Schuderer, N. Kuster, A.A. Borbely, P. Achermann, Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep EEG, *Neuroreport* 11 (15) (2000) 3321–3325.
- [90] K. Mann, Effects of pulsed high-frequency electromagnetic fields on human sleep, *Neuropsychobiology* 33 (1996) 41–47.
- [91] G. Oberfeld, The microwave syndrome—further aspects of a Spanish study, in: *Proceedings of the Third International Workshop on Bioelectromagnetic Effects of Electromagnetic Fields*, Kos, Greece, 2004.
- [92] R. Santini, M. Seigne, L. Bonhomme-Faivre, S. Bouffet, E. Defrasne, M. Sage, Symptoms experienced by users of digital cellular phones: a pilot study in a French engineering school, *Pathol. Biol. (Paris)* 49 (3) (2001) 222–226.
- [93] R. Santini, P. Santini, P. Le Ruz, J.M. Danze, M. Seigne, Survey study of people living in the vicinity of cellular phone base stations, *Electromag. Biol. Med.* 22 (2003) 41–49.
- [94] TNO Physics and Electronics Laboratory, The Netherlands, Effects of Global Communication System Radio-frequency Fields on Well-being and Cognitive Functions of Human Beings With and Without Subjective Complaints, Netherlands Organization for Applied Scientific Research (2003), pp. 1–63.
- [95] E.S. Altpeter, T.H. Krebs, Study on health effects of the shortwave transmitter station of Schwarzenburg, Bern, Switzerland, University of Bern BEW Publications Study No. 56, The Federal Office of Energy, Switzerland, 1995.
- [96] S.E. Chia, Prevalence of headache among handheld cellular telephone users in Singapore: a community study, *Environ. Health Persp.* 108 (11) (2000) 1059–1062.
- [97] H. Chiang, et al., Health effects of environmental electromagnetic fields, *J. Bioelectr.* 8 (1989) 127–131.
- [98] H. D'Costa, et al., Human brain wave activity during exposure to radiofrequency field emissions from mobile phones, *Aust. Phys. Eng. Sci. Med.* 26 (4) (2003).
- [99] H.P. Hutter, H. Moshhammer, P. Wallner, M. Kundi, Subjective symptoms, sleeping problems and cognitive performance in subjects living near mobile phone base stations, *Occup. Env. Med.* 63 (2006) 307–313.
- [100] M. Koivisto, et al., Effects of 902 MHz electromagnetic field emitted by cellular telephones on response times in humans, *Neuroreport* 11 (2000) 413–415.
- [101] M. Koivisto, et al., The effects of electromagnetic field emitted by GSM phones on working memory, *Neuroreport* 11 (2002) 1641–1643.
- [102] A.A. Kolodynski, V.V. Kolodynska, Motor and psychological functions of school children living in the area of the Skrunda radio location station in Latvia, *Sci. Total Environ.* 180 (1996) 87–93.
- [103] C.M. Krause, L. Sillanmaki, M. Koivisto, A. Haggqvist, C. Saarela, A. Revonsuo, M. Laine, H. Hamalainen, Effects of electromagnetic field emitted by cellular phones on the EEO during a memory task, *Neuroreport* 11 (4) (2000) 761–764.
- [104] C.M. Krause, L. Sillanmaki, M. Koivisto, A. Haggqvist, C. Saarela, A. Revonsuo, M. Laine, H. Hamalainen, Effects of electromagnetic fields emitted by cellular phones on the electroencephalogram during a visual working memory task, *Int. J. Radiat. Biol.* 76 (12) (2000) 1659–1667.
- [105] J. Lass, et al., Effects of 7 Hz-modulated 450 MHz electromagnetic radiation on human performance in visual memory tasks, *Int. J. Radiat. Biol.* 73 (10) (2002) 937–944.
- [106] A.A. Marino, E. Nilsen, C. Frilot, Nonlinear changes in brain electrical activity due to cell phone radiation, *Bioelectromagnetics* 24 (5) (2003) 339–346.
- [107] G. Oberfeld, et al., The microwave syndrome—further aspects of a Spanish study, in: *Proceedings of the Third International Workshop on Bioelectromagnetic Effects of Electromagnetic Fields*, Kos, Greece, 2004.
- [108] A. Preece, et al., Effect of a 915-MHz simulated mobile phone signal on c function in man, *Int. J. Radiat. Biol.* 75 (1999) 447–456.
- [109] S.J. Regel, S. Negovetic, M. Roosli, V. Berdinas, J. Schuderer, A. Huss, U. Lott, N. Kuster, P. Achermann, UMTS base station-like exposure, well-being and cognitive performance, *Environ. Health Persp.* 114 (2006) 1270–1275.
- [110] S. Eliti, D. Wallace, A. Ridgewell, K. Zougkou, R. Russo, F. Sepulveda, D. Mirshakar-Syahkal, P. Rasor, R. Deebie, E. Fox, Does short-term exposure to mobile phone base station signals increase symptoms in individuals who report sensitivity to electromagnetic fields? A double-blind randomized provocation study, *Environ. Health Persp.* 115 (2007) 1603–1608.
- [111] E.F. Pace-Schott, J.A. Hobson, The neurobiology of sleep: genetics, cellular physiology and subcortical networks, *Nat. Rev. Neurosci.* 3 (2002) 591–605.
- [112] J.B. Burch, J.S. Reif, C.W. Noonan, T. Ichinose, A.M. Bachand, T.L. Koleher, M.G. Yost, Melatonin metabolite excretion among cellular telephone users, *Int. J. Radiat. Biol.* 78 (2002) 1029–1036.
- [113] M.L. Clark, J.B. Burch, M.G. Yost, Y. Zhai, A.M. Bachand, C.T. Fitzpatrick, J. Ramaprasad, L.A. Cragin, J.S. Reif, Biomonitoring of estrogen and melatonin metabolites among women residing near radio and television broadcasting transmitters, *J. Occup. Environ. Med.* 49 (2007) 1149–1156.
- [114] L.M. Wang, N.A. Suthana, D. Chaudhry, D.R. Weaver, C.S. Colwell, Melatonin inhibits hippocampal long-term potentiation, *Eur. J. Neurosci.* 22 (2005) 2231–2237.

- [115] M. Ozcan, B. Yilmaz, D.O. Carpenter, Effects of melatonin on synaptic transmission and long term potentiation in two areas of mouse hippocampus, *Brain Res.* 1111 (2006) 90–94.
- [116] R.J. Reiter, L. Tang, J.J. Garcia, A. Munoz-Hoyos, Pharmacological actions of melatonin in oxygen radical pathophysiology, *Life Sci.* 60 (1997) 2255–2271.
- [117] C.W. Noonan, J.S. Reif, J.B. Burch, T.Y. Ichinose, M.G. Yost, K. Magnusson, Relationship between amyloid beta protein and melatonin metabolite in a study of electric utility workers, *J. Occup. Environ. Med.* 44 (2002) 769–775.
- [118] L.G. Salford, A.E. Brun, et al., Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones, *Environ. Health Persp.* 111 (2003) 881–883.
- [119] J.L. Eberhardt, B.R.R. Persson, A.E. Brun, L.G. Salford, L.O.G. Malmgren, Blood–brain barrier permeability and nerve cell damage in rat brain 14 and 28 days after exposure to microwaves from GSM mobile phones, *Electromagn. Biol. Med.* 27 (2008) 215–229. doi:10.1080.1536370802344037.
- [120] L.G. Salford, et al., Permeability of the blood brain barrier induced by 915 MHz electromagnetic radiation continuous wave and modulated at 8, 16, 50 and 200 Hz, *Microsc. Res. Tech.* 27 (1994) 535–542.
- [121] A. Schirmacher, Electromagnetic fields (1.8 GHz) increase the permeability of sucrose of the blood–brain barrier in vitro, *Bioelectromagnetics* 21 (2000) 338–345.
- [122] Y. Hamnerius, Microwave exposure from mobile phones and base stations in Sweden, in: *Proceedings of the International Conference on Cell Tower Siting*, sponsored by the University of Vienna & Land Salzburg, Salzburg, Austria, June 7–8, 2000.
- [123] E.D. Mantiply, Summary of measured radiofrequency electric and magnetic fields (10 kHz–30 GHz) in the general and work environment, *Bioelectromagnetics* 18 (1997) 563–577.
- [124] Sage Associates, 2004. An Overview of Low-intensity radiofrequency/microwave radiation studies relevant to wireless communications and data, *Bioelectromagnetics Society Annual Meeting*, Washington DC, June, 2004.
- [125] C. Sage, O. Johansson, S.A. Sage, Personal digital assistant (PDA) cell phone units produce elevated extremely-low frequency electromagnetic field emissions, *Bioelectromagnetics* 28 (7) (2007) 581–582.
- [126] C. Sage, O. Johansson, Response to comment on personal digital assistant (PDA) cell phone units produce elevated extremely-low frequency electromagnetic field emissions, *Bioelectromagnetics* (July) (2007), 17654541.
- [127] C. Sage, O. Johansson, Response to comment on measuring ELF fields produced by mobile phones and personal digital assistants (PDAs), *Bioelectromagnetics* (July) (2007), 17654536.
- [128] M. Tuor, S. Ebert, J. Schuderer, N. Kuster, Assessment of ELF magnetic fields from five mobile handsets, in: *ITIS Foundation, Conference Presentation*, Monte Verita, 2005.
- [129] T. Linde, K.H. Mild, Measurement of low frequency magnetic fields from digital cellular telephones, *Bioelectromagnetics* 18 (1997) 184–186 (Brief communication).
- [130] B. Armstrong, G. Theriault, P. Guenel, J. Deadman, M. Goldberg, P. Heroux, Association between exposure to pulsed electromagnetic fields and cancer in electrical utility workers in Ontario and Quebec, Canada, and France 1970–1989, *Am. J. Epidemiol.* 140 (1994) 805–820.
- [131] G.M. Lee, R.R. Neutra, L. Hristova, M. Yost, R.A. Hiatt, A nested case–control study of residential and personal magnetic field measures and miscarriages, *Epidemiology* 13 (1) (2002) 21–31.
- [132] K.D. Li, R. Oudouli, S. Wi, T. Janevic, I. Golditch, T.D. Bracken, R. Senior, R. Rankin, R. Iriye, A population-based prospective cohort study of personal exposure to magnetic fields during pregnancy and the risk of spontaneous abortion, *Epidemiology* 13 (2002) 9–20.
- [133] G. Theriault, M. Goldberg, A.B. Miller, B. Armstrong, P. Guenel, J. Deadman, E. Imbemon, T. To, A. Chevalier, D. Cyr, C. Wall, Cancer risks associated with occupational exposure to magnetic fields among utility workers in Ontario and Quebec, Canada and France: 1970–1989, *Am. J. Epidemiol.* 139 (1994) 550–572.
- [134] P. Michelozzi, A. Capon, U. Kirchmayer, F. Forastiere, A. Biggeri, A. Barca, C.A. Perucci, Adult and childhood leukemia near a high-power radio station in Rome, Italy, *Am. J. Epidemiol.* 155 (2002) 1096–1103.
- [135] M. Ha, H. Im, M. Lee, H.J. Kim, B.C. Kim, Y.M. Gimm, J.K. Park, Radio-frequency radiation exposure from AM radio transmitters and childhood leukemia and brain cancer, *Am. J. Epidemiol.* 166 (2007) 270–279.
- [136] C. Sage, D.O. Carpenter (Eds.), *BioInitiative Working Group BioInitiative Report: A Rationale for a Biologically-based Public Exposure Standard for Electromagnetic Fields (ELF and RF)*, 2007. www.bioinitiative.org.; C. Blackman, H. Lai, Chapters 6 and 14 of the BioInitiative Report, 2007.
- [137] A. Chiabrera, B. Bianco, E. Moggia, J.J. Kaufman, *Bioelectromagnetics* 21 (4) (2000) 312–324.
- [138] D.J. Panagopoulos, L.H. Margaritis, in: A.C. Harper, R.V. Bures (Eds.), *Mobile Telephony Radiation Effects on Living Organisms*, 2008, pp. 107–149. ISBN 978:1-60456-436-5.
- [139] W.R. Adey, in: P.J. Rosch, M.S. Markov (Eds.), *Potential Therapeutic Applications of Nonthermal Electromagnetic Fields: Ensemble Organization of Cells in Tissue as a Factor in Biological Field Sensing*, *Bioelectromagnetic Medicine*, 2004, pp. 1–16.
- [140] S. Engstrom, in: P.J. Rosch, M.S. Markov (Eds.), *Magnetic Field Generation and Dosimetry*, *Bioelectromagnetic Medicine*, 2004, pp. 39–50.
- [141] A. Pilla, in: F.S. Barnes, B. Greenebaum, (Eds.), *Mechanisms and Therapeutic Applications of Time-varying and Static Magnetic Fields in Biological and Medical Aspects of Electromagnetic Fields*, third edition, 2007, pp. 351–412.
- [142] M. Blank, R. Goodman, Initial interactions in electromagnetic field-induced biosynthesis, *J. Cell. Physiol.* 199 (2004) 359–363.
- [143] M. Blank, R. Goodman, A biological guide for electromagnetic safety: the stress response electromagnetic initiation of transcription at specific DNA sites, *Bioelectromagnetics* 25 (2004) 642–646.
- [144] M. Blank, *BEMS Soc. Newslett.* (January–February) (2008) 6–7.
- [145] G. Lotz, Letter from Greg Lotz, PhD, Chief Physical Agents Effects Branch, Division of Biomedical and Behavioral Science, National Institute of Occupational Safety and Health to Richard Tell, Chair, IEEE SCC28 (SC4) Risk Assessment Work Group, June 17, 1999.
- [146] P. Grandjean, Implications of the precautionary principle for primary prevention and research, *Am. Rev. Public Health* 25 (2004) 199–223.
- [147] D. Davis, *The Secret History of the War on Cancer*, Basic Books, 2008.
- [148] R. Proctor, *Cancer Wars*, Harper Collins Publishers, 1995.
- [149] European Environmental Agency, *Late Lessons from Early Warnings. The Precautionary Principle 1896–2000*. Copenhagen, Denmark, 2001.
- [150] P. Landrigan, How much do chemicals affect our health? *Discover Mag.* (2008).
- [151] California Air Resources Board, Appendix III Proposed Identification of Environmental Tobacco Smoke as a Toxic Air Contaminant. Part B—Health Effects, 2005.

The London Resolution

Olle Johansson*

The Experimental Dermatology Unit, Karolinska Institute, Department of Neuroscience, Stockholm, Sweden

At a scientific conference on 27th November 2007 entitled—“Are Present ICNIRP EMF Exposure Recommendations Adequate?”, hosted by Roger Coghill and Robert Verkerk, at the Royal Society, London, scientists endorsed the BioInitiative Report, extended the 2006 Benevento Resolution and resolved that:

“We, the undersigned, do call on the UK Health Protection Agency (HPA), UK Government and all the health protection agencies and governments world-wide, to take note of the findings and recommendations in the BioInitiative Report (2007) [1] and its predecessors the Benevento Resolution (2006) [2], the Catania Resolution (2002) [3] and the Salzburg Resolution (2000) [4] to immediately reduce the guidelines for exposure to radiofrequency radiation (RF) and extremely low-frequency electromagnetic¹ fields (ELF-EMF) for the following reasons:

- The overwhelming evidence of adverse non-thermal health effects at exposures many times below the current guidelines.
- The near 100% penetration of the market in Europe, the USA and many other markets by mobile phones and increasing penetration elsewhere.
- The vast proliferation of wireless networks and devices beyond those envisaged at the time the current guidelines were set.

We call for the ICNIRP to reconvene as a matter of urgency to reassess the exposure guidelines and to develop and implement biologically based public safety limits reflecting the overall scientific evidence that existing ICNIRP guidelines are not sufficiently protective against health effects from chronic exposures to the rapidly increasing environmental-level ELF-EMF and RF...

* Tel.: +46 8 52487073; fax: +46 8 303904.

E-mail address: olle.johansson@ki.se.

¹ Magnetic fields at 50–60 Hz.

Failing that:

- We call for the setting up of an independent body to define new biologically based public exposure limits and/or preventative actions, for ELF-EMF and RF, that address reported biological effects, which, with prolonged exposure, can reasonably be presumed to result in adverse health consequences.
- In the absence of such recommendations we suggest as an intermediate step that the HPA and UK Government immediately implement the ELF-EMF and RF recommendations of the BioInitiative Report 2007 and strive for the recommendations of the Public Health Department of the Government of Salzburg (2002) of 0.06 V/m for outdoor and 0.02 V/m for indoor RF exposure.

Based on the precautionary principle, children and vulnerable groups (such as people with epilepsy and heart conditions) should not be exposed to a risk of harm, thus we propose that

- Children under 16 should use mobile phones and cordless phones for emergency calls only.
- No Wi-fi, WiMax or other forms of wireless networking are placed in homes, schools or public areas or promoted for use thereof.
- That regular and frequent independent audits are undertaken of emissions to ensure that base stations (“masts”) do not exceed the new biologically based guidelines at any locality either singly or by accumulation. Such audits should be widely publicised and made available for public scrutiny.

The precautionary principle needs to be implemented.”

Signed:

Prof. Christopher Busby, School of Biomedical Sciences,
Faculty of Life and Health Sciences, University of Ulsters
Coleraine, UK

Mr. Roger Coghill, MA (Cantab) C Biol Mi Biol MA (Environ Mgt), Coghill Research Laboratories, Pontypool, Wales, UK

Dr. Andrew Goldsworthy, Honorary Lecturer, Imperial College, London, UK

Dr. Lennart Hardell, University Hospital, Orebro University, Orebro, Sweden

Prof. Olle Johansson, Experimental Dermatology Unit, Department of Neuroscience, Karolinska Institute, and The Royal Institute of Technology, Stockholm, Sweden

Dr. Gerd Oberfeld, Public Health Department, Salzburg Government, Salzburg, Austria

Mr. Alasdair Philips, B.Sc. (Eng), Director, Powerwatch, Sutton in the Isle, UK

Mr. Graham Philips, MBCS, Technical Manager, Powerwatch, Ely, UK

Ms Cindy Sage, Co-Editor, BioInitiative Report, Santa Barbara CA, USA

Dr. John Walker, Chartered Physicist, Sutton Coldfield, UK

References

- [1] BioInitiative Report, <http://www.bioinitiative.org/report/docs/report.pdf>.
- [2] Benevento Resolution, *Electromagnetic Biology and Medicine*, 25, 2006, pp. 197–200, <http://www.icems.eu/docs/BeneventoResolution.pdf>.
- [3] Catania Resolution, http://www.plattform-mobilfunk-initiativen.at/englisch/CATANIA_eng.pdf.
- [4] Salzburg Resolution, http://www.salzburg.gv.at/salzburg_resolution_e.pdf.